



Contents lists available at ScienceDirect

## Journal of Archaeological Science: Reports

journal homepage: [www.elsevier.com/locate/jasrep](http://www.elsevier.com/locate/jasrep)

## Capturing Roman dietary variability in the catastrophic death assemblage at Herculaneum

R.E.V. Martyn<sup>a,\*</sup>, P. Garnsey<sup>b</sup>, L. Fattore<sup>c</sup>, P. Petrone<sup>d</sup>, A. Sperduti<sup>e</sup>, L. Bondioli<sup>e</sup>, O.E. Craig<sup>a</sup>

<sup>a</sup> Department of Archaeology, BioArCh, University of York, York YO10 5DD, UK

<sup>b</sup> Faculty of History, University of Cambridge, Cambridge CB3 9EF, UK

<sup>c</sup> Dipartimento Ingegneria Chimica Materiali Ambiente, Università di Roma 'La Sapienza', Rome, Italy

<sup>d</sup> Dipartimento di Scienze Biomediche Avanzate, Sezione di Medicina Legale, Istologia e Anatomia, Università degli Studi di Napoli Federico II, Naples, Italy

<sup>e</sup> Museo delle Civiltà, Servizio di Bioarcheologia, Rome, Italy

### ARTICLE INFO

#### Keywords:

Herculaneum  
Stable isotopes  
Palaeodiet  
Vesuvius

### ABSTRACT

Here we present a comparative study of stable carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) isotope data from 81 individuals from the catastrophic death assemblage at Herculaneum (79 CE) and compare these with the attritional sites of Velia (Salerno, Italy, 1st–2nd century CE) and Isola Sacra (Rome, Italy, 1st–2nd century AD). The instantaneous deposition of the Herculaneum assemblage highlights some interesting differences in our contextual and methodological understanding of stable dietary isotopes, suggesting that isotopic variation between sites may sometimes be a result of greater temporal variability rather than truly comparable differences. Our results suggest that the people of Herculaneum obtained a relatively small proportion (ca. 30%) of their dietary carbon from marine foods; the majority originating from terrestrial foodstuffs of a similar carbon isotopic composition, most likely cereals. Also observed is a generally greater dietary isotopic enrichment in male individuals than females. We infer that males had greater access to fish which may be reflective, in part, of the sociodemographic framework characteristic of Roman society. Finally, we highlight the methodological challenges which may be faced when undertaking comparisons of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  data between the various age-related strata of a population, particularly due to the slow and variable rate of collagen turnover.

### 1. Introduction

The health and economic ‘well-being’ of the Roman world is a fundamental benchmark in the historic investigation of past civilisations. Although the study of the Roman productive economy is extensive, our knowledge regarding the distribution of wealth and differences in living conditions in Roman society is limited to partial and incomplete records (Garnsey and Saller, 2015). We do not yet know how food was distributed to different elements of the population, between households, villages or towns. Historical accounts (Rackham, 1967; Edwards, 2001; Wolf, 2010) and archaeological evidence from animal and plant remains (Meyer, 1980; Pagano, 1994; Reese, 2002; Rowan, 2014; Robinson and Rowan, 2015) provide specific information regarding the types of foods that were eaten but lack the resolution required to quantify dietary content, or to study dietary variability within societies. Such information is crucial if we are to make meaningful comparisons between Roman and other pre-modern and developing societies, and to clarify relationships between social status, health

and nutrition.

Stable carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) isotope analysis of bone collagen offers a direct approach to the inter- and intra-population study of ancient diet. Isotopic signals represent a direct measure of an individual's average dietary intake during the period of bone collagen formation. These analyses are particularly useful for discriminating diets of coastal inhabitants with access to mixed marine and terrestrial diets, and where the major dietary sources (e.g. marine fish, terrestrial herbivores, terrestrial omnivores and cereal grains) have distinct isotope values. So far, the analyses of over 500 individuals from Roman Imperial period necropolises in southern Italy have succeeded in identifying relative isotopic differences within and between assemblages, attributed to differences in occupation, age and sex, and mainly relating to the differential consumption of marine foods (Prowse et al., 2004; Craig et al., 2009; Killgrove and Tykot, 2013; Killgrove and Tykot, in press). Nevertheless, the analysis of diet in such attritional death assemblages is heavily burdened by methodological and interpretative limitations. Unlike census data, skeletal assemblages from burial

\* Corresponding author.

E-mail addresses: [revm500@york.ac.uk](mailto:revm500@york.ac.uk) (R.E.V. Martyn), [pdag1@cam.ac.uk](mailto:pdag1@cam.ac.uk) (P. Garnsey), [luciano.fattore@uniroma1.it](mailto:luciano.fattore@uniroma1.it) (L. Fattore), [pipetron@unina.it](mailto:pipetron@unina.it) (P. Petrone), [alessandra.sperduti@beniculturali.it](mailto:alessandra.sperduti@beniculturali.it) (A. Sperduti), [luca.bondioli@beniculturali.it](mailto:luca.bondioli@beniculturali.it) (L. Bondioli), [oliver.craig@york.ac.uk](mailto:oliver.craig@york.ac.uk) (O.E. Craig).

<http://dx.doi.org/10.1016/j.jasrep.2017.08.008>

Received 31 March 2017; Received in revised form 1 August 2017; Accepted 7 August 2017  
2352-409X/© 2017 Elsevier Ltd. All rights reserved.

grounds are palimpsests that gradually accumulate over time, and their fidelity to any living population is undermined by both selective burial and selective mortality (Wood et al., 1992; Roberts and Grauer, 2001; Jackes, 2011; DeWitte and Stojanowski, 2015). For example, individuals who were afforded cremation, a common Roman funerary custom, cannot be studied, whilst frail individuals who succumbed to disease are likely to be over-represented in the younger age classes (Wood et al., 1992).

In studying stable isotopic data from a sample of 81 individuals from the catastrophic death assemblage at Herculaneum (Bisel, 1991; Capasso and Domenicantonio, 1998; Capasso and Capasso, 1999; Capasso, 2000; Mastrolorenzo et al., 2001; Mastrolorenzo et al., 2010; Petrone, 2011), we hope to circumvent these problems and derive a clearer picture of dietary variability in at least one Roman town. All were victims of the 79 CE eruption of Vesuvius and were discovered within 9 *fornci* (stone vaults) running adjacent to the seafront (Fattore et al., 2012). The stable isotope data for 72 individuals were originally reported in Craig et al. (2013) but here we investigate these data with respect to new osteological information regarding the age and sex of the skeletons. Notably, this revision identified one of the 72 individuals (F8I10) as a juvenile. In addition, we also report new isotopic data from 9 infants and juveniles (< 20 years of age). Albeit a modest sample of a small Imperial coastal town of ca. 4–5000 residents (Wallace-Hadrill, 2011), the assemblage contains a broadly equal mixture of adult males and females, with juveniles and infants also represented (Capasso, 2000; Mastrolorenzo et al., 2001). Whilst some selectivity in those sheltering in the vaults is to be expected, the assemblage offers a rare glimpse of contemporary Roman life, where sudden and collective death negated the selective biases usually faced in osteoarchaeological analysis. Therefore, we are able for the first time to quantify the differential access to foods within an ancient ‘living’ population.

## 2. Methods

Collagen for the new 9 samples was extracted from bone and analysed by EA-IRMS exactly as described previously (Craig et al., 2013). In the majority of for both these samples and those presented in Craig et al. (2013), rib samples were chosen (Craig et al., 2013; see Supporting information, Table 1) and any samples showing signs of pathological change were excluded. Briefly, bone samples (0.5–1 g) were coarsely ground and demineralised (0.6 M HCl, 4 °C, 3–12 days), samples were rinsed with distilled water and then gelatinised (pH 3 [0.001 M] HCl, 80 °C, 48 h). The supernatant containing the collagen was filtered (30 kDa, Amicon® Ultra-4 Centrifugal Filter Units, Millipore, Billerica, MA, USA), frozen, and lyophilised. Collagen samples (1 mg) were analysed in duplicate or triplicate by EA/IRMS in a Sercon GSL analyser coupled to a Sercon 20–22 Mass Spectrometer (Sercon, Crewe, UK) at the University of York, or a Roboprep Combustion Device coupled to a Europa 20–20 Mass Spectrometer (PDZ-Europa, Crewe, UK). The analytical error, calculated from repeated measurements of each sample and measurements of the bovine control from multiple extracts, was < 0.2‰ (1σ). Accuracy was determined by measurements of international standard reference materials (IAEA 600, IAEA N2, IA Cane) within each analytical run, with the error being less than < 0.5‰ in all instances. The difference in the <sup>15</sup>N/<sup>14</sup>N ratio between the sample and the internationally defined standard AIR (atmospheric air) in ‰ units is referred to as δ<sup>15</sup>N, and δ<sup>13</sup>C refers to the difference in <sup>13</sup>C/<sup>12</sup>C ratio between the sample and the internationally defined standard, PDB (Vienna Pee Dee Belemnite Limestone). The reported ratios are calculated using the equation:

$$\delta x = ((R_{\text{sample}} - R_{\text{standard}})/R_{\text{standard}}) \times 1000$$

For Herculaneum, the <sup>14</sup>C offset attributable to the marine reservoir effect was estimated for each sample using the following regression equation derived from radiocarbon dating and stable isotope analysis of 9 samples (Craig et al., 2013):

$$y = 34.3 - 300x, R^2 = 9.1 \text{ where } y = {}^{14}\text{C offset (years) and } x = \delta^{15}\text{N value (‰)} \quad (1)$$

These 9 individuals are a sub-sample of the 81 individuals analysed for δ<sup>13</sup>C and δ<sup>15</sup>N in the current study.

The calculated <sup>14</sup>C offset from the above equation was used to estimate the % of total carbon derived from a marine source, assuming a maximum reservoir age of 390 years corresponding to 100% marine derived carbon. The % of marine protein contribution to collagen was derived through linear interpolation of values between the terrestrial endpoint (+ 7.2‰) and marine endpoint (+ 16‰). The latter were derived from measurements of contemporary herbivore and marine fish values, using similar assumptions as previously reported (Craig et al., 2013). All statistical analysis was carried out using R version 3.1.2.

The human osteological material was analysed according to the common standards reported in the literature (Krogman and İşcan, 1986; Buikstra and Ubelaker, 1994; White and Folkens, 2005). Sex determination in the adults was obtained by the application of the visual assessment of the morphological traits of skull and pelvis (Ferembach et al., 1980; White and Folkens, 2005). Age at death was determined using multiple age indicators. For adult individuals, methods included: degenerative changes of the pubic symphysis (Todd, 1921), the auricular surface of the innominate (Buikstra and Ubelaker, 1994), and the sternal ends of ribs (İşcan et al., 1984); ecto- and endo-cranial suture closure (Buikstra and Ubelaker, 1994). For individuals still growing at the time of death the following criteria were applied: stages of epiphyseal fusion (Scheuer et al., 2010), long bone dimensions (Scheuer et al., 2010), and the stages of formation and eruption of teeth (AlQahtani et al., 2010). The analyses were independently performed by three observers (PP, LF, AS) and cases of discrepancy were resolved by a fourth joint and consensual analysis (on the reliability of the age-at-death assessment see (Baccino et al., 1999; Garvin and Passalacqua, 2012). The extraordinary preservation state of the skeletal and dental material allowed for the age at death to be determined by 5 year intervals for subadults and 10 year intervals for adult individuals (the last age class being 50 +), thus permitting comparison with almost contemporaneous central Italian skeletal series (Prowse et al., 2004; Prowse et al., 2005; FitzGerald et al., 2006; Craig et al., 2009; Crowe et al., 2010; Petrone et al., 2011).

The Herculaneum sample set reported in this paper is composed of 81 individuals: 28 females, 37 males, 6 unsexed individuals older than 15 years and 10 individuals (< 15 years) which were unsexed, see Supplementary information, Table 1.

For the dietary reconstruction, we included the biological sub-adults (age 15–20, 5 males, 2 females, and 4 unsexed) within the analysis of the adult individuals on the grounds that they probably ate an adult diet, being classed ‘social’ adults in accordance with the trend of traditional Roman life (Treggiari, 1993).

## 3. Results and discussion

### 3.1. Dietary variation at Herculaneum and other coastal roman sites

The carbon and nitrogen stable isotope data for the Herculaneum population are reported in Supporting information, Table 1. These include all the data reported in Craig et al. (2013) plus those from an additional 9 infants and juveniles. Overall, the isotope data for all individuals > 15 years fall within the range of similar age cohorts from other coastal Imperial necropolises (Fig. 2). These are Isola Sacra (Prowse et al., 2004; Crowe et al., 2010), the cemetery that served Portus Romae - the gateway to Rome, and Velia - a small coastal town south of Naples (Craig et al., 2009) (Fig. 1). The δ<sup>13</sup>C values at each of the three sites have comparable ranges (Herculaneum = − 18.2‰ to − 20.2‰; Isola Sacra = − 17.8‰ to − 19.5‰; Velia = − 18.7‰ to − 20.0‰) but the variances are significantly different between sites (Fligner-Killeen test of homogeneity of variances;  $\chi^2 = 6.8, p = 0.03$ ).



Fig. 1. Map showing approximate locations of Italian Roman Imperial period sites referred to in the text (after Craig et al., 2013).

It is noticeable, however, that the  $\delta^{15}\text{N}$  values for Herculaneum show a narrower range (8.2‰ to 11.7‰) than for Isola Sacra (7.5‰ to 15.3‰) or Velia (6.4‰ to 14.1‰), despite similar sample sizes (Velia = 117; Isola Sacra = 94; Herculaneum = 71). Conversely, the variances within samples are not significantly different (Fligner-Killeen test of homogeneity of variances;  $\chi^2 = 3.4$ ,  $p = 0.18$ ). The bagplots (Fig. 2 (Rousseuw et al., 1999)) clearly show that Herculaneum has an “intermediate” position between the two other coastal sites both for  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$ , and a much narrower distribution of  $\delta^{15}\text{N}$  values. One explanation for the relatively reduced dietary variation at Herculaneum compared to Isola Sacra and Velia is the nature of the assemblage formations. As the latter are individuals from cemeteries that were used for many generations (ca. 150 years for Velia and ca. 300 years for Isola Sacra), greater isotopic variation may simply reflect greater dietary variation through time, rather than real differences in the diet of the living populations, as is commonly assumed when such comparisons are made.

To test for inter-site differences in  $\delta^{15}\text{N}$ , a robust ANOVA model was used. As diet is significantly affected by sex in each of these assemblages (see Section 3.2.), it was particularly important to examine whether differences in the demographic profiles are a more likely explanation for the amplitude of isotopic variation between sites. The  $\delta^{15}\text{N}$  values are significantly different by site ( $F = 129.4$ ,  $p \leq 0.001$ ) as expected but not when the interaction between sex and site is considered ( $F = 0.1$ ,  $p = 0.89$ ). Therefore, the distribution of  $\delta^{15}\text{N}$  values genuinely reflects greater dietary variation at the attritional assemblages, compared to Herculaneum. Interestingly the core distributions, containing 25% to 75% of the  $\delta^{15}\text{N}$  data (Fig. 2), at each assemblage are

comparable in terms of amplitude of variance. The main difference between the sites is that Velia and Isola Sacra have a greater number of outliers, particularly individuals with high marine protein diets (i.e. high  $\delta^{15}\text{N}$  values).

Finally, the amplitude of variance in  $\delta^{15}\text{N}$  between the sites is not easily explained by greater absolute differences in dietary end-points (plants and fish) as discussed previously (Craig et al., 2009), although temporal variation in these, particularly changes in location of grain supply, would be interesting to check. The consumption of leguminous vegetables, thought to be integral to the Roman diet (Garnsey, 1999) and with ample evidence from Vesuvian cities (Meyer, 1980; Wolf, 2010), should also be explored. These may have a large effect on the isotopic endpoints since they are relatively depleted in  $^{15}\text{N}$ . Finally, the presence in the Velia assemblage of a specific subset of individuals, possibly fishermen, has been observed (Crowe et al., 2010) and contributes to the broad range of  $\delta^{15}\text{N}$  for this site.

### 3.2. Variation by sex

The distribution of  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  is significantly different between the sexes at Herculaneum (Kruskal-Wallis,  $\chi^2 = 4.6$ ,  $p = 0.03$  and  $\chi^2 = 5.1$ ,  $p = 0.02$  for each isotope, respectively) with males typically enriched in  $^{15}\text{N}$  and  $^{13}\text{C}$  compared to females (Fig. 3a). From these data, it is proposed that males consumed more fish with relatively elevated  $\delta^{15}\text{N}$  values. This is not to suggest that other low trophic level species were not consumed at Herculaneum, either fresh or as commodities such as garum. Indeed there are ample remains of small fish such as sardine, anchovy, and marine shellfish from sewer deposits (Rowan,




Fig. 2. Bagplot of three Roman Imperial period mortuary assemblages. Comparison of human stable isotope data between Velia (left) and Isola Sacra (middle) by means of bagplots. A bagplot is a bivariate generalization of the boxplot. The central darker shaded area contains 50% of all data points. The outer lighter shaded area is three times the area of the central part and is fenced by a line connecting data points that lie on the periphery of this area. Points outside the fence are considered outliers. Medians are represented with a gold star. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

2014) but these are less likely to be distinguished isotopically.

At Herculaneum, since all the individuals died simultaneously (Mastrolorenzo et al., 2001), we can exploit differences in individual radiocarbon dates to independently quantify marine food consumption with much more certainty. At this site it has been previously shown that both carbon and nitrogen isotopes in human bone collagen are positively linearly correlated with the amount of ‘old’ carbon derived from the marine reservoir (Craig et al., 2013). On this basis, it is estimated that across the Herculaneum sample a relatively small proportion (0–30%) of the total carbon in bone collagen, broadly equivalent to the weight % or calorific contribution to the diet, was derived from marine foods (Craig et al., 2013). Given their richer protein content, marine foods make a much greater contribution to total dietary protein (nitrogen) which at Herculaneum is estimated to range between 20 and 50% (Craig et al., 2013). These estimates are also supported by the application of a Bayesian mixing model, which takes into account the macronutrient composition of different food groups (Fernandes, 2015).

Apart from fish, the remainder of the diet at Herculaneum - at least 70% by weight - was made up of terrestrial foodstuffs of similar isotopic composition and of substantially less protein content than fish. Although other low-protein terrestrial foods, even fatty meats or legumes, cannot be ruled out (Fernandes et al., 2014), the most likely contenders are cereals. Carbonized remains of cereals, mainly naked wheats and barley, have been found in abundance at Herculaneum and in the Villa dei Papiri nearby (Meyer, 1988; Ciarallo, 1994; Pagano,




Fig. 3. Stable isotope measurements of human remains from Herculaneum. a.) - compared with Isola Sacra and Velia; b.) - showing the Kernel density plot of all adults from Herculaneum by sex (F = 28; M = 37) against the estimation of % marine carbon to total dietary carbon.

1994). A cache of 117 wooden writing tablets (the ‘Murecine Tablets’) found in a wicker basket just outside the walls of Pompeii and dating from the mid-first century (29–61 CE) reveals that ‘Alexandrian wheat’ was stored in large quantities in warehouses at Puteoli (Camodeca, 1999; Wolf, 2010). Overall, the high consumption of cereals with a relatively low protein concentration, and variable contribution of marine foods, explains the narrow range of  $\delta^{13}\text{C}$  values compared with  $\delta^{15}\text{N}$  values observed at Herculaneum and other Italian Roman Imperial period coastal sites (Fig. 3a) (Craig et al., 2013).

The observed isotopic differences between the sexes at Herculaneum could simply be a matter of biology; the calorific requirements of males are known to be greater than those of females, and the undertaking of hard labour would undoubtedly exacerbate such needs leading to quantitative and qualitative dietary discrepancies. However, it is terrestrial products - mainly cereals - that provided the majority of calories regardless of sex, so this is less likely. Rather, it is the consumption of high trophic level marine fish that isotopically distinguishes males from females. In Fig. 3b we have used the  $\delta^{15}\text{N}$  to indicate the % contribution of marine foods to dietary carbon (an approximation to their weight contribution to total diet) using Eq. (1)

(above). The distributions (Fig. 3b) show that a small proportion of the males obtained a slightly greater % of their total diet from marine foods. The differences between males and females with respect to marine consumption is great (typically < 5% contribution to total diet) but the effect on their  $\delta^{15}\text{N}$  values is much more pronounced, since fish makes a disproportional contribution to dietary protein.

It is reasonable to suppose that occupation is a key variable which determines these sex-related dietary differences. Men had primary access to marine foods in as much as fishing and trade in fish products were male-dominated activities. In general, the uneven distribution of power, which in a traditional society lay with males, and other social factors, would have played a part in permitting or restricting access to fish, both within the families of fishermen, and in the wider community (Garnsey, 1999).

### 3.3. Variation by age

When the sample is subdivided into specific age classes (15–20, 20–30, 30–40, 40–50, 50+ years) there are no significant differences in  $\delta^{15}\text{N}$  values (Kruskal-Wallis  $\chi^2 = 7.0$ ,  $p = 0.13$ ) or in  $\delta^{13}\text{C}$  values (Kruskal-Wallis  $\chi^2 = 6.4$ ,  $p = 0.17$ ). If the data are first disaggregated by sex and then compared by age, there are no significant differences between males and females in any of the age classes, or between males and females of different age classes (Robust ANOVA  $\delta^{15}\text{N}$  interaction between age classes and sex  $F = 2.6$ ,  $p = 0.05$ ; Robust ANOVA  $\delta^{13}\text{C}$  interaction between age classes and sex  $F = 1.1$ ,  $p = 0.37$ ). Overall, the intra-population stable isotopic variation at Herculaneum is related to sex but seems to be less dependent on an individual's age at death. However,  $\delta^{15}\text{N}$  values are significantly different between adults < 30 years old (i.e. 15–30) compared with those older than 30 years (Wilcoxon rank sum test with continuity correction  $W = 385.5$ ,  $p = 0.04$ ). When testing for the interaction with sex within these age classes, the robust ANOVA shows no significant interaction for  $\delta^{15}\text{N}$  ( $F = 2.6$ ,  $p = 0.05$ ). Boxplots in Fig. 4 show that older males at Herculaneum tended to have diets richer in marine foods. Conversely, females and younger males have diets more similar to each other. There are no significant differences in  $\delta^{13}\text{C}$  values between these broader (15–30, 30+ years) age ranges (Wilcoxon rank sum test with continuity correction  $W = 469.5$ ,  $p = 0.30$ ).

Certainly, we would expect some age and sex related differences at Herculaneum. By 30 years of age, most men might be supposed to have received a boost in their disposable income, allowing access to greater

quantities of more expensive commodities such as fish. By 30 years old most men would have entered into their first marriage (Saller, 1996; Aldrete, 2008; Garnsey and Saller, 2015) and most sons are likely to have lost their fathers, becoming *sui iuris* ('of one's own right'), and had themselves inherited the role – including the legal and financial independence – of the head of the household (*paterfamilias*). A second consideration is the high prevalence of slaves and freedmen in the city. Demographic estimates based on the Marble Album of Herculaneum suggest that a significant proportion of the town's urban population (ca. 23%) were freedmen (de Ligt and Garnsey, 2012). The study proposes that ca. 69% of the adult male citizen population were ex-slaves, and that ca. 60% of the entire urban slave population at Herculaneum were manumitted by the age of 30. With manumission came possible elevation to the rank of Roman citizen in accordance with the laws passed in the time of Augustus. Freedmen were normally involved *ipso facto* in a patronage relationship with their ex-masters, supposing the latter were still alive – in which case the freedman might benefit from a legacy (Aldrete, 2008; Garnsey and Saller, 2015). In either eventuality, their standard of living and subsistence is likely to have improved following manumission, again permitting access to new foods. In comparison, female slaves were manumitted later in life, if at all. Furthermore if, as seems probable, freedmen were involved in the processing and trade of fish (Curtis, 2005), they are also likely to have had preferential access to this resource, and be well-represented among those in the sample with high  $\delta^{15}\text{N}$  values.

A potential methodological explanation for the absence of strong isotopic differences by narrower age classes at Herculaneum is that the measurements are of collagen which is synthesised at different times within an individual's lifespan. As bone collagen turnover rate is relatively slow, a substantial proportion of collagen derived from earlier in life will still be present at death. For example, from studies of collagen turnover rates in femoral bone (Hedges et al., 2007) we estimate that 63% of collagen in a 45 year old male, or 53% in a female of the same age, is derived from foods consumed before 30 years of age. Furthermore, the rate of bone turnover slows dramatically following adolescence, meaning that younger individuals' skeletons contain relatively more collagen synthesised from foods consumed closer to the time of death than older individuals. A slightly faster turnover rate may be anticipated in the rib samples analysed in this study, nevertheless, these measurements are unlikely to reflect true differences between the age classes. Indeed, the differences that we observed are probably underestimations of the true dietary differences between the old and the younger adults at their time of death.

Age related dietary differences were observed at Isola Sacra. Here, Prowse et al. (2005) showed that age and  $\delta^{15}\text{N}$  are positively correlated for both sexes. Explaining this observation is far from straightforward as it is not possible to distinguish whether individuals consumed a greater proportion of fish in later life or whether high fish consumers simply lived longer. Given the difficulties in interpreting isotopic data from bone collagen due to its slow turnover rate, and the lack of strong evidence of age related differences within the 'living population' at Herculaneum, we suggest that the latter explanation is more likely. At Velia, there are no overall significant differences by age class (Craig et al., 2009), however one group of adult males ( $n = 11$ ) are relatively enriched in  $^{15}\text{N}$  (i.e. > 9.6‰) compared to the rest of the population. Interestingly, 10 are over 30 years of age and also have a much higher prevalence of external auricular exostosis (Crowe et al., 2010), a pathology caused by regular exposure to cold water which is most likely linked to sea-related occupations (Crowe et al., 2010).

## 4. Conclusion

Overall, the data from the catastrophic assemblage at Herculaneum emphasizes the difficulty in interpreting intra-population isotopic variability in attritional cemetery populations, as are commonly encountered in archaeological research. There is less overall variation in




Fig. 4. Boxplot of  $\delta^{15}\text{N}$  values at Herculaneum by age in years and sex.

$\delta^{15}\text{N}$  at Herculaneum compared to Velia and Isola Sacra regardless of sample size or demographic composition. This result is most easily explained by the short-lived nature of the population. Diets change over generations as the result of changes in the economy and food supply, as well as cultural shifts. Therefore, the range of foods eaten by individuals living contemporary lives may be considerably narrower than revealed through isotopic analysis of individuals buried in cemeteries, which are also influenced by selective mortality and selective burial. This has important implications for considering the *durée* of cemetery populations before making comparisons of any osteological datasets. Despite these interpretative issues, underlying trends are still observable between osteological and isotopic datasets, for example due to occupation (Crowe et al., 2010). However, we suggest that these correlations are probably related to an individual's long-term diet rather than directly attributable to specific periods of their life, given the attenuated dietary record represented by adult bone collagen. At the very least, such direct associations need to be questioned. Further comparison of stable isotope values of collagen from tissues with different turnover rates is needed to help resolve these issues. Finally, we confirm there is clear differentiation of diet by sex as observed in attritional Roman populations, related to differential access of males and females to marine foods.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.jasrep.2017.08.008>.

## Acknowledgements

We thank Andrew Millard, University of Durham, for his most valuable assistance with estimating collagen turnover rates, Dr. Pietro Guzzo and Dr. Teresa Elena Cinquantaquattro of the former Soprintendenza Speciale per i Beni Archeologici di Napoli e Pompei for their continued support and two anonymous referees for their valued thoughts and comments.

## References

- Aldrete, G., 2008. Daily Life in the Roman City: Rome, Pompeii, and Ostia. Oklahoma University Press, Norman, Oklahoma.
- AlQahatani, S.J., Hector, M.P., Liversidge, H.M., 2010. Brief communication: the London atlas of human tooth development and eruption. *Am. J. Phys. Anthropol.* 142, 481–490.
- Baccino, E., Ubelaker, D.H., Hayek, L.-A., Zerilli, A., 1999. Evaluation of seven methods of estimating age at death from mature human skeletal remains. *J. Forensic Sci.* 44, 931–936.
- Bisell, S., 1991. The human skeletons of Herculaneum. *Int. J. Anthropol.* 6, 1–20.
- Buikstra, J., Ubelaker, D., 1994. Standards for Data Collection from Human Skeletal Remains: Proceedings of a Seminar at the Field Museum of Natural History. Arkansas Archaeological Survey Press, Fayetteville.
- Camodeca, G., 1999. *Tabulae Pompeianae Sulpiciorum*, Editione critica dell'archivio pu-teolano dei Sulpicii I/II. Quasar, Rome.
- Capasso, L., 2000. Herculaneum victims of the volcanic eruptions of Vesuvius in 79 AD. *Lancet* 356, 1344–1346.
- Capasso, L., Capasso, L., 1999. Mortality in Herculaneum before volcanic eruption in 79 AD. *Lancet* 354, 1826.
- Capasso, L., Domenicantonio, L.D., 1998. Work-related syndesmoses on the bones of children who died at Herculaneum. *Lancet* 352, 1634.
- Ciarallo, A., 1994. Il frumento nell'area vesuviana. In: *Le ravitaillement en blé de Rome et des centres urbaines des débuts de la République jusqu'au haut empire. Actes du colloque international organisé par le Centre Jean Bérard et L'URA 994 du CNRS. Naples, 14–16 Février 1991. Centre Jean Bérard: Ecole française de Rome*, pp. 137–140.
- Craig, O.E., Biazio, M., O'Connell, T.C., Garnsey, P., Martinez-Labarga, C., Lelli, R., Salvadei, L., Tartaglia, G., Nava, A., Renò, L., Fiammenghi, A., Rickards, O., Bondioli, L., 2009. Stable isotopic evidence for diet at the imperial Roman coastal site of Velia (1st and 2nd centuries AD) in southern Italy. *Am. J. Phys. Anthropol.* 139, 572–583.
- Craig, O.E., Bondioli, L., Fattore, L., Higham, T., Hedges, R., 2013. Evaluating marine diets through radiocarbon dating and stable isotope analysis of victims of the AD79 eruption of vesuvius. *Am. J. Phys. Anthropol.* 152, 345–352.
- Crowe, F., Sperduti, A., O'Connell, T.C., Craig, O.E., Kirsanow, K., Germoni, P., Macchiarelli, R., Garnsey, P., Bondioli, L., 2010. Water-related occupations and diet in two roman coastal communities (Italy, first to third century AD): correlation between stable carbon and nitrogen isotope values and auricular exostosis prevalence. *Am. J. Phys. Anthropol.* 142, 355–366.
- Curtis, R.I., 2005. Sources for production and trade of Greek and Roman processed fish. In: Bekker-Nielsen, T. (Ed.), *Ancient Fishing and Fish Processing*. University Press, Aarhus, pp. 31–46.
- DeWitte, S.N., Stojanowski, C.M., 2015. The osteological paradox 20 years later: past perspectives, future directions. *J. Archaeol. Res.* 23, 397–450.
- Edwards, J., 2001. Philology and cuisine in de re coquinaria. *Am. J. Philol.* 122, 255–263.
- Fattore, L., Bondioli, L., Garnsey, P., Rossi, P., Sperduti, A., 2012. The human skeletal remains from Herculaneum: New evidence from the excavation of fornici 7, 8, 9, 10 and 11. Poster presented at the AAPA 81st Annual Meeting, Portland: Oregon, 11–14 April 2012. *Am. J. Phys. Anthropol.* 142.
- Ferembach, D., Schwidetzky, I., Stoukal, M., 1980. Recommendation for age and sex diagnoses of skeletons. *J. Hum. Evol.* 9, 517–549.
- Fernandes, R., 2015. A simple(R) model to predict the source of dietary carbon in individual consumers. *Archaeometry* 58, 500–512.
- Fernandes, R., Millard, A.R., Brabec, M., Nadeau, M.-J., Grootes, P., 2014. Food reconstruction using isotopic transferred signals (FRUITS): a Bayesian model for diet reconstruction. *PLoS One* 9, e87436.
- FitzGerald, C., Saunders, S., Bondioli, L., Macchiarelli, R., 2006. Health of infants in an imperial roman skeletal sample: perspective from dental microstructure. *Am. J. Phys. Anthropol.* 130, 179–189.
- Garnsey, P., 1999. *Food and Society in Classical Antiquity*. Cambridge University Press, Cambridge.
- Garnsey, P., Saller, R., 2015. *The Roman Empire: Economy, Society and Culture*, 2nd ed. University of California Press, Oakland, CA.
- Garvin, H.M., Passalacqua, N.V., 2012. Current practices by forensic anthropologists in adult skeletal age estimation. *J. Forensic Sci.* 57, 427–433.
- Hedges, R.E.M., Clement, J.G., Thomas, C.D.L., O'Connell, T.C., 2007. Collagen turnover in the adult femoral mid-shaft: modeled from anthropogenic radiocarbon tracer measurements. *Am. J. Phys. Anthropol.* 133, 808–816.
- İşcan, M.Y., Loth, S.R., Wright, R.K., 1984. Age estimation from the rib by phase analysis: white males. *J. Forensic Sci.* 29, 1094–1104.
- Jacks, M., 2011. Representativeness and bias in Archaeological skeletal samples. In: Argarwal, S., Gencross, B. (Eds.), *Social Bioarchaeology*. Chichester, Wiley-Blackwell, pp. 107–146.
- Killgrove, K., Tykot, R.H., 2013. Food for Rome: a stable isotope investigation of diet in the imperial period (1st–3rd centuries AD). *J. Anthropol. Archaeol.* 32, 28–38.
- Killgrove, K., Tykot, R.H., 2017. Diet and Collapse: A Stable Isotope Study of Imperial-era Gabii (1st–3rd Centuries AD). Reports: *J. Archaeol. Sci.* (in press).
- Krogman, W.M., İşcan, M.Y., 1986. *The Human Skeleton in Forensic Medicine*. C. C. Thomas, Springfield.
- de Ligt, L., Garnsey, P., 2012. The album of Herculaneum and a model of the town's demography. *J. Rom. Archaeol.* 25, 69–94.
- Mastrolorenzo, G., Petrone, P., Pagano, M., Incoronato, A., Baxter, P., Canzanello, A., Fattore, L., 2001. Herculaneum victims of Vesuvius in AD 79. *Nature* 410, 769–770.
- Mastrolorenzo, G., Petrone, P., Pappalardo, L., Guarino, F., 2010. Lethal impact at periphery of pyroclastic surges: evidences at Pompeii. *PLoS One* 5, e11127.
- Meyer, F., 1980. Carbonized food plants of Pompeii, Herculaneum, and the villa at Torre Annunziata. *Econ. Bot.* 34, 401–437.
- Meyer, F., 1988. Food Plants Identified from Carbonized Remains at pompeii and Other Vesuvian Sites. In: Curtis, R.I. (Ed.), *Studia pompeiana and Classica in Honor of Wilhelmina F. Jashemski*. Aristide D. Caratzas, New Rochelle, N.Y., pp. 183–229.
- Pagano, M., 1994. Commercio e consumo del grano ad Ercolano. In: *Le ravitaillement en blé de Rome et des centres urbaines des débuts de la République jusqu'au haut empire. Actes du colloque international organisé par le Centre Jean Bérard et L'URA 994 du CNRS. Naples, 14–16 Février 1991. Centre Jean Bérard: Ecole française de Rome*, pp. 141–148.
- Petrone, P.P., 2011. Human corpses as time capsules: new perspectives in the study of past mass disasters. *J. Anthropol. Sci.* 89, 3–6.
- Petrone, P., Giordano, M., Giustino, S., Guarino, F., 2011. Enduring Flouride health hazard for the vesuvius area population: the case of AD 79 *Herculaneum*. *PLoS One* 6, e21085.
- Prowse, T., Schwarcz, H.P., Saunders, S., Macchiarelli, R., Bondioli, L., 2004. Isotopic paleodiet studies of skeletons from the imperial Roman-age cemetery of Isola Sacra, Rome, Italy. *J. Archaeol. Sci.* 31, 259–272.
- Prowse, T.L., Schwarcz, H.P., Saunders, S.R., Macchiarelli, R., Bondioli, L., 2005. Isotopic evidence for age-related variation in diet from Isola Sacra, Italy. *Am. J. Phys. Anthropol.* 128, 2–13.
- Rackham, H., 1967. *Pliny the Elder: Natural History*. Harvard University Press, Massachusetts.
- Reese, D., 2002. Fish: evidence from specimens, mosaics, wall paintings, and Roman authors. In: Jashemski, W., Meyer, F. (Eds.), *The Natural History of Pompeii*. Cambridge University Press, Cambridge, pp. 274–291.
- Roberts, C.A., Grauer, A., 2001. Commentary: bones, bodies and representivity in the archaeological record. *Int. J. Epidemiol.* 30, 109–110.
- Robinson, M., Rowan, E., 2015. Roman food remains in Archaeology and the contents of a Roman sewer at Herculaneum. In: Wilkins, J., Nadeau, R. (Eds.), *Companion to Food in the Ancient World*. Wiley-Blackwell, Oxford.
- Rousseeuw, P.J., Ruts, I., Tukey, J.W., 1999. The bagplot: a bivariate boxplot. *Am. Stat.* 53, 382–387.
- Rowan, E., 2014. The fish remains from the Cardo V sewer: new insights into taphonomy, consumption and the fish economy of Herculaneum. In: *Fish and Ships: Proceedings of the Fish and Ships Conference, Rome, 18–22 June 2012*. Errance.
- Saller, R., 1996. *Patriarchy, Property and Death in the Roman Family*. Cambridge University Press, Cambridge.
- Scheuer, L., Black, S., Schaefer, M.C., 2010. *Juvenile Osteology: A Laboratory and Field Manual*. Elsevier Science.
- Todd, T.W., 1921. Age changes in the pubic bone. *Am. J. Phys. Anthropol.* 4, 1–70.
- Treggiari, S., 1993. *Roman Marriage: Iusti Coniuges from the Time of Cicero to the Time*

- of Ulpian. Oxford University Press.
- Wallace-Hadrill, A., 2011. *Herculaneum: Past and Future*. Frances Lincoln Ltd., London.
- White, T.D., Folkens, P.A., 2005. *The Human Bone Manual*. Elsevier Academic.
- Wolf, J., 2010. *Neue Rechtsurkunden aus Pompeji: Tabulae Pompeianae Novae*. Wissenschaftliche Buchgesellschaft, Darmstadt.
- Wood, J.W., Milner, G.R., Harpending, H.C., Weiss, K.M., Cohen, M.N., Eisenberg, L.E., Hutchinson, D.L., Jankauskas, R., Cesnys, G., Česnys, G., Katzenberg, M.A., Lukacs, J.R., McGrath, J.W., Roth, E.A., Ubelaker, D.H., Wilkinson, R.G., 1992. The osteological paradox: problems of inferring prehistoric health from skeletal samples [and comments and reply]. *Curr. Anthropol.* 33, 343–370.