
DIAGNOSTICA DI LABORATORIO Presidio Ospedaliero S. Eugenio - CTO

Tel. 06-51008889-2335 / Fax 06-51002972



DIAGNOSTICA INFETTIVOLOGICA MOLECOLARE

Servizio Igiene Sanità Pubblica.

Materiale: Tampone nasofaringeo

Diagnosi Molecolare

SARS-CoV-2-Interpretazione

SARS-CoV-2 gene E SARS-CoV-2 gene RdRP/S

SARS-CoV-2 gene N

Materiale: Tampone nasofaringeo Nota **POSITIVO**

Rilevato

Rilevato Rilevato

Si raccomanda di comunicare il risultato al Medico curante che valuterà se NEGATIVO l'eventuale sospensione dell'isolamento fiduciario, se POSITIVO le misure di isolamento domiciliare, in attesa di essere eventualmente contattato dal

Il Dirigente:


PCR Real Time


Referto Completo

Referto id. firmato digitalmente da Dott. in data alle ore come da normativa vigente.

Questa rappresentazione è conforme all'originale conservato presso la Regione Lazio.

Struttura del Sars-Cov2





Correlazione tra Ct e gravità della malattia

Tso CF, Garikipati A, Green-Saxena A, Mao Q, Das R. Correlation of Population SARS-CoV-2 Cycle Threshold Values to Local Disease Dynamics: Exploratory Observational Study. JMIR Public Health Surveill. 2021 Jun 3;7(6):e28265. doi: 10.2196/28265. PMID: 33999831; PMCID: PMC8176948.

Duration of infectiousness and correlation with RT-PCR cycle threshold values in cases of COVID-19, England, January to May 2020

Anika Singanayagam^{1,2}, Monika Patel^{1,2}, Andre Charlett³, Jamie Lopez Bernal⁴, Vanessa Saliba⁴, Joanna Ellis¹, Shamez Ladhani⁴, Maria Zambon¹, Robin Gopal¹

- 1. Virus Reference Department, Public Health England, Colindale, United Kingdom
- These authors contributed equally
- 3. Data and Analytical Services, Public Health England, Colindale, United Kingdom
 4. Immunisation and Countermeasures, Public Health England, Colindale, United Kingdom

Correspondence: Maria Zambon (maria.zambon@phe.gov.uk)

FIGURE 1

Detection of SARS-CoV-2 by RT-PCR targeting the RdRp gene, England, January-April 2020 (n = 754)





FIGURE 2

Relationship between RT-PCR Ct value and culture positivity in mixed effects logistic regression analysis, SARS-CoV-2, England, January–May 2020 (n = 324)



Ct: cycle threshold.




Understanding cycle threshold (Ct) in SARS-CoV-2 RT-PCR

Protecting and improving the nation's health

A guide for health protection teams

COVID-19 symptom onset schematic diagram



Whilst high Ct values may be associated with reduced infectivity, a swab taken at a single point in time does not provide information about the trajectory or subsequent course of illness.

Ct values are not directly comparable between assays and may not be reported by some RT-PCR platforms in use. Interpreting single positive Ct values for staging infectious course, prognosis, infectivity or as an indicator of recovery must be done with context about the clinical history.