
Corso di Immunologia - III anno Prof. Paolini


Lezione 24/10/2025

"L'attivazione dei linfociti T"

Il materiale presente in questo documento viene distribuito esclusivamente ad uso interno e per scopi didattici.

Mature DCs provide naive T cells with all the signals they need to be activated






How to communicate? Starting by touching each other

Intercellular Adhesion Molecules (ICAMs) of the integrin-binding immunoglobulin superfamily

The molecules involved in signal propagation are located in the contact area between the APC and the T lymphocyte

LFA-1 integrin receptor

Principal function



- Belongs to the β 2 family of integrins
- Its ligands are ICAM-1, 2, 3, expressed on APCs,and endothelium
- It is responsible for cell-cell adhesion between:

T cell/APC


CTL/target cell

Leukocyte/endothelium

LFA-1 affinity & avidity for ligands are regulated

Low Ag recognition High

Adhesion molecules participate to T cell/APC interaction




How to communicate? Starting by touching each other

Intercellular Adhesion Molecules (ICAMs) of the integrin-binding immunoglobulin superfamily

The molecules involved in signal propagation are located in the contact area between the APC and the Tlymphocyte


Antigen recognition induces a re-organization of T cell surface receptors



The region where the T cell contacts the APC is called the


immunological synapse.

In the synapse, all the interacting proteins form a structure known as the SupraMolecular Adhesion Complex. **SMAC** = SupraMolecular Activation Cluster









LFA-1


TCR

The immunological synapse


An organized membrane structure which facilitates, stabilizes and prolongs TCR/MHC/peptide interactions by:

- 1. Excluding long receptors and inhibitory receptors from the interaction area
- 2. Facilitates the aggregation and the accumulation of TCR and MHC molecules (higher local ligand concentration)
- 3. Promotes cytoskeleton rearrangement


Cytoskeletal rearrangement is involved in the formation and the stabilization of the immunological synapse




TCR aggregation upon antigen recognition




After interacting with the MHC-peptide complex, the TCR translocates to a microdomain of the membrane (LIPID RAFT) rich in cholesterol, unsaturated fatty acids, sphingomyelin and glycosphin




The immunological synapse in different types of T cell interactions



La trasduzione del segnale da parte di recettori di membrana prevede una fase citosolica e una nucleare




CD3 complex associates with TcR and initiates signal transduction



Immunoreceptor Tyrosine-based Activation Motif

Src family tyrosine kinases initiate TCR/CD3 signal transduction

Engagement of MHC-peptide initiates processes that lead to assembly of signaling complex



CD4/8-associated p56^{Lck} phosphorylates ITAMs of zeta chains, creates docking site for ZAP-70


Sito di legame alla membrana

Src family tyrosine kinases (fyn and lck) initiate signal transduction in T cells


Their activity is regulated by inhibitory and activating tyrosine residues

Csk= tyrosine kinase


CD45= tyrosine phosphatase




1. lck and fyn phosphorylate CD3 ITAMs




2. ZAP70 tyrosine kinase is recruited and activated at the cell membrane




Principali enzimi coinvolti nell'attivazione dei linfociti T



Structure of the related Syk and Zap-70 tyrosine kinases




ZAP70 activates many intracellular pathways. 1. adaptors and $PLC\gamma$




ation (n)


A rapid augmentation of cytosolic Ca²⁺ concentration is one of the earliest activation events upon antigen recognition




RAS pathway in T cell activation



TCR signal transduction leads to the activation of multiple transcription factors



T lymphocytes need 2 signals for activation







Figure 8-10 Immunobiology, 6/e. (© Garland Science 2005)

Cell surface molecules involved in T cell activation



What is the second signal?

CYTOKINE

produced by APC

MEMBRANE INTERACTION

receptor T cell ligand APC

The expression of the costimulatory signal can be:

CONSTITUTIVE

or

INDUCIBLE (inflammation, infection, tissue injury)

CD28 costimulatory molecule






Figure 8-11 Immunobiology, 6/e. (© Garland Science 2005)

- Expressed on almost all CD4+ 50% CD8+
- *Binds to CD80 and CD86 (B7.1 and B7.2), expressed on mature DC, activated B cells and macrophages
- Integrates signals of TCR/CD3 to gene activation


The signal transduced by CD28 amplifies the first signal by increasing the transcription levels of the IL-2 gene



Only "mature", "activated" APCs can provide the costimulatory signal



Why T cells need TWO signal for activation? (1)



Usually, only professional APC can give costimulatory signal

This limits the possibility of self-antigen recognition on tissue cells...

..and represents a mechanism of peripheral TOLERANCE

Figure 8-13 Immunobiology, 6/e. (© Garland Science 2005)


Why T cells need TWO signal for activation? (2)

Costimulatory signals are induced by pathogens, inflammation, tissue injury


They carry information about DANGEROUSNESS.....

...and focus T cell responses against AGGRESSIVE NON SELF

Antigen-specific T cells proliferate before becoming EFFECTORS (= clonal expansion)



Generation of effector T lymphocytes



Nature Reviews Immunology, 2004


Only antigen specific T cells proliferate!

T cell activation by antigen + costimulator

Secretion of IL-2


Expression of IL-2Rα chain; formation of high-affinity IL-2Rαβγ complex

IL-2-induced T cell proliferation




The expression of the high affinity IL-2 receptor (\alpha\beta\gamma) is tightly regulated

© Elsevier. Abbas & Lichtman: Basic Immunology, Updated 2e - www.studentconsult.com




IL-2: a growth factor for innate and adaptive lymphocytes



© Elsevier 2005. Abbas & Lichtman: Cellular and Molecular Immunology 5e www.studentconsult.com


Effector T cells migrate to the infection site to perform their function




Naive and effector lymphocytes differ phenotypically and functionally

Property	Naive T cells	Effector T cells
Cell adhesion molecules (CD2 and LFA-1)	Low	High
Trafficking patterns	HEVs* in secondary lymphoid tissue	Tertiary lymphoid tissues; inflam- matory sites
Costimulatory signal (CD28-B7 interaction)	Required for activation	Not required for activation


Effector T lymphocytes recognize the antigen again: eradication of the infection




Development of effector and memory T cells




NEGATIVE SIGNALS INACTIVATE OR KILL EFFECTOR T LYMPHOCYTES




Switching off T-specific signaling: The role of Cbl ubiquitin ligase and ubiquitination




Activation of naive T lymphocytes induces the expression of the inhibitory receptor CTLA4




Co-stimulatory and co-inhibitory molecules set the threshold for T cell activation



Principal members of B7-family proteins




Major functions of selected B7-CD28 family members




CD28-B7: initiation of immune responses

CTLA-4-B7: inhibits early T cell responses in lymphoid organs

PD-1:PD-L1,2: inhibits effector T cell responses in peripheral tissues





Inhibitory functions of CTLA-4 e PD-1

CTLA-4

PD-1

Action site

Lymphoid organs

Peripheral tissues

Inhibition phase

Upon initial activation

During effector phase

Inhibited signals

CD28 co-stimulation


Chronic stimulation

Inhibited lymphocyte subsets

CD4+ and CD8+

CD8+ > CD4+

TWO different mechanisms for peripheral TOLERANCE of T cells!



Turning off T-cell specific signaling: The role of regulatory lymphocytes (Tregs)

