Corso di Immunologia - III anno Prof. Paolini

Lezione 31/10/2025


"I linfociti T citotossici"

Il materiale presente in questo documento viene distribuito esclusivamente ad uso interno e per scopi didattici.

### Lymphocyte cytotoxicity

- Effector mechanism of cell-mediated immunity
- Directed against:
  - Infected cells (intracellular pathogens)
  - \*Tumor cells
  - \*Allogeneic cells
- 2 major lymphocyte populations:
  - □CD8+ T lymphocytes CTL
  - □Natural Killer cells NK
- Different mechanisms for target cell recognition
- Same lytic mechanisms


# Why do exist different cytotoxic lymphocyte populations?




Complementary strategies in the defense against intracellular pathogens!


Figure 2-49 Immunobiology, 6/e. (© Garland Science 2005)


#### CTL participate to the response against intracellular pathogens





...and tumor cells






CD8 T cell recognition is

specific

and

MHC class Irestricted

## Differentiation of pre-CTL to effector cells




# CD8 T CELLS MATURE IN THE THYMUS AND ARE PRE-CTLS, ABLE TO RECOGNIZE ANTIGENS BUT NOT TO KILL INFECTED TARGET CELLS


HOW ARE THEY ACTIVATED AND HOW DO THEY BECOME CTLs?

Different mechanisms for CTL differentiation

# I° Naive CD8+ lymphocytes are activated by a dendritic cell infected by an intracellular pathogen.




# ... or from an uninfected dendritic cell that cross-presents viral antigens




#### Alternative mechanisms requires help from CD4+ T cells

II°




CD4 produces IL-2 which activates CD8

III°



CD4 activates DC to express ligands that induce CD8 to produce IL-2 driving its own proliferation

# Cytotoxic mediators are acquired during differentiation: CTLs acquire tha ability to kill!!




# The synthesis of mediators is induced:

- Cytotoxic granules (perforin and granzymes)
- Cytokines (IFN $\gamma$  e TNF $\beta$ )
- Expression of FasL


EOMES = Eomesodermin EGR2 = early growth response 2

Nature Reviews | Immunology

### CTLs migrate to the site of infection...




.... where they are activated by antigen recognition and kill the infected cell!



#### MECHANISMS OF CTL-MEDIATED LYSIS

#### CYTOTOXICITY DEPENDENT ON:

Direct contact:



#### Secretory Mechanism

- · Calcium-Dependent
- · Granule Exocytosis

#### Non secretory mechanism


- · Calcium-Independent
- Fas-FAsL


Cytokine secretion: IFN $\gamma$  TNF  $\alpha/\beta$ 

#### TARGET CELL DEATH

- Osmotic lysis
- · Apoptosis

# 1° CELL-CONTACT-DEPENDENT CYTOTOXICITY: SECRETORY MECHANISM





# Antigen recognition augments LFA-1 affinity & avidity: Formation of a stable effector/target conjugate








Figure 8-28 Immunobiology, 6/e. (© Garland Science 2005)



# The secretory synapse: Immunological synapse formed by CTL and target cell







## The different phases of the secretory mechanism







Figure 8-29 Immunobiology, 6/e. (© Garland Science 2005)






# What's in CTL cytotoxic granules?


Protein in granules of cytotoxic T cells	Actions on target cells	
Perforin	Aids in delivering contents of granules into the cytoplasm of target cell	
Granzymes	Serine proteases, which activate apoptosis once in the cytoplasm of the target cell	
Granulysin	Has antimicrobial actions and can induce apoptosis	
Proteoglycans	a structural role and a protective function	



### Perforin polymerization creates pores on target cell membrane




# Model of perforin and granzyme synergy in target-cell death



Perforin polymerizes into transmembrane pores (Ca<sup>++</sup>- and pH-dependent), through which granzymes might diffuse into the cell

### Models of granzyme entry into the target cell


Perforin allows the diffusion of granzymes into the cytoplasm of the target cell by creating pores on its plasma membrane



Granzymes are recognized by specific receptors and internalized along with perforin.

Inside endosomes, perforin allows granzymes to escape into the cytosol.

### Perforin-Granzyme mediated target-cell death




Perforin induces pores and osmotic death!!

Granzyme B activates caspases and causes apoptotic death!!!

Granzyme A
releases an
endonuclease
that cuts DNA
and
induces apoptosis

### Why isn't the CTL killed?



### Codice OPIS assegnato al corso integrato tenuto da:

#### Prof. ROSSELLA PAOLINI


#### Biotecnologie (29887)

Modulo	Insegnamento	Codice OPIS
IMMUNOLOGIA I (1051487_2)	IMMUNOLOGIA (10514879)	1BH7HMGQ
IMMUNOLOGIA II (1051487_1)	IMMUNOLOGIA (10514879)	MH4E21B4

# 2° CELL-CONTACT DEPENDENT CTL CYTOTOXICITY: NON-SECRETORY MECHANISM (FAS/FASL)

#### FAS ligand (FASL):

- a TM protein induced on <u>activated</u> T cells (CD8, Th1) and released from cell surface in soluble form



# FAS engagement initiates a cascade of caspases 1.Caspase 8






Figure 6-24 part 1 of 2 Immunobiology, 6/e. (© Garland Science 2005)

# FAS engagement initiates a cascade of caspases 1.Caspase 8 2.Caspase 3




Multiple caspase 3 substrates lead to: cell shrinkage, and membrane blebbing DNA degradation, and block of DNA repair




FAS-activated caspase 8 cleaves BID, a member of the BCL-2 family

#### Presence of trophic factor




# Proapoptotic and antiapoptotic members of BCL-2 family regulate cell survival




BCL-2 BCL-XL (anti-apoptotic)

#### Absence of trophic factor

BAX




Bax, Bad Bim, Bak (pro-apoptotic) Cooperation between the extrinsic (death receptor-mediated) and the intrinsic (mitochondrion-mediated) pathways to cell




#### FAS-MEDIATED CYTOTOXICITY:

- · Mechanism of immune response regulation
- Mechanism used by tumors to evade immunological surveillance
- ·Defense mechanism of immunologically privileged organs

# NEGATIVE SIGNALS REGULATING T LYMPHOCYTE ACTIVATION FASL-induced T cell death



### FAS/FASL: a homeostatic mechanism of the immune system



#### a FasL+ tumor FasL-Soluble FasL Fas Fas+ T cell Caspase activation T cell death b Caspase activation Suicide Fratricide as L induction Fas+ T cell TCR MHC class I Fas Ltumor

# Fast-induced T cell death upon recognition of tumor cells

Some tumors evade the immune response by expressing Fas ligand (FasL) and a soluble isoform of FasL

Activated T cells can kill themselves ('suicide') and each other ('fratricide') through the same caspase-based mechanism

Nicholas P. Restifo Nature Medicine 1/6, 493 - 495 (2000)

# Cells from "immunologically privileged organs" express Fas ligand







Figure 13-12 Immunobiology, 6/e. (© Garland Science 2005)

## Two major lytic mechanisms of lymphocytes







### Cooperation between the granule-dependent and the FASmediated mechanism of cytotoxicity



### 3° CTL also produce CYTOTOXIC cytokines






Tumor Necrosis Factor

TNF  $\alpha$ 


TNF  $\beta$  (LT)

that can induce apoptosis (through TNF-RI)


### CTL-mediated cytotoxicity



Direct killing by release of granzyme B and perforin into the intercellular space between CTL and target cell




Induction of apoptosis in target cells via death receptor triggering




Indirect killing of target cells by release of Tumor Necrosis Factor-alpha and IFN-gamma


### Naive-to-memory progression after acute infection or vaccination



# The development of memory cells requires CD4+ T-cell help (cytokines, etc.)




Cellular antigen (weakly immunogenic)




Viral or bacterial antigen (strongly immunogenic)

### Memory CTL are less dependent on CD4+ T help



### Two types of cytotoxic lymphocytes



- Different recognition strategies
- Distinct activation mechanisms
  - Common lytic mechanisms

They use complementary defense strategies against intracellular pathogens and tumor cells!