


ACIDI NUCLEICI

Purine and Pyrimidine

Pyrimidine contains two pyridine-like nitrogens in a six-membered aromatic ring


Purine has 4 N's in a fused-ring structure. Three are basic like pyridine-like and one is like that in pyrrole



Nucleic Acids and Nucleotides

Deoxyribonucleic acid (DNA) and ribonucleic acid (RNA), are the chemical carriers of genetic information

Nucleic acids are biopolymers made of nucleotides, aldopentoses linked to a purine or pyrimidine and a phosphate




© 2004 Thomson/Brooks Cole

Sugars in DNA and RNA

RNA is derived from ribose


DNA is from 2'-deoxyribose

- (the ' is used to refer to positions on the sugar portion of a nucleotide)



Ribose

© 2004 Thomson/Brooks Cole




2'-Deoxyribose

Heterocycles in DNA and RNA


Adenine, guanine, cytosine and thymine are in DNA

RNA contains uracil rather than thymine




Adenine (A)
DNA
RNA


© 2004 Thomson/Brooks Cole




Guanine (G)
DNA
RNA



Cytosine (C)
DNA
RNA




Thymine (T)
DNA




Uracil (U)
RNA

Nucleotides


In DNA and RNA the heterocycle is bonded to C1' of the sugar and the phosphate is bonded to C5' (and connected to 3' of the next unit)




The Deoxyribonucleotides




2'-Deoxyadenosine 5'-phosphate



2'-Deoxyguanosine 5'-phosphate




2'-Deoxycytidine 5'-phosphate





2'-Deoxythymidine 5'-phosphate


The Ribonucleotides



Adenosine 5'-phosphate





Cytidine 5'-phosphate



Uridine 5'-phosphate

COFATTORI ENZIMATICI DI NATURA NUCLEOTIDICA



cofattori NAD^+ e NADP^+

