

Ion-Trapping, Microsomal Binding, and Unbound Drug Distribution in the Hepatic Retention of Basic Drugs

Gerhard A. Siebert, Daniel Y. Hung, Ping Chang, and Michael S. Roberts

Department of Medicine, University of Queensland, Princess Alexandra Hospital, Woolloongabba, Queensland, Australia

Received July 13, 2003; accepted September 18, 2003

ABSTRACT

This study investigated the relative contribution of ion-trapping, microsomal binding, and distribution of unbound drug as determinants in the hepatic retention of basic drugs in the isolated perfused rat liver. The ionophore monensin was used to abolish the vesicular proton gradient and thus allow an estimation of ion-trapping by acidic hepatic vesicles of cationic drugs. In vitro microsomal studies were used to independently estimate microsomal binding and metabolism. Hepatic vesicular ion-trapping, intrinsic elimination clearance, permeability-surface area product, and intracellular binding were derived using a physiologically based pharmacokinetic model. Modeling showed that the ion-trapping was significantly lower after monensin treatment for atenolol and propranolol, but not for antipyrine. However, no changes induced by monensin treatment were

observed in intrinsic clearance, permeability, or binding for the three model drugs. Monensin did not affect binding or metabolic activity in vitro for the drugs. The observed ion-trapping was similar to theoretical values estimated using the pHs and fractional volumes of the acidic vesicles and the pK_a values of drugs. Lipophilicity and pK_a determined hepatic drug retention: a drug with low pK_a and low lipophilicity (e.g., antipyrine) distributes as unbound drug, a drug with high pK_a and low lipophilicity (e.g., atenolol) by ion-trapping, and a drug with a high pK_a and high lipophilicity (e.g., propranolol) is retained by ion-trapping and intracellular binding. In conclusion, monensin inhibits the ion-trapping of high pK_a basic drugs, leading to a reduction in hepatic retention but with no effect on hepatic drug extraction.

Basic lipophilic compounds are characterized by a high volume of distribution as a result of extensive tissue uptake. The main mechanisms of such a distribution pattern are nonspecific binding to membrane phospholipids (Bickel and Steele, 1974; Francesco and Bickel, 1977; Romer and Bickel, 1979), binding to microsomal protein (Hung et al., 2002), and the sequestration of the compounds into acidic vesicular compartments such as lysosomes or mitochondria (Daniel et al., 1995). A potential consequence of an apparent irreversible sequestration of basic drugs into acidic vesicles is a potentially reduced drug bioavailability (de Duve et al., 1974; Ohkuma and Poole, 1978) or drug interactions (Daniel and Wojcikowski, 1999b; Nebbia et al., 1999). Lysosomal trapping of basic lipophilic drugs has also been demonstrated to be an

important determinant of disposition for desipramine and chloroquine and psychotropic compounds such as the piperidine and piperazine-type neuroleptics (Daniel et al., 2001). The lysosomotropic properties of basic drugs are particularly important determining drug disposition and pharmacokinetics in lysosome-rich organs such as lungs, kidneys, or the liver.

Specific studies determining the relative contribution of ion-trapping and microsomal binding to the hepatic retention of drugs or relating the relative uptake to the physicochemical properties of drugs do not seem to have been undertaken so far and are the focus of the present study. Fluorescence microscopy, acridine orange staining, and other studies using primary hepatocyte cultures have shown that H^+ ionophores, such as the carboxylic antibiotic monensin, abolish the pH gradient in the approximately 170 acidic vesicles in hepatocytes (Lake et al., 1987). Such studies have not been carried out in the isolated perfused rat liver, which not only retains intact cellular integrity and fully functional lysosomes but allows the effects of drug structure on hepatocyte wall per-

This study was supported by the National Health and Medical Research Council of Australia and the Queensland and New South Wales Lions Kidney and Medical Research Foundation.

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.

DOI: 10.1124/jpet.103.056770.

ABBREVIATIONS: DMO, dimethylloxazolidine-2,4-dione; CL_{int} , intrinsic elimination clearance; K_v , equilibrium amount ratio characterizing the vesicular ion-trapping sites (ion-trapping parameter); CV^2 , normalized variance; k_{vc} , rate constant for transport from acidic vesicles into cytosol; E, hepatic extraction ratio; MTT, mean transit time; f_{ub} , drug fraction unbound in perfusate; $f_{u,MP}$, drug fraction unbound in microsomal protein; PS, permeability-surface area product; K_b , equilibrium amount ratio characterizing the intracellular binding sites; Q, flow rate; k_{cv} , rate constant for transport from cytosol into acidic vesicles; k_e , elimination rate constant; V_B , extracellular reference space; k_{in} , influx rate constant; V_C , cellular water volume; V_{max} , maximum velocity; k_{out} , efflux rate constant.

meability, hepatic binding, sequestration, and metabolism to be described (Hung et al., 2001, 2002). Monensin infusions should allow the acidic vesicle pH gradient (and thus the ability for ion-trapping of basic drugs) to be experimentally abolished without affecting other permeation, binding, and clearance processes.

To evaluate the importance of physicochemical drug characteristics such as lipophilicity and pK_a on ion-trapping and microsomal binding and thus drug retention, we compared two drugs with similar pK_a values but different lipophilicity (apparent log octanol-water partition coefficient, $\log P_{app}$, atenolol and propranolol) and two drugs with a similar lipophilicity but different pK_a (atenolol and antipyrine). The study follows up on the recently reported structure-hepatic disposition relationships of cationic drugs in normal and diseased rat liver (Hung et al., 2001, 2002). Pharmacokinetic parameters such as hepatocellular influx, efflux, binding and elimination for the three drugs were determined and kinetic parameters were derived from an improved two-phase physiologically based organ pharmacokinetic model (Weiss and Roberts, 1996; Hung et al., 2001), and the relative contribution of ion-trapping, intracellular binding, and unbound drug to hepatic drug retention was calculated.

Materials and Methods

Chemicals

Atenolol (4-[2-hydroxy-3-[(1-methylethyl)amino]propoxy]benzeneacetamide), antipyrine (1,2-dihydro-1,5-dimethyl-2-phenyl-3H-pyrazol-3-one), monensin sodium (2-[5-ethyltetrahydro-5-[tetrahydro-3-methyl-5-[tetrahydro-6-hydroxy-6-(hydroxymethyl)-3,5-dimethyl-2H-pyran-2-yl]-2-furyl]-2-furyl]-9-hydroxy- β -methoxy- $\alpha,\gamma,2,8$ -tetramethyl-1,6-dioxaspiro[4,5]decane-7-butyric sodium salt), propranolol (1-[(1-methylethyl)amino]-3-(1-naphthalenyl)-2-propanol) all were obtained from Sigma Aldrich (St. Louis, MO). [U - ^{14}C]Dimethylloxazolidine-2,4-dione (DMO), [^{14}C]sucrose, and [3H]water were purchased from Amersham Biosciences UK, Ltd. (Little Chalfont, Buckinghamshire, UK).

In Situ Perfusion of the Isolated Rat Liver

The experimental protocol was approved by the University of Queensland Animal Ethics Committee. Perfusion of the isolated rat liver used in this study was performed as described previously (Cheung et al., 1996). Briefly, male Wistar rats, weighing 200 to 250 g, were anesthetized using an intraperitoneal injection of xylazine/ketamine (10/80 mg kg^{-1}). The laparotomized rats were heparinized with 200 units of heparin injected into the inferior vena cava. The bile duct and the portal vein were cannulated using polyethylene-10 (Clay Adams, Franklin Lakes, NJ) and an intravenous 16-gauge catheter, respectively. The liver was then perfused with 3-(*N*-morpholino-)propanesulfonic acid (MOPS) buffer containing 2% bovine serum albumin and 15% washed canine red blood cells, adjusted to pH 7.4, and oxygenated via a silastic tubing lung with 100% pure oxygen. A noncirculating peristaltic pump was used as perfusion system. Once perfusion was established the animals were sacrificed by thoracotomy and the inferior vena cava was cannulated for sample collection. The animals were placed in a temperature-controlled environment at 37°C. Assessment of liver viability was by macroscopic appearance, measurement of bile flow, oxygen consumption, and portal resistance pressure (Cheung et al., 1996).

Perfusions were adjusted to a flow rate of 15 ml/min and given a 10-min stabilizing period before the injection of the first bolus. Each bolus consisted of 50 μ l of buffer containing a given concentration of radiolabeled compound ([^{14}C]sucrose, [3H]water, and [^{14}C]DMO) or cationic drug, in the presence and absence of 0.5 mM monensin. The

injection was timed to coincide with the start of a computer-controlled fraction collector (samples were collected over 4 min: 20×1 s, 5×4 s, 5×10 s, and 4×30 s).

The collected samples were centrifuged and 75- μ l aliquots of the supernatant containing [3H]water, [^{14}C]sucrose, [^{14}C]antipyrine, or [^{14}C]DMO were taken for scintillation counting (MINAXI beta TRI-CARB 4000 series liquid scintillation counter; PerkinElmer Life Sciences, Boston, MA). The remainder was vortexed for high-performance liquid chromatography analysis and the atenolol and propranolol outflow concentrations were determined (Hung et al., 2001).

In Vitro Binding of Cationic Drugs to a Microsomal Protein Preparation

To assess the effect of monensin on hepatic drug binding and metabolism, an in vitro study with a microsomal protein preparation in the presence and absence of monensin was carried out, using buffer containing 0.35 mg/ml microsomal protein from normal livers. The unbound fraction of cationic drug in each buffer solution was estimated using an ultrafiltration method. A known concentration of the cationic drug stock solution was added to 500 μ l of each buffer solution to make final concentration of 0.05 μ M and placed in a centrifugal filter device (Microcon YM-30, 30,000 molecular weight cut-off; Millipore Corporation, Bedford, MA) and then centrifuged at 3000g for 10 min. The ultrafiltrate (in triplicate) was assayed by high-performance liquid chromatography. The fraction of drug unbound in the microsomal protein preparation (f_{UMP}) was determined as the ratio of the free concentration to total concentration of solute.

The impact of monensin on hepatic drug metabolism was assessed in these studies using intrinsic elimination clearance CL_{int} in the presence and absence of monensin estimated by the formula $CL_{int} = V_{max}/(K_m + C)$, where V_{max} is the maximum velocity, K_m is the Michaelis-Menten constant, and C the concentration of the drug. The respective K_m values were taken from the literature: antipyrine (Roberts and Rowland, 1986), atenolol (Bagwell et al., 1989), and propranolol (Ishida et al., 1992). V_{max} was determined by drug incubation with the microsomal protein preparation at 37°C and sampling at time points 0, 5, 10, and 20 min. The concentrations of drugs were analyzed by high-performance liquid chromatography (Hung et al., 2002). Data were fitted to the formula and CL_{int} estimated using Scientist (Micromath Scientist, Salt Lake City, UT) (Hung et al., 2001, 2002). The perfused liver intrinsic clearance was based on the assumption that 70% of the harvested microsomal protein was metabolically active and an average amount of 50 mg of microsomal protein was obtained per gram of liver (Roberts and Rowland, 1986).

Analytical Procedure

The high-performance liquid chromatography method used in this study has been described and validated previously (Hung et al., 2001).

Modeling and Data Fitting of the Outflow Concentration-Time Profiles of Extracellular and Cellular References

Calculation of Intracellular pH (pH_i). Changes in intracellular pH were calculated according to the method of Le Couteur et al. (1993).

Data Analysis

The two-phase physiologically based organ pharmacokinetic model describing intersinusoidal mixing (Roberts et al., 1988), transfer across the hepatocyte membrane, and the intracellular distribution and elimination kinetics (Weiss and Roberts, 1996; Weiss et al., 1997) has been further developed to account for both the ion-trapping effect (K_v , characterizing the acidic vesicular ion-trapping sites) and intracellular binding (K_b , characterizing the total of intracellular binding) for the three model cationic. Briefly, as shown in Fig. 1 the model assumes drug transfer across the cytoplasmic membrane with influx and efflux rate constants k_{in} and k_{out} , respectively, recognizing




Fig. 1. Schematic overview of hepatocellular drug transport and intracellular drug distribution, including ion-trapping, intracellular binding, and drug metabolism. D, drug; DH^+ , protonated drug; f_{uB} , fraction of drug unbound in the perfusate; k_{in} , influx rate constant; k_{out} , efflux rate constant; k_e , elimination rate constant; K_b , equilibrium amount ratio characterizing the intracellular binding sites (microsomal and non-specific binding sites); k_{cv} and k_{vc} , rate constant for transport from cytosol into acidic vesicles (lysosomes and mitochondria) or from acidic vesicles into cytosol, respectively, determining the equilibrium amount ratio K_v ($K_v = k_{vc}/k_{cv}$) that characterizes ion-trapping; and V_B , extracellular volume (vascular + Disse space).

that solute concentrations change in space and time in both phases. The stochastic approach represents the transit of a molecule through the organ as a series of sojourns in one of the two regions described by density functions. The density of cellular residence times $\hat{f}_y(s)$ describes the hepatocellular distribution and elimination kinetics.

The retention time distribution $f_y(t)$ of a drug after a single excursion in the cellular space for the resulting two-compartment cell model was obtained by standard methods in the Laplace domain, $\hat{f}_y(s) = L^{-1}[f_y(t)]$, as described previously (Weiss, 1999; Weiss et al., 2000).

$$\hat{f}_y(s) = \frac{(s + k_{vc})k_{in}}{s^2(k_{in}/k_{out})(1 + K_b) + s((k_{in}/k_{out})(k_{vc} + K_b k_{vc}) + k_e + k_{cv} + k_{in}) + (k_{in}/k_{out})k_e k_{vc} + k_{in}k_{vc}} \quad (1)$$

where the influx rate constant $k_{in} = f_{uB}PS/V_B$, is the permeation clearance per extracellular volume V_B , PS is the permeability-surface product, f_{uB} is the unbound fraction of solute in the perfusate, and k_{cv} and k_{vc} represent the rate constant for transport from cytosol into the acidic compartment (lysosomes and mitochondria) or from acidic vesicles into cytosol, respectively, determining the equilibrium amount ratio $K_v = k_{vc}/k_{cv}$ characterizing ion-trapping. K_b is the equilibrium amount ratio characterizing the intracellular binding sites (microsomal and nonspecific binding) and the elimination rate constant defined as $k_e = CL_{int}/V_C$ is the intrinsic elimination clearance normalized per cellular volume V_C (Pang et al., 1995). On comparison of the CL_{int} obtained in microsomal metabolism studies with the one determined when K_b was assigned as a slow distribution process, it was found that this resulted in an unrealistically high clearance rate. Therefore it was assumed that K_b is characterized by an instantaneous distribution process (eliminating one fitting parameter at the same time).

The hepatic transit time density function $\hat{f}(s)$ of solutes can be derived in terms of the extracellular transit time density of a non-permeating reference molecule (sucrose) $\hat{f}_B(s)$ (see eq. 2, and the density function of successive sojourn times $\hat{f}_y(s)$ of the drug molecules into the cellular space

$$\hat{f}(s) = \hat{f}_B[s + k_{in}(1 - \hat{f}_y(s))] \quad (2)$$

The fractional outflow versus time data were fitted in the time domain using a numerical inverse Laplace transformation of the appropriate transit time density function using the nonlinear regression program SCIENTIST (MicroMath Scientific Software). Data were analyzed by a sequential procedure. First, the fractional out-

flow curve $C_{sucr}(t)$ of the extracellular marker $[U-^{14}C]$ sucrose was fitted by eq. 3, whereby $\hat{f}_{cath}(s)$ accounts for the catheter and $\hat{f}_B(s)$ includes the large vessel transit time,

$$C_{sucr}(t) = \frac{\text{Dose}}{Q} L^{-1}\{\hat{f}_{cath}(s)\hat{f}_B(s)\} \quad (3)$$

and the transit time density (TTD) of the nonpermeating indicator is given by

$$\hat{f}_B(s) = p\hat{f}_1(s) + (1 - p)\hat{f}_2(s) \quad (4)$$

with

$$\hat{f}_i(s) = \exp\left\{\frac{1}{CV_i^2} \left[\frac{MTT_i}{CV_i^2/2} \left(s + \frac{1}{2MTT_i CV_i^2} \right) \right]^{1/2}\right\} \quad (i=1,2) \quad (5)$$

Equation 5 is the Laplace transform of the inverse Gaussian density function with mean MTT_i and relative dispersion. Equations 3 to 5 have been shown to adequately describe the TTD of vascular markers in the perfused rat liver (Weiss et al., 1997).

Catheter transit time density was calculated by an independent experiment fitting eqs. 4 and 5 to the catheter system outflow profile. The four parameters describing $\hat{f}_{cath}(s)$ were then fixed while fitting the liver outflow data.

The mean transit time of the extracellular reference, $MTT_B = \int_0^\infty t\hat{f}_B(t)dt$, is given by

$$MTT_B = pMTT_1 + (1 - p)MTT_2 \quad (6)$$

$V_B = MTT_B Q(1 - \text{hematocrit})$ is the sum of the sinusoidal plasma space volume accessible to sucrose, and the Disse space, $V_B = V_{Plasma} + V_{Disse}$. $Q(1 - \text{hematocrit})$ denotes the plasma flow rate. Sucrose does not distribute into erythrocytes; therefore this extracellular space value has to be corrected for hematocrit (Varin and Huet, 1985). Second, using this information the outflow concentration data of the permeating drugs, $C(t)$, were analyzed, i.e., the parameters MTT_i ($i = 1, 2$), and p of the individual fits of $[U-^{14}C]$ sucrose data were substituted as fixed parameters in $\hat{f}_B(s)$ of the model (eq. 2):

$$C(t) = \frac{\text{Dose}}{Q} L^{-1}\{\hat{f}_{cath}(s)\hat{f}(s)\} \quad (7)$$

and the parameters k_{in} , k_{out} , K_v , K_b , and k_e were estimated.

In this study, a uniform $K_v = 0.25$ was assumed in monensin

treatment (the K_v value was estimated from the volume ratios of the subcellular compartments to the remaining cytoplasmic fraction: 0.21 for mitochondria plus lysosomes and 0.79 for cytosol (Rhoades and Pflanzner, 1996) and used as a fixed parameter in data fitting to obtain k_{in} , k_e , k_{out} , and K_b . The parameters k_{in} , k_{out} , k_e , and K_b were then fixed to estimate K_v for the data in the not monensin-treated groups.

The relative contribution of ion-trapping, intracellular binding and distribution of unbound drug for hepatic sequestration of the three model drugs is therefore given by

$$\frac{K_v}{1 + K_v + K_b}, \quad \frac{K_b}{1 + K_v + K_b} \quad \text{and} \quad \frac{1}{1 + K_v + K_b} \quad (8)$$

respectively.

The cellular distribution volume of water was estimated by fitting the ^3H -water outflow data with eq. 7 using the density function for water $\hat{f}_w(s)$ instead of $\hat{f}(s)$. The latter differ only with regard to the respective tissue retention time densities $\hat{f}_v(s)$ assuming no cytoplasmic binding of water and $k_{out,w} = k_{in,w}v_{c,w}$ where $v_{c,w}$ denotes the normalized cellular water volume eq. 1 for well mixed intracellular distribution reduces (Weiss et al., 2000)

$$\hat{f}_{y,w}(s) = \frac{k_{in,w}/v_{c,w}}{k_{in,w}/v_{c,w} + s} \quad (9)$$

and is substituted in eqs. 2 and 7.

Nonparametric estimates of hepatic availability, mean transit time and normalized variance were determined from the outflow concentration versus time profiles for the reference from eqs. 10 to 13 using the trapezoidal method with exponential tail approximation.

$$F = \frac{Q \cdot \text{AUC}}{D} \quad (10)$$

AUC = $\int_0^\infty C(t)dt$ is the area under the solute concentration versus time curve, Q is the perfusate flow rate, and D is the drug dose administered (expressed in molar equivalents).

$$\text{MTT} = \frac{\int_0^\infty t C(t) dt}{\text{AUC}} \quad (11)$$

$$\text{CV}^2 = \frac{\sigma^2}{\text{MTT}^2} \quad (12)$$

where

$$\sigma^2 = \frac{\int_0^\infty t^2 C(t) dt}{\int_0^\infty C(t) dt} - \text{MTT}^2 \quad (13)$$

Statistical Analysis

All data are presented as mean \pm S.D. unless otherwise stated. Statistical analysis was performed using Tukey's post hoc or Student's t test where appropriate. A $p < 0.05$ was taken as significant.

Results

Figure 2 shows typical logarithms of measured and predicted (fitted data) time-outflow fraction profiles before and after monensin treatment for the three model cationic drugs antipyrine, atenolol, and propranolol. The peak outflow concentrations were in the order atenolol < antipyrine < propranolol and are inversely related to drug lipophilicity as




Fig. 2. Outflow profile fractions (log scale) for antipyrine, atenolol, and propranolol in controls and after treatment. Empty symbols represent controls and solid symbols represent monensin treatment. Dashed and solid lines stand for fitted data in control and treatments, respectively.

defined by their individual $\log P_{app}$ (atenolol 0.14, antipyrine 0.33, and propranolol 3.10). Monensin treatment did not affect the outflow concentration-time profile of antipyrine (with the lowest pK_a and a low $\log P_{app}$) but changed the outflow profiles of atenolol and propranolol (Fig. 2). The effect of

monensin on the outflow profile of propranolol is dramatic (Fig. 2) and in the presence of monensin the profile assumes a shape similar to antipyrine (Fig. 2). The outflow profiles of antipyrine (as a positive control that does not show ion-trapping) remained unchanged and were virtually superimposed before and after treatment (Fig. 2).

Table 1 shows the nonparametric moments parameters hepatic extraction, mean transit time, and normalized variance for the three model drugs. The mean transit time of propranolol showed treatment-induced significant differences ($p < 0.01$). No other differences for comparison of controls and monensin treatment groups were observed ($p > 0.05$; Table 1).

The model-derived kinetic parameters K_v , K_b , CL_{int} , and PS for hepatic drug disposition of the three model drugs are summarized in Table 2. The vesicular ion-trapping constant K_v was significantly lower after monensin treatment for atenolol and propranolol (K_v control/ K_v treatment: 30- and 35-fold, respectively; $p < 0.001$), but not for antipyrine. However, no changes induced by monensin treatment were observed for the intracellular binding constant, the intrinsic elimination clearance, or the permeability-surface area product values for the three model drugs (Table 2).

A comparison of the predicted and observed model-derived ion-trapping parameter K_v values are presented in Table 3. The theoretical and observed values were in good agreement. Atenolol and propranolol both showed considerable ion-trapping (high K_v values), whereas the corresponding value for antipyrine was relatively small (Table 3).

The results of an in vitro binding, the calculated microsomal association constant and in vitro intrinsic elimination clearance for the three drugs in a microsomal protein preparation are shown in Table 4. Monensin treatment had no effect on binding behavior or metabolic drug elimination. No differences before and after treatment were found to exist for the three model drugs (Table 4). It is also apparent that the in vitro microsomal estimates of K_b (Table 4) compare favorably with those obtained from modeling the perfused liver data (Table 2). The microsomal CL_{int} values obtained in vitro are also in general good agreement with the model-derived CL_{int} values in the in situ isolated perfused liver study (Tables 2 and 4).

The relative contribution of ion-trapping, intracellular (microsomal and unspecific) binding, and the distribution of unbound drug to the hepatic retention of the three drugs are summarized in Fig. 3. Ion-trapping is the most important determinant of hepatic drug retention for atenolol and the least important contributor for antipyrine, whereas ion-trapping and intracellular binding are equally important for hepatic drug retention for propranolol (Fig. 3). The distribution

of unbound drug is the least important determinant for propranolol and the most important for antipyrine (Fig. 3).

Discussion

The study presented here was undertaken to define the relative importance of intracellular binding, vesicular (lysosomal and mitochondrial) drug sequestration, and distribution of unbound drug as determinants of hepatic solute retention and also to investigate the relative importance of pK_a lipophilicity for this process. Monensin was used to determine the relative contribution of vesicular ion-trapping in the hepatic retention of three model cationic drugs: atenolol, antipyrine, and propranolol. In vitro microsomal studies produced independent estimates of microsomal binding and intrinsic clearance that showed that monensin treatment does not affect these parameters and therefore can be used to estimate the contribution of vesicular ion-trapping to hepatic drug retention. The difference in K_b calculated from in vitro microsomal data (Table 4) and the model-derived values from the perfused liver (Table 2) suggests that microsomal binding accounts for most of hepatic binding, as has been previously suggested for the normal rat and human livers (McLure et al., 2000; Hung et al., 2002). A comparison of CL_{int} values obtained in the in vitro study (Table 4) and results from the in situ liver perfusion (Table 2) show the reliability and suitability of the model used in the study.

In this investigation, the intact, viable, perfused in situ rat liver was used to study sequestration of basic drugs in acidic organelles of the liver. By using monensin as an inhibitor of vesicular sequestration, the processes governing the pharmacokinetic events in the disposition of these basic drugs were quantified. This is, as far as we are aware, the first attempt to determine the differential contribution of ion-trapping, microsomal binding, and distribution of unbound solute to the hepatic retention of basic drugs with differing physicochemical properties in the isolated perfused rat liver.

Previous studies with ionophores have used cultured hepatocytes (Lake et al., 1987; Myers et al., 1995) or liver slices (Daniel et al., 1995; Daniel and Wojcikowski, 1999a). Lake et al., (1987) used monensin to characterize the number of acidic compartments in intact hepatocytes (Lake et al., 1987; Myers et al., 1995), and Myers et al. (Lake et al., 1987; Myers et al., 1995) applied flow cytometry to make dynamic measurements of the lysosomal pH in living hepatocytes. The studies of Daniel et al. (1995) have been focused on the contribution of two factors, nonspecific drug binding to membrane phospholipids and ion-trapping of cationic drugs in tissue slices (Daniel et al., 1995; Daniel and Wojcikowski, 1999a) and the contribution of lysosomal trapping to the total

TABLE 1

Non-parametric moments for model cationic drugs (mean \pm S.D., $n = 6$)

There were significant differences between controls and treatment groups for the mean transit time of propranolol (** $p < 0.01$). No other differences for the non-parametric moment parameters were found between control and treatment groups ($p > 0.05$).

	log P_{app}	pK_a	Hepatic Extraction Ratio		Mean Transit Time (s)		Normalized Variance	
			Control	Treatment	Control	Treatment	Control	Treatment
Atenolol	0.14	9.60	0.26 \pm 0.10	0.16 \pm 0.02	30.9 \pm 2.76	29.1 \pm 2.88	1.44 \pm 0.14	1.52 \pm 0.39
Antipyrine	0.33	1.45	0.20 \pm 0.03	0.17 \pm 0.09	67.5 \pm 10.1	55.8 \pm 7.34	0.48 \pm 0.12	0.62 \pm 0.10
Propranolol	3.10	9.45	0.95 \pm 0.03	0.92 \pm 0.05	119 \pm 20.3	79.5 \pm 15.3**	1.07 \pm 0.25	0.86 \pm 0.39

log P_{app} , log octanol/water partition coefficient; pK_a , negative logarithm of the ionization constant.

TABLE 2

Kinetic parameters derived from the two-phase stochastic model fitting for model cationic drugs (mean \pm S.D., $n = 6$)
There were significant differences for comparison of control and treatment groups for atenolol and propranolol ($p < 0.001$) but not for antipyrine.

	K_v		K_b		CL_{int} (ml min ⁻¹ g ⁻¹ liver)		PS (mL min ⁻¹ g ⁻¹ liver)		p
	Control	Treatment	Control	Treatment	Control	Treatment	Control	Treatment	
Atenolol	7.37 \pm 1.89	0.25	0.81 \pm 0.45	0.81 \pm 0.45	1.68 \pm 1.73	1.70 \pm 1.73	3.01 \pm 1.54	3.14 \pm 1.66	
Antipyrine	0.24 \pm 0.06	0.25	0.51 \pm 0.14	0.51 \pm 0.14	0.47 \pm 0.41	0.49 \pm 0.43	26.2 \pm 7.87	27.4 \pm 7.87	
Propranolol	8.79 \pm 1.79	0.25	8.65 \pm 1.13	8.65 \pm 1.13	11.8 \pm 2.44	12.1 \pm 2.31	35.9 \pm 8.64	37.3 \pm 9.07	

K_v ($= k_{off}/k_{off}^{app}$), the equilibrium amount ratio characterizing the slowly accessible pool for ion-trapping; K_b , the rapidly equilibrating intracellular binding sites (microsomal and nonspecific binding); PS, permeability-surface area product; CL_{int} , intrinsic elimination clearance.

tissue uptake of neuroleptics (Daniel et al., 1995; Daniel and Wojcikowski, 1999a). The extent of uptake of drugs into tissue slices was found to be tissue-specific, and the contribution of the two uptake mechanisms to be strongly drug-dependent (the contribution of ion-trapping to the total drug uptake of perazine in the liver was given as 40%).

The model used in our investigation differs from previous studies in that a perfused liver, in which the vascular hepatic architecture remains intact and the liver is oxygenated by red blood cells, was used to ensure optimal hepatic vitality and drug extraction in the perfused organ in situ. In our present study, we characterized both hepatic extraction and retention, including the respective relative contribution of ion-trapping, and also those of microsomal protein binding (the major hepatic factor contributing to drug binding; Hung et al., 2002) and of unbound drug uptake into the liver. The results are of considerable clinical relevance, especially for drugs that show high first pass hepatic extraction and are targeted to the liver, such as the statins (Garcia et al., 2003) or cytostatic drugs.

Analysis of hepatic extraction, mean transit time, and normalized variance by moments showed that there were no differences in these parameters for the three drugs before and after monensin treatment (Table 1) with the exception of propranolol that showed a decrease in mean transit time from 119 to 79.5 s post-treatment. This change is also evident in Fig. 2, where the change in outflow profile and especially the different tail sections of the curves are indicative of this change. Together with monensin not affecting in vitro microsomal binding and intrinsic clearance, it is evident that the major impact of monensin is on ion-trapping of basic drugs in hepatic acidic vesicles, as has been shown previously by histology (Lake et al., 1987).

According to the hepatic transport model used in this work (Fig. 1), hepatic extraction is affected by metabolism, permeability, and perfusate flow, whereas the mean transit time is defined not only by these parameters but also by ion-trapping (K_v) and microsomal binding (K_b). Confirmation of the relationships is provided by an independent assessment of the effects of monensin on hepatic extraction and mean transit time (Table 1), relative to controls. Furthermore, the in vitro study suggests that the drug fraction unbound by microsomal protein (surrogate of K_b) remains unchanged by monensin treatment (Table 4). Changes in mean transit time after monensin treatment are most evident for propranolol using a model-independent approach (Table 1). The change in K_v (Table 2) is consistent with this change in mean transit time due to ion-trapping.

The comparison of kinetic parameters before and after monensin treatment showed no differences for intracellular binding, intrinsic elimination clearance, or permeability (Table 2). Differences between control and treatment groups were observed for the vesicular ion-trapping of atenolol and propranolol (with their relatively high pK_a) but not for antipyrine (Table 2). The post-treatment ion-trapping was estimated from the volume ratios of the subcellular compartments to the remaining cytoplasmic fraction and fixed at 0.25. This calculated value is very close to the ion-trapping found for antipyrine in the control group. Because ion-trapping is defined by the relative permeation rates into and out of the acidic vesicles, it becomes obvious that this parameter is a potent marker of vesicular ion-trapping.

TABLE 3

Comparison of predicted and observed, model-derived K_v (equilibrium amount ratio characterizing the vesicular ion-trapping) values for the three model drugs

Drug	pK_a	Lysosomal to Intracellular Concentration Ratio ^a	Mitochondrial to Intracellular Concentration Ratio ^a	Predicted K_v ^b	Observed Model-Derived K_v ^c
Atenolol	9.60	738	3.97	8.17	7.37 ± 1.89
Antipyrine	1.45	1.00	1.00	0.21	0.24 ± 0.06
Propranolol	9.45	736	3.96	8.16	8.97 ± 1.79

^a Vesicular to cytosolic concentration ratio = $(1 + 10^{pK_a - pH_v}) / (1 + 10^{pK_a - pH_i})$ (Myers et al., 1995), where $pH_i \approx 7.27$ is the assumed cytosolic pH (Le Couteur et al., 1993), $pH_v \approx 4.4$ is the assumed lysosomal pH (Daniel et al., 2001) and $pH_m \approx 6.67$ is the assumed mitochondrial pH in the fasted state (Pietri et al., 2001).

^b Given that of lysosomal and mitochondrial to the cytosolic fraction is 1 and 20% (Rhoades and Pflanzner, 1996), the overall unbound drug vesicles/intracellular distribution ratio (v_i) for the three model drugs can be estimated from the individual organelle volume fraction and concentration ratio above using the equation: $K_v = f_{lys} \times l_i + f_{mito} \times m_i$.

^c Data fitting results using the physiological two-phase organ pharmacokinetic model.

TABLE 4

In vitro binding and intrinsic elimination clearance for the model cationic drugs in a microsomal protein preparation (mean ± S.D., $n = 4$)

No significant differences before and after treatment were found to exist for drug fraction unbound by microsomal protein, calculated microsomal association constant and internal elimination clearance for the three drugs ($p > 0.05$).

log P_{app}	pK_a	Drug Fraction Unbound by Microsomal Protein		Microsomal K_b ($ml^{-1} mg$)		Microsomal CL_{int} ($ml \min^{-1} g^{-1}$ liver)	
		Control	Treatment	Control	Treatment	Control	Treatment
log P_{app}	pK_a						
Atenolol	9.60	0.91 ± 0.06	0.91 ± 0.07	0.43 ± 0.09	0.46 ± 0.09	1.08 ± 0.35	1.11 ± 0.24
Antipyrine	1.45	1.00 ± 0.09	0.98 ± 0.07	0.61 ± 0.04	0.60 ± 0.06	0.75 ± 0.28	0.68 ± 0.13
Propranolol	9.45	0.49 ± 0.05	0.47 ± 0.02	5.93 ± 0.71	5.07 ± 0.89	9.44 ± 1.14	9.46 ± 0.73

log P_{app} , log octanol/water partition coefficient; pK_a , negative logarithm of the ionization constant; microsomal K_b , calculated association constant for the microsomal protein preparation; microsomal CL_{int} , intrinsic elimination clearance for the microsomal protein preparation.




Fig. 3. Comparison of the relative contribution (percentage) of vesicular ion-trapping (black columns), intracellular binding (lighter columns) and the fraction of unbound drug (white bars) to hepatic disposition of the three model drugs.

Ion-trapping may be estimated from considerations of the pHs and fractional volumes of the various cellular components, assuming that distribution is instantaneous and the resulting steady-state ratios are reflect in the observed ratios from dynamic (nonsteady-state) studies. When the unbound drug concentration is assumed to be identical in both the intracellular and perfusate compartments, the intracellular to perfusate concentration ratio for a drug with a given pK_a is given by $1 + 10^{pK_a - pH_i} / 1 + 10^{pK_a - pH_p}$ (Goldstein et al., 1974), where pH_i is the intracellular pH and pH_p is the perfusate pH (Table 3). A comparison of the predicted ion-trapping and the experimental ratios obtained by data fitting showed good similarity (Table 3). It is recognized that the predicted ion-trapping is only an approximation as a range of pHs have been reported for intracellular pH (7.19–7.29) (Le Couteur et al., 1993; Burns et al., 1999; Pietri et al., 2001), mitochondrial pH (6.7–7.0) (Soboll et al., 1980), and lysosomal pH (4–5) (MacIntyre and Cutler, 1988; Myers et al., 1995; Proost et al., 1997) as well as differing lysosomal fractional volumes (0.68–1%) (Rhoades and Pflanzner, 1996).

The relative contributions of ion-trapping, microsomal binding and distribution of unbound drug to the hepatic sequestration of the three model drugs is summarized in Fig. 3. It is apparent that neither pK_a nor lipophilicity is a sole determinant of drug distribution. Whereas the retention of the polar drug atenolol with its high pK_a is mainly determined by ion-trapping, and intracellular binding and distribution of unbound drug contribute only in a minor way, the retention of the neutral drug antipyrine is mainly decided by the distribution of intracellular unbound drug. The combination of high lipophilicity and high pK_a of propranolol results in intracellular binding and ion-trapping determining its retention in equal parts (Fig. 3).

It has been suggested that lysosomes account for as much as 10% of the distribution of the basic drugs imipramine, biperiden, and chlorpromazine in the rat liver (Ishizaki et al., 1996). This effect is likely to be reduced by the neutralization of the lysosomal and mitochondrial pH at the higher therapeutic plasma concentration of the drugs. Furthermore it has been shown that external propranolol concentrations of 10^{-6} , 10^{-5} , and 10^{-4} M increased the intralysosomal pH in highly purified rat liver, from pH 5.5 to approximate pHs of 5.6, 5.75, and 6.1, respectively (Ishizaki et al., 2000). Given that the uptake of imipramine into purified lysosomes shows a lack of dependence on either temperature or countertransport (Ishizaki et al., 2000), it is probable that a passive ion-trapping mechanism is responsible for the uptake of cationic drugs into lysosomes. The effect of neutralization of pH on uptake into mitochondria is less marked because their pH is only an order of magnitude different to physiological pH, whereas the intralysosomal differs by several orders of magnitude. Ishizaki et al. (1996) also showed that weak bases suppressed uptake of bases into lysosomes more than into mitochondria.

Could abolishing of the ion-trapping effect be justified therapeutically? Monensin is routinely used in the cattle and

poultry industry as feed additive, and lysosomal inhibitors have already been proposed as possible potent antimalarial therapeutics (Gumila et al., 1997) and also been considered as anticancer drugs (Singh et al., 1999; Park et al., 2002). However, monensin has been characterized by a narrow safety margin and may cause lethal toxicoses, especially when coadministered with other drugs (Nebbia et al., 1999). Furthermore, accidental poisoning in animals (Nebbia et al., 1999) and an isolated case of lethal monensin ingestion in a human patient suggest that large doses of lysosomal inhibitors must be considered potentially dangerous for humans (Caldeira et al., 2001). Nevertheless the possibility to use lysosomal inhibitors to manipulate hepatic (or pulmonary) drug extraction to achieve a higher systemic availability for drugs that show a pronounced first pass effect is of interest.

In conclusion, the relative contributions of ion-trapping, intracellular binding, and distribution of unbound drug of the basic drugs atenolol, antipyrine, and propranolol and the importance of pK_a and lipophilicity for total hepatocellular retention were determined. Ion-trapping is the dominant determinant of intracellular distribution of a polar drug with high pK_a and low lipophilicity (i.e., atenolol) but for a drug with a comparable high pK_a and high lipophilicity (i.e., propranolol) intracellular binding and ion-trapping are equally important for hepatic drug retention. The intracellular distribution of a drug with low pK_a and low lipophilicity (i.e., antipyrine) is characterized mainly by the distribution of intracellular unbound drug. The methodology used in this study may become a powerful and important tool in assessing and predicting differential intracellular drug distribution and retention pharmacokinetics.

Acknowledgments

We thank Prof. Michael Weiss and Dr. Yuri Anissimov for comments on aspects of the mathematical modeling.

References

- Bagwell EE, Webb JG, Walle T, and Gaffney TE (1989) Stereoselective uptake of atenolol into storage granules isolated from PC12 cells. *J Pharmacol Exp Ther* **249**:476–482.
- Bickel MH and Steele JW (1974) Binding of basic and acidic drugs to rat tissue subcellular fractions. *Chem Biol Interact* **8**:151–162.
- Burns SP, Cohen RD, Iles RA, Bailey RA, Desai M, Germain JP, and Going TC (1999) Zonation of gluconeogenesis, ketogenesis and intracellular pH in livers from normal and diabetic ketoacidotic rats: evidence for intralobular redistribution of metabolic events in ketoacidosis. *Biochem J* **343**:273–280.
- Caldeira C, Neves WS, Cury PM, Serrano P, Baptista MA, and Burdman EA (2001) Rhabdomyolysis, acute renal failure and death after monensin ingestion. *Am J Kidney Dis* **38**:1108–1112.
- Cheung K, Hickman PE, Potter JM, Walker NI, Jericho M, Haslam R, and Roberts MS (1996) An optimized model for rat liver perfusion studies. *J Surg Res* **66**:81–89.
- Daniel WA, Bickel MH, and Honegger UE (1995) The contribution of lysosomal trapping in the uptake of desipramine and chloroquine by different tissues. *Pharmacol Toxicol* **77**:402–406.
- Daniel WA and Wojcikowski J (1999a) Lysosomal trapping as an important mechanism involved in the cellular distribution of perazine and in pharmacokinetic interaction with antidepressants. *Eur Neuropsychopharmacol* **9**:483–491.
- Daniel WA and Wojcikowski J (1999b) The role of lysosomes in the cellular distribution of thioridazine and potential drug interactions. *Toxicol Appl Pharmacol* **158**:115–124.
- Daniel WA, Wojcikowski J, and Palucha A (2001) Intracellular distribution of psychotropic drugs in the grey and white matter of the brain: the role of lysosomal trapping. *Br J Pharmacol* **134**:807–814.
- de Duve C, de Barse T, Poole B, Trouet A, Tulkens P, and Van Hoof F (1974) Commentary. Lysosomotropic agents. *Biochem Pharmacol* **23**:2495–2531.
- Francesco CD and Bickel MH (1977) Membrane lipids as intracellular binders of chlorpromazine and related drugs. *Chem Biol Interact* **16**:335–346.
- Garcia MJ, Reinoso RF, Sanchez Navarro A, and Prous JR (2003) Clinical pharmacokinetics of statins. *Methods Fund Exp Clin Pharmacol* **25**:455–479.
- Goldstein A, Aronow I, and Kalman SM (1974) Principles of drug action, in *The Basis of Pharmacology*, pp 227–300. John Wiley & Sons, New York.
- Gumila C, Ancelin ML, Delort AM, Jeminet G, and Vial HJ (1997) Characterization of the potent in vitro and in vivo antimalarial activities of ionophore compounds. *Antimicrob Agents Chemother* **41**:523–529.
- Hung DY, Chang P, Cheung K, McWhinney B, Masci PP, Weiss M, and Roberts MS (2002) Cationic drug pharmacokinetics in diseased livers determined by fibrosis index, hepatic protein content, microsomal activity and nature of drug. *J Pharmacol Exp Ther* **301**:1079–1087.
- Hung DY, Chang P, Weiss M, and Roberts MS (2001) Structure-hepatic disposition relationships for cationic drugs in isolated perfused rat livers: transmembrane exchange and cytoplasmic binding process. *J Pharmacol Exp Ther* **297**:780–789.
- Ishida R, Obara S, Masubuchi Y, Narimatsu S, Fujita S, and Suzuki T (1992) Induction of propranolol metabolism by the azo dye sudan III in rats. *Biochem Pharmacol* **43**:2489–2492.
- Ishizaki J, Yokogawa K, Hirano M, Nakashima E, Sai Y, Ohkuma S, Ohshima T, and Ichimura F (1996) Contribution of lysosomes to the subcellular distribution of basic drugs in the rat liver. *Pharm Res (NY)* **13**:902–906.
- Ishizaki J, Yokogawa K, Ichimura F, and Ohkuma S (2000) Uptake of imipramine in rat liver lysosomes in vitro and its inhibition by basic drugs. *J Pharmacol Exp Ther* **294**:1088–1098.
- Lake JR, Van Dyke RW, and Scharschmidt BF (1987) Acidic vesicles in cultured rat hepatocytes. Identification and characterization of their relationship to lysosomes and other storage vesicles. *Gastroenterology* **92**:1251–1261.
- Le Couteur DG, Rivory LP, and Pond SM (1993) Hepatic intracellular pH during the prereplicative period following partial hepatectomy. *Am J Physiol* **264**:G767–G773.
- MacIntyre AC and Cutler DJ (1988) The potential role of lysosomes in tissue distribution of weak bases. *Biopharm Drug Dispos* **9**:513–526.
- McLure JA, Miners JO, and Birkett DJ (2000) Nonspecific binding of drugs to human liver microsomes. *Br J Clin Pharmacol* **49**:453–461.
- Myers BM, Tietz PS, Tarara JE, and LaRusso NF (1995) Dynamic measurements of the acute and chronic effects of lysosomotropic agents on hepatocyte lysosomal pH using flow cytometry. *Hepatology* **22**:1519–1526.
- Nebbia C, Ceppal L, Dacasto M, Carletti M, and Nachtmann C (1999) Oxidative metabolism of monensin in rat liver microsomes and interactions with tiamilin and other chemotherapeutic agents: evidence for the involvement of cytochrome P-450 3A subfamily. *Drug Metab Dispos* **27**:1039–1044.
- Ohkuma S and Poole B (1978) Fluorescence probe measurement of the intralysosomal pH in living cells and the perturbation of pH by various agents. *Proc Natl Acad Sci USA* **75**:3327–3331.
- Pang KS, Barker F, Simard A, Schwab AJ, and Goresky CA (1995) Sulfation of acetaminophen by the perfused rat liver: the effect of red blood cell carriage. *Hepatology* **22**:267–282.
- Park WH, Seol JG, Kim ES, Kang WK, Im YH, Jung CW, Kim BK, and Lee YY (2002) Monensin-mediated growth inhibition in human lymphoma cells through cell cycle arrest and apoptosis. *Br J Haematol* **119**:400–407.
- Pietri S, Martel S, Culcasi M, Delmas-Beauvieux MC, Canioni P, and Gallis JL (2001) Use of diethyl(2-methylpyrrolidin-2-yl)phosphonate as a highly sensitive extra- and intracellular 31P NMR pH indicator in isolated organs. Direct NMR evidence of acidic compartments in the ischemic and reperfused rat liver. *J Biol Chem* **276**:1750–1758.
- Proost JH, Roggeveld J, Wierda JM, and Meijer DK (1997) Relationship between chemical structure and physicochemical properties of series of bulky organic cations and their hepatic uptake and biliary excretion rates. *J Pharmacol Exp Ther* **282**:715–726.
- Rhoades R and Pflanzner R (1996) *Human physiology*. Harcourt Brace College Publishers, Philadelphia, PA.
- Roberts MS, Donaldson JD, and Rowland M (1988) Models of hepatic elimination: comparison of stochastic models to describe residence time distributions and to predict the influence of drug distribution, enzyme heterogeneity and systemic recycling on hepatic elimination. *J Pharmacokin Biopharm* **16**:41–83.
- Roberts MS and Rowland M (1986) Correlation between in-vitro microsomal enzyme activity and whole organ hepatic elimination kinetics: analysis with a dispersion model. *J Pharm Pharmacol* **38**:177–181.
- Romer J and Bickel MH (1979) Interactions of chlorpromazine and imipramine with artificial membranes investigated by equilibrium dialysis, dual-wavelength photometry and fluorimetry. *Biochem Pharmacol* **28**:799–805.
- Singh M, Ferdous AJ, and Jackson TL (1999) Stealth monensin liposomes as a potentiator of adriamycin in cancer treatment. *J Control Release* **59**:43–53.
- Soboll S, Elbers R, Scholz R, and Heldt HW (1980) Subcellular distribution of di- and tricarboxylates and pH gradients in perfused rat liver. *Hoppe Seylers Z Physiol Chem* **361**:69–76.
- Varin F and Huet PM (1985) Hepatic microcirculation in the perfused cirrhotic rat liver. *J Clin Invest* **76**:1904–1912.
- Weiss M (1999) Cellular pharmacokinetics: effects of cytoplasmic diffusion and binding on organ transit time distribution. *J Pharmacokin Biopharm* **27**:233–256.
- Weiss M, Kuhlmann O, Hung DY, and Roberts MS (2000) Cytoplasmic binding and disposition kinetics of diclofenac in the isolated perfused rat liver. *Br J Pharmacol* **130**:1331–1338.
- Weiss M and Roberts MS (1996) Tissue distribution kinetics as determinant of transit time dispersion of drugs in organs: application of a stochastic model to the rat hindlimb. *J Pharmacokin Biopharm* **24**:173–196.
- Weiss M, Stedtler C, and Roberts MS (1997) On the validity of the dispersion model of hepatic drug elimination when intravascular transit time densities are long-tailed. *Bull Math Biol* **59**:911–929.

Address correspondence to: Professor Michael S. Roberts, Department of Medicine, University of Queensland, Princess Alexandra Hospital, Woolloongabba Qld 4102, Australia. E-mail: mroberts@soms.uq.edu.au