



# Biomolecules: lipids


# Lipids: functions.

## Structure





## Energy storage

## Signalling.




## *Organic biomolecules: lipids*

- ◆ Organic amphiphilic compounds insoluble in water
- ◆ Easily extracted from animal and vegetal cells using apolar solvents
- ◆ Fundamental to build cell's shape and organelles
- ◆ They form a heterogeneous class, difficult to classify in a simple manner




# Compartmentalization is essential for a living cell



# Lipids Classification

■ The major classes of lipids are:

1. Fatty acids.
2. Triglycerols.
3. Glycerphospholipids.
4. Sphingolipids.
5. Steroids.



## *A short list of lipid classes*


- ◆ **Fatty acids (FA):** organic carboxylic acids with long carbon chains ( $C > 7$ ), saturated, unsaturated, subjected to esterification
- ◆ **Glycerides:** mono-, di-, tri-esters of fatty acids and glycerol
- ◆ **Ceramides:** esters of sphingosine and fatty acids
- ◆ **Phospholipids:** di-esters of glycerol-3-Phosphate or esters of sphingosine-phosphate
- ◆ **Cerebrosides (glycolipids):** esters of sphingosine with 1 FA and 1 sugar with 6-C (hexose)
- ◆ **Gangliosides:** glycolipids in which the sugar moiety is complex and branched
- ◆ **Steroids:** derivatives of cholesterol (cicle-pentane-peri-hydro-phenanthrene)
- ◆ **Terpenoids:** compounds made by repetitions of isoprenes units (2-methyl-butadiene)

## *Heat (energy) production*

Compound	Kcal/g
Methane	13.3
Octane	11.5
Stearic Acid (lipids)	9.5
Alanine (protein)	4.4
Glucose (sugars)	3.7

Lipids allow efficient energy storage, also due to the fact that they are stored in an anhydrous manner.

## The most common FAs in biological membranes



In the membrane there are always FA with an even number of C.

In case there are, double bonds are always in cis configuration.

The old notation  $\omega$ -3,  $\omega$ -6 and  $\omega$ -9 refers to the position of the first double bond, counting how many carbon atoms are left before the end



## Some naturally occurring Fatty Acids


Carbon skeleton	Structure*	Systematic name†	Common name (derivation)	Melting point (°C)
12:0	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>10</sub> COOH	<i>n</i> -Dodecanoic acid	Lauric acid (Latin <i>laurus</i> , "laurel plant")	44.2
14:0	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>12</sub> COOH	<i>n</i> -Tetradecanoic acid	Myristic acid (Latin <i>Myristica</i> , nutmeg genus)	53.9
16:0	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>14</sub> COOH	<i>n</i> -Hexadecanoic acid	Palmitic acid (Latin <i>palma</i> , "palm tree")	63.1
18:0	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>16</sub> COOH	<i>n</i> -Octadecanoic acid	Stearic acid (Greek <i>stear</i> , "hard fat")	69.6
20:0	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>18</sub> COOH	<i>n</i> -Eicosanoic acid	Arachidic acid (Latin <i>Arachis</i> , legume genus)	76.5
24:0	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>22</sub> COOH	<i>n</i> -Tetracosanoic acid	Lignoceric acid (Latin <i>lignum</i> , "wood" + <i>cera</i> , "wax")	86.0
16:1(Δ <sup>9</sup> )	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> CH=CH(CH <sub>2</sub> ) <sub>7</sub> COOH	<i>cis</i> -9-Hexadecenoic acid	Palmitoleic acid	-0.5
18:1(Δ <sup>9</sup> )	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>7</sub> CH=CH(CH <sub>2</sub> ) <sub>7</sub> COOH	<i>cis</i> -9-Octadecenoic acid	Oleic acid (Latin <i>oleum</i> , "oil")	13.4
18:2(Δ <sup>9,12</sup> )	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub> CH=CHCH <sub>2</sub> CH=CH(CH <sub>2</sub> ) <sub>7</sub> COOH	<i>cis</i> -, <i>cis</i> -9,12-Octadecadienoic acid	Linoleic acid (Greek <i>linon</i> , "flax")	-5
18:3(Δ <sup>9,12,15</sup> )	CH <sub>3</sub> CH <sub>2</sub> CH=CHCH <sub>2</sub> CH=CHCH <sub>2</sub> CH=CH(CH <sub>2</sub> ) <sub>7</sub> COOH	<i>cis</i> -, <i>cis</i> -, <i>cis</i> -9,12,15-Octadecatrienoic acid	α-Linolenic acid	-11
20:4(Δ <sup>5,8,11,14</sup> )	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub> CH=CHCH <sub>2</sub> CH=CHCH <sub>2</sub> CH=CHCH <sub>2</sub> CH=CH(CH <sub>2</sub> ) <sub>3</sub> COOH	<i>cis</i> -, <i>cis</i> -, <i>cis</i> -, <i>cis</i> -5,8,11,14-Icosatetraenoic acid	Arachidonic acid	-49.5

INSATURATED FATTY ACIDS always have the *cis* form of the double bonds.

Angle of about 30°.


The larger the number of *cis* bonds, the lower the possibility to form Vdw interactions (lower melting points).

Free fatty acids are normally absent in the cell.




## *FAs packing*

- ◆ 3D packing or assembly of FAs depends on the degree of unsaturation (no. of double bonds)
- ◆ The main consequence of this packing is the melting temperature: saturated FAs are solid at RT, unsaturated FAs have lower  $T_m$ )
- ◆ FAs spontaneously assemble in aqueous solutions in order to decrease the hydrophobic effect of water




## *FAs packing*

- ◆ 3D packing or assembly of FAs depends on the degree of unsaturation (no. of double bonds)
- ◆ The main consequence of this packing is the melting temperature: saturated FAs are solid at RT, unsaturated FAs have lower  $T_m$ )
- ◆ FAs spontaneously assemble in aqueous solutions in order to decrease the hydrophobic effect of water




Melting point increases with molecular mass and decreases with the degree of insaturation.





Fatty acids from plants tend to have a higher degree of insaturation and are liquid at room temperature,

By industrial hydrogenation, oils from plants become solid at r.t. (higher melting point).  
Trans double bonds are introduced.

Acido elaidico (trans)



# Lipid hydrogenation




The partial hydrogenation of fats with industrial methods leads to the formation of unsaturated lipids in *trans* configuration.

***Reaction with numerous applications, especially in the treatment of oils and fats in the food industry; through a partial hydrogenation the vegetable oils, normally liquid, are converted into solid or semi-solid fats (such as margarine).***



Typically, the reagent used is gaseous hydrogen,  $H_2$ , in the presence of metal catalysts such as Ni, Pd or Pt. The reaction is generally carried out under rather drastic conditions, at high temperatures and pressures.

Main dietary sources of trans fats for the adult American population. Average intake of trans fat equal to 5.8 g (2.6% of total calories).






As early as 1988 there were suggestions in the scientific literature that trans fats could be a cause of the large increase in coronary artery disease. In 1994, it was estimated that trans fats caused 20,000 deaths annually in the US from heart disease.

1. Food and nutrition board, institute of medicine of the national academies (2005). [Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids \(Macronutrients\)](#). National Academies Press. p. 423.
2. Food and nutrition board, institute of medicine of the national academies (2005). [Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids \(Macronutrients\)](#). National Academies Press. p. 504.
3. Mozaffarian D, Katan MB, Ascherio A, Stampfer MJ, Willett WC (April 13, 2006). "Trans Fatty Acids and Cardiovascular Disease". *New England Journal of Medicine* 354 (15): 1601–1613.

Trans unsaturated fats increase the risk of chronic cardiovascular disease by inducing an increase in LDL cholesterol and a decrease in HDL cholesterol.

Atherosclerotic deposit



The biochemical mechanism that leads to the increased risk of chronic cardiovascular disease related to the intake of trans fats is not clear. It is hypothesized that since they are not recognized by lipases, they remain in circulation for longer, increasing the formation of atherosclerotic plaques.



Pancreatic Lipase

"On a per-calorie basis, trans fats appear to increase the risk of CHD more than any other macronutrient, conferring a substantially increased risk at low levels of consumption (1 to 3% of total energy intake)". The study (3) estimates that between 30,000 and 100,000 cardiac deaths per year in the United States are attributable to the consumption of trans fats.


- Denmark (since 2003) has regulated the trans fat content, it cannot exceed 2% in the single ingredient. A 50% reduction in chronic cardiovascular diseases is expected.
- Switzerland adopted the same limits in 2008.
- New York City and California imposed a declaration of hydrogenated fats contents in restaurants.

## ***FATTY ACIDS REACTIONS***

- ◆ They are weak acids with  $K_a$   $10^{-5} - 10^{-6}$ , decreasing with increasing chain length
- ◆ They can form:
  - esters after reacting with alcohols
  - amides after reacting with amines
  - salts after reacting with a strong base (NaOH / KOH)
- ◆ They can be oxidised by enzymes to release energy (beta-oxidations)

## *Which reactions do FAs undergo?*

- ◆ Unsaturated FAs can also be oxidised by oxygen radicals, giving rise to epoxides, di-oles, peroxides
- ◆ These byproducts are degraded into aldehydes and short chain FA, and are responsible for the bad smell/taste of out-of-date fat







epoxide

di-ole

peroxide

# Carboxylic acid + alcohol = ester









## *Mono-, di- and tri-glycerides*


- ◆ Esters of FA(s) with glycerol (1,2,3-propan-tri-ole)
- ◆ Triglycerides are neutral lipids used for storage, they are confined in specialised vacuoles of special cells (adipose cells)




1-stearoyl-2-linoleyl-3-palmitoyl-glycerol

Triglycerides are energy storage molecules in the adipocytes.



The following diagram may help you remember the components of a triglyceride.



© 2001 Brooks/Cole - Thomson Learning



Glycerol





1-Stearoyl, 2-linoleoyl, 3-palmitoyl glycerol, a mixed triacylglycerol

## *Soap production*


- ◆ Soap is the sodium / potassium salt of long chain FAs
- ◆ It is produced from either animal or vegetal fat
- ◆ The reaction is an alkaline hydrolysis performed at high temperature
- ◆ This reaction produces 1 mol of glycerol and 3 mol of salt per 1 mol of triglyceride
- ◆ The % of unsaturated FA and presence of additives give solid or liquid soap at room temperature

# Saponification Reaction (alkaline hydrolysis)






## Detergent activity (soap) (soap=fatty acid salt)




micella




## *Phospholipids*

- ◆ These molecules are the scaffold of the membrane lipid bilayer and contain a polar head
- ◆ The term phospholipids include two types of polar lipids present in biological membranes:
  - **Glycerol-phospholipids**, synthesized starting from glycerol-3-phosphate (G3P, an intermediate of glucose metabolism)
  - **Sphingolipids**, synthesized from sphingosine (a complex aliphatic amino-alcohol)




Phosphatidic acid: ester of glycerol with phosphoric acid and two fatty acids



# *Phosphoglycerides*


Glycerol + 2 fatty acids + phosphate + alcohol





In phosphoglycerides a fourth (polar/charged) component is bound to the phosphate..




## Common classes of glycerophospholipids




Name of X—OH	Formula of -X	Name of Phospholipid
Water	—H	Phosphatidic acid
Ethanolamine	—CH <sub>2</sub> CH <sub>2</sub> NH <sub>3</sub> <sup>+</sup>	Phosphatidylethanolamine
Choline	—CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>3</sub> <sup>+</sup>	Phosphatidylcholine (lecithin)
Serine	—CH <sub>2</sub> CH(NH <sub>3</sub> <sup>+</sup> )COO <sup>-</sup>	Phosphatidylserine
<i>myo</i> -Inositol		Phosphatidylinositol
Glycerol	—CH <sub>2</sub> CH(OH)CH <sub>2</sub> OH	Phosphatidylglycerol
Phosphatidylglycerol		Diphosphatidylglycerol (cardiolipin)


***A typical mammalian phospholipid:  
phosphatidyl-choline (PC)***



**Phosphatidic acid** is the simplest phospholipid.



*Phospholipases are enzymes specialised in the hydrolysis of single ester bonds*




## *Sphingolipids*


- ◆ These lipids account for 20-30% of plasma membrane, especially in cells of the central nervous system
- ◆ In this case the scaffold is the trans unsaturated 18-C amino alcohol called **sphingosine**

# Sphingosine and its derivatives


18 Carbon atoms amino-alcohol




Sphingosine + fatty acid = ceramide



A ceramide looks like a diglyceride




Sphingomyelin looks like a phospholipid  
and it is abundant in neurons







Glycosphingolipids are abundant in the SNC.




# Sphingolipids

Are important components of the myeline sheath. In humans they make up to 25% of lipids.




sphingomyelin

# Sphingolipids





# Membrane lipids are amphipathic (amphiphilic)




## *Glycerol-phospholipids vs sphingolipids*

- ◆ Despite the different chemical origin, the two molecules have a similar 3D spatial arrangement and a similar charge (polar) distribution




## Gangliosides




- ◆ Both Gangliosides and Cerebrosides contain sugars → they are always present only on the external leaflet of the membrane bilayer
- ◆ Gangliosides are responsible for the blood groups

# Formation of lipid bilayer: an energy minimization process

shape of lipid molecule      packing of lipid molecules




(A)



lipid micelle

lipid bilayer

(B)






ENERGETICALLY UNFAVORABLE



planar phospholipid bilayer  
with edges exposed to water




sealed compartment  
formed by phospholipid  
bilayer

ENERGETICALLY FAVORABLE

# Detergents

**Figure of a Detergent Monomer**




**Figure 1A**

**Figure 1B**

**Molecular Shapes of Detergents and Lipids**



**Figure 2A**



**Figure 2B**








**Figure 2C**



**Figure 2D**


# Hydrophobic effect: lipid sequestration from water increases entropy






# Sterols

- ◆ This class is found almost exclusively in eukaryotes
- ◆ They have a common cyclic non-aromatic scaffold derived from cicle-pentane-peri-hydro-phenanthrene
- ◆ They have both structural and hormonal/signalling function
- ◆ The presence of 4 fused rings rigidifies the molecule → once inserted into the membranes it affects fluidity
- ◆ In mammals the precursor form all steroid hormones is cholesterol, whose OH group in C3 makes it slightly polar



# *Sterols*


- ◆ This class is found almost exclusively in eukaryotes
- ◆ They have a common cyclic non-aromatic scaffold derived from cicle-pentane-peri-hydro-phenanthrene
- ◆ They have both structural and hormonal/signalling function
- ◆ The presence of 4 fused rings rigidifies the molecule → once inserted into the membranes it affects fluidity
- ◆ In mammals the precursor of all steroid hormones is cholesterol, whose OH group in C3 makes it slightly polar



## ***Steroid hormones***

- ◆ **Glucocorticoids** (ex. Cortisol): hormones synthesised by the surrenal cortex glands, affecting the metabolism of carbohydrates, proteins and lipids. They also affect inflammatory and stress responses.
- ◆ **Mineral-corticoids** (ex. Aldosterone): hormones synthesised by the surrenal cortex glands, affecting water and salt excretion from kidneys.
- ◆ **Androgens** (ex. Testosterone) and **estrogens** (ex. beta-estradiol): crucial hormones for a correct sexual development and functionality, they are synthesised by either testes (male) or ovaries (female)
- ◆ They are all slightly more soluble than cholesterol





# Cholesterol




- From the diet and hepatic synthesis.
- insoluble, found in membranes.
- Steroid hormones synthesis.
- Bile salts synthesis.
- High blood levels-> cardiovascular diseases.



# Membranes fluidity



**(b) Transizione con e senza colesterolo**






Copyright 1999 John Wiley and Sons, Inc. All rights reserved.

Cholesterol transportation in lipoproteic complexed (LDL)

## Vitamin D


- ◆ Vitamin D is a regulator of  $\text{Ca}^{2+}$  homeostasis
- ◆ It is formed after photolysis of C9-C10 bond by UV light, followed by a spontaneous isomerization to vitamin D2 or D3
- ◆ Both these forms must be hydroxylated in the kidney (position C1) and in the liver (position C25) to give the active compound: 1- $\alpha$ -25-di-hydroxy-colecalciferol.



R = X vitamin D3  
R = Y vitamin D2


## Vitamin D

- ◆ The active form of vitamin D (1,25-dihydroxyvitamin D) induces an increase in the plasma concentration of  $\text{Ca}^{2+}$ , thus favouring its uptake by intestinal cells.
- ◆ Without vitamin D, only 10 to 15% of dietary calcium and about 60% of phosphorus is absorbed. The interaction with its receptor increases the efficiency of intestinal calcium absorption to 30 to 40% and phosphorus absorption to approximately 80%
- ◆ In this way these ions are deposited into bones and teeth.




## *Terpenes and terpenoids*

- ◆ Heterogenous class of hormones, co-enzymes, vitamins sharing only a part of the scaffold: a repeat of isoprene units




## Coenzyme Q (ubiquinone)

- ◆ Shuttle of electrons in the respiratory chain in mitochondria, chloroplasts and in prokaryotes
- ◆ It is stably inserted in the membranes



## *Vitamin A (retinol)*



- ◆ Vitamin A is a derivative of beta-carotene, once oxidised into the aldehyde form (retinal) it is incorporated into the proteins of vision, where it helps transducing the light impulse into images
- ◆ Low levels supplied by the diet induce a reduced by-night vision
- ◆ Extremely low levels induce blindness (poverty-related blindness)



**X** = CH<sub>2</sub>OH retinol  
**X** = CHO retinal

## Vitamin K

- ◆ This vitamin is only produced by plants and bacteria
- ◆ In humans it is produced by the intestinal commensals
- ◆ It is crucial for a proper blood coagulation




Vitamin K1 (plants)

Vitamin K2 (bacteria)




## *Vitamin E*

- ◆ Also known as alpha-tocopherol is an anti-oxidant, able to scavenge reactive oxygen species (ROS)
- ◆ It prevents oxidation of membranes and proteins




Prostaglandins are synthesized from arachidonic acid.



Hormone-like effect:

Fever, contraction, inflammation, asthma...

Their synthesis is inhibited by anti-inflammatory drugs such as aspirin.



**Figure 2.** Arachidonic acid is metabolized to produce inflammatory mediators. Many current anti-inflammatory and pain medicines are inhibit some portion of the arachidonic acid pathways.

