


<https://doi.org/10.1038/s42003-024-06001-2>

The consequences of viral infection on protists



Victoria Fulgencio Queiroz¹, Juliana Miranda Tatará², Bruna Barbosa Botelho¹,
Rodrigo Araújo Lima Rodrigues¹, Gabriel Magno de Freitas Almeida^{1,2,3}✉ &
Jonatas Santos Abrahão^{1,3}

Protists encompass a vast widely distributed group of organisms, surpassing the diversity observed in metazoans. Their diverse ecological niches and life forms are intriguing characteristics that render them valuable subjects for in-depth cell biology studies. Throughout history, viruses have played a pivotal role in elucidating complex cellular processes, particularly in the context of cellular responses to viral infections. In this comprehensive review, we provide an overview of the cellular alterations that are triggered in specific hosts following different viral infections and explore intricate biological interactions observed in experimental conditions using different host-pathogen groups.

The taxon “Protozoa” was formalized by the English biologist, comparative anatomist, and paleontologist Richard Owen in 1858¹. At the time, the taxon comprised “*numerous organisms of minute size retaining the form of nucleated cells, which manifest the common organic characters, but without the distinctive superadditions of true plants or animals*”¹. Since then, the terms protozoa, protozoans, protista and protists have been used to designate a variety of eukaryotic groups^{2,3}. As this taxon does not encompass well established clades that diverged from it, protists form a paraphyletic group. Similar to other paraphyletic taxa, it proves challenging to identify universal characteristics among all its members². Nevertheless, protists can be defined as predominantly unicellular eukaryotic organisms, which do not develop their tissues through the process of embryonic stratification^{4,5}. Their cell morphology showcases an astonishing diversity of forms, functions, and survival strategies. They are widespread worldwide, and comprise marine, freshwater, terrestrial, symbiotic, and pathogenic strains^{2,6,7}. However, taxonomists estimate that the known representatives compose only a small portion of the total of protists on Earth today, and that the diversity of this group is greater than that of metazoans^{6,7}.

With such abundance and richness, these eukaryotes are part of communities with a wide diversity of organisms and are especially studied for their interactions with pathogens or regarding their own pathogenicity⁷. A frequent example in literature are amoebae. Most amoebae are free-living protists of great diversity that have garnered significant attention due to their intricate associations with a plethora of microorganisms from all domains of life^{8–19}. Free-living amoebae occupy different ecological niches and have already been found in a wide range of natural and anthropized environments⁷. The most notorious feature

of amoebae is their ability to alter their cell shape by creating temporary extensions of cytoplasm known as pseudopods, which serves both for feeding and movement⁷. Due to their foraging behaviors and predatory nature, amoebae are often considered as professional phagocytes, mainly consuming microorganisms to fulfill their nutritional needs⁹. However, many microorganisms successfully evade the phagocytic pathway and thrive within the amoebae, turning them into transmission vehicles or incubators and enabling various types of ecological relationships to occur, from symbiotic to parasitic⁹. Besides, their cell biology and active grazing behavior serve as a widely used infection route for giant viruses, although miscellaneous entry mechanisms have already been described for this phylum of viruses^{10,20,21}. Given the substantial size of their viral particles, they can trigger phagocytosis in amoebae, creating entry opportunities^{11–14}. In fact, the efficiency of this route lead other giant viruses that lack the requisite size for phagocytosis to employ a mimicry strategy to use the phagocytic pathway as an entry route¹⁵.

The myriad of interactions involving amoebae renders them exceptional experimental models to explore and investigate a diversity of scientific fields¹⁶. Over the years, amoebae have contributed to the understanding of a wide array of subjects. These encompass the unraveling of mechanisms related to the resistance and pathogenicity of microorganisms, the intricacies of cell locomotion, the functioning of non-muscle contractile systems, the dynamics of populations and communities, the implications of cell nucleus removal and transplantation, events involving horizontal gene transfer, and the evolution of organelles, that may provide valuable insights into the origin and evolution of eukaryotic cells^{9,22–29}. This shows that the study of protists can help advance different fields of science. However, to

¹Federal University of Minas Gerais, Institute of Biological Sciences, Department of Microbiology, Belo Horizonte, Minas Gerais, Brazil. ²The Norwegian College of Fishery Science, Faculty of Biosciences, Fisheries and Economics, UiT - The Arctic University of Norway, Tromsø, Norway. ³These authors contributed equally: Gabriel Magno de Freitas Almeida, Jonatas Santos Abrahão. ✉e-mail: gabriel.d.almeida@uit.no


gain a more comprehensive understanding, it is important that we delve deeper into the basic aspects of protist cell biology.

Viruses are unique tools to study protists biology

Viruses have been unique tools in helping to comprehend complex biological processes in different host models. As mandatory intracellular parasites, viruses have an intimate relationship with their host cells, often depending on and controlling their cell structure, metabolism, biochemical

machinery and behavior^{30,31}. The diversity of structures, genomes, and replication strategies that viruses exhibit reflects thousands-to-billions of years of coevolution with their hosts, some older than the origin of the eukaryotic cell³². This is a key point in biology, as over the centuries, the study of host-virus interactions has led to important discoveries. For instance, in genetics, these findings include the discovery of DNA as the source of heritage, mRNA, mRNA processing, RNA interference, as well as the regulation of gene expression, transcriptional control elements, and

Fig. 1 | Overview of the current extent of the known protist virosphere. a Protist supergroups are shown on the x-axis while viral groups are shown on the y-axis. Highlighted squares on the matrix represent a group of viruses infecting a respective group of protists. The clades Viridiplantae, Fungi and Metazoa were removed from the dendrogram (latest eukaryotic phylogeny was retrieved and adapted from Keeling & Eglit, 2023²²³). Viral groups at family or genus level (unclassified cressdnavirus and preplasmivirus at phylum level and unclassified imitervirus and picornavirus at order level) are also on the y-axis. **b** The viral counts at genus level for each virus class (DNA or RNA, single-stranded or double-stranded) were retrieved from VirusHostdb²²⁴ and manual curation of the literature. Abbreviations: double stranded (ds), single stranded (ss), negative sense (−), positive sense (+), double-stranded DNA genome that has an RNA intermediate (dsDNA-RT). Detailed information regarding viruses and respective hosts used to make this figure can be found in supplementary data file (tab 1).






Fig. 2 | Overview of the main changes in the *A. castellanii* cell during different giant viruses' infection. a "Trophozoite cell shape" to represent viruses that did not interfere in the cell shape structure. b "Rounded cell shape" to represent only the viruses reported to trigger cell rounding. Red text: cell modifications.

transcription factors^{30,31,33,34}. Additionally, these studies have contributed significantly to our comprehension of biochemistry, elucidating tyrosine kinases and signal transduction pathways^{30,31,33}. Furthermore, they have shed light on immunology and many other intrinsic aspects of host cells. These include our understanding of histocompatibility antigen function, interferons, cellular oncogenes, tumor suppressor proteins, and apoptosis in eukaryotic hosts^{30,31,33,35}. They've also contributed to our knowledge of the CRISPR/Cas9 mechanism in bacterial and archaeal hosts, which is now widely used for genome editing for many biotechnological purposes³⁶.

Undoubtedly, the field of virology has made substantial contributions to science, extending beyond its significance in diseases. Throughout history, the discovery of viruses has primarily been driven by the concerns of human health, and the health of other animals and plants of economic importance^{30,33,37}. This implies that most of the described viruses have direct connections to humans, whether in the realms of economics, medicine, or biotechnology, resulting in an extremely anthropocentric known virosphere³⁷. However, the viruses known to humankind represent only a small fraction of the viral diversity on Earth. As viruses infect organisms of the three domains of life, and some studies estimate the count of viral particles on Earth to be of the order of 10^{31} , there remains a large number of undiscovered viruses^{35,38}. Still, due to the absence of a universal molecular marker for viruses, the true extent of global viral diversity remains largely unknown.

Despite having a small number of identified representatives so far, viruses that infect protists exhibit a remarkable diversity (Fig. 1). One of the earliest mentions of virus-like particles (VLP) in protozoa seems to have been documented in 1960 within *Entamoeba histolytica*³⁹. Over the years, many taxa of viruses that infect protists have been identified, with representatives containing DNA or RNA genomes of various topologies and sizes, which can exist either in a naked form or be enclosed within capsids, that also exhibit diverse sizes and symmetries^{40–47}. Among these viruses, a group that garners significant attention due to the complexity of their genomes and the size of their particles are the viruses of the phylum *Nucleocytoviricota*. Regardless of their hypothetical monophyletic origin, the viruses belonging to this phylum display a broad host range within eukaryotes. This group includes the giant viruses (GVs) that infect amoebae, the ones holding the distinction of possessing the largest genomes and viral particles found in the virosphere⁴⁸. Although these viruses infect amoebae in the laboratory, their natural hosts are still uncertain. Over the past two decades, there has been dedicated research into the exceptional biological traits of these viruses, the enigmatic composition of their genomes and particles, and their ecological significance.

Virology is an integrative science. Host-parasite interactions can reveal valuable information about cellular alterations and host defense mechanisms that can be used to study basic cellular processes^{30,33}. During a viral infection, the virion and host experience modifications of such magnitude that both start to present a different nature, sometimes even shifting the cellular role to solely producing the viral progeny. In consideration of this, the term 'virocell' was coined to refer to this specific moment when a virus is actively infecting a cell⁴⁹. One of the most characteristic changes during a viral infection is the establishment of a viral factory (VF), that can be formed in both the cytoplasm or nucleus of the cell. It is the place where viral progeny is assembled, and often, the genetic material is replicated. Its formation is usually accompanied by extensive cytoplasmic rearrangements⁵⁰. Therefore, studying host-pathogen interactions can uncover how protists respond to viral infections and highlight the features that were evolutionarily selected to counter viral threats. Moreover, viruses can be engineered for various biotechnological purposes, including gene delivery and genome editing. This technology can help elucidate protist biology for basic research or practical purposes⁵¹. Besides, these studies can also provide enlightenment into general cell biology that may apply to a wider range of eukaryotic organisms. In this review, we provide an overview of the effects of virus infection and replication on protists and discuss some prospects that the investigation into viral infections in protists could offer.

Cell machinery and morphology modulation in protist cells during viral infections

Over the past two decades, studies concerning giant viruses have highlighted virus-protist interactions⁵². Amoeba, particularly the genus *Acanthamoeba* and the species *Vermamoeba vermiformis*, are commonly studied hosts that suffer cell machinery and morphological changes after infection⁵³. The ciliate *Tetrahymena* sp. also shows alterations following GV inoculation. Limited details exist for protists infected with GV and other viruses, including diatoms, flagellates, dinoflagellates and algae. A list of publications mentioning cellular alterations derived from viral infections in protists is shown in the supplementary data file (tab 2). Below, we discuss these alterations in detail for each main host group.

Acanthamoeba castellanii and their broad response to different viruses

Most known GV groups were isolated in *A. castellanii*⁵⁴. A successful giant virus infection triggers the formation of VFs inside the host cells. However, some studies mention cytopathic effects (CPEs) without evident VFs^{55–57}. Nevertheless, GV's DNA replication may also happen in the nucleus, with assembly in cytoplasm⁵⁸. Noumeavirus, for instance, recruits the host's nuclear transcription machinery to the cytoplasmic VF⁵⁹. An overview of some of the main changes using the *A. castellanii* model as an example can be seen in Fig. 2.

Some GV infections can disorganize and deform the nucleus of *A. castellanii*, leading to loss of its spherical appearance, membrane invaginations, or degradation^{55,60–66}. The cytoplasm may undergo significant changes including acidification, organelle rearrangement, cytoskeleton modulation with fragmentation and shortening of microtubules, lower quantity of vacuoles, viral particles present inside vacuoles, formation of Golgi apparatus-like vesicles, among others^{59,61,63,64,67–72} (Fig. 2a).

Viral infection can also modulate the energy of the host, redirecting it to the replication site, often involving the recruitment of mitochondria to VFs^{60,65,73,74}. Mollivirus sibericum infection results in smaller and elongated mitochondria, although host cells did not present changes in cellular adhesion and morphology⁶². Despite that, cell rounding is one common CPE, observed after GV infection (Fig. 2b)^{75–78}. Studies also reported loss of cell adhesion decreased motility, and intercellular bridges or rotational behavior^{55,56,76,79,80}. A peculiar CPE is cellular aggregation ("bunches")^{77,81,82}. This mechanism was suggested to increase viral dissemination during Tupanvirus infection, as mannose-binding protein gene transcripts are significantly increased at earlier times of infection (1, 2 and 4 h post-infection [HPI]) and precedes the formation of bunches (6 HPI) between infected and uninfected cells.

A. polyphaga and *V. vermiformis* morphology modifications upon giant viruses' infection

Fewer studies described CPE in *A. polyphaga* during GV infection. *Acanthamoeba polyphaga* mimivirus (APMV) infection leads to nucleus enlargement, prominent VF formation in the cytoplasm, multivesicular bodies and endoplasmic reticulum (ER) membranes appearance around VF, stimulus of ER synthesis, significant structural alteration in microtubules and actin microfilaments resulting in cells becoming smooth, rounded, and losing motility^{83–85}. Marseillevirus infection induces similar VFs to the observed in *A. castellanii*, with lack of a clear boundary with cytoplasm^{67,86}. Furthermore, cell rounding with formation of cell chains was reported for the first time after a viral infection in amoeba⁸⁷.

Over the past decade, studies implemented *V. vermiformis* as a new isolation platform, leading to the discovery of new GV^{12,13,88–94}. Their infection resulted in cell rounding and loss of adhesion^{88,94}, and cell stretching with increased motility before cell rounding¹². The formation of "blebs" and cell bunches seen in *V. vermiformis*, were previously reported in *A. castellanii*^{12,73,91}. A distinct response was the encystment of neighbor non-infected cells after the exposure to infected cell-released non-protein soluble factors, serving as a defense mechanism, as those cysts become unviable

Table 1 | Overview of the proteomic and transcriptomic studies of different protist organisms infected by their respective viruses

References	Protist organism	Virus	Technique	Time post infection	FoldChange	Down	Up
Rodrigues et al. (2020) ¹³⁷	<i>A. castellanii</i>	Marseillevirus	RNAseq	0, 1, 2, 4, 5, 6, 8, 10, and 12HPI	<−1 and >1	28	19
Legendre et al. (2015) ⁶²	<i>A. castellanii</i>	Mollivirus sibericum	LC-MS/MS	0 to 6HPI	<−2 and >2	38	30
Zhang et al. (2021) ¹³⁸	<i>A. castellanii</i>	Medusavirus	RNAseq	8 to 16HPI	<−0.445 and >0.364	7970	2657
Moniruzzaman et al. (2018) ¹⁶⁰	<i>A. anophagefferens</i>	Aureococcus anophagefferens virus	RNAseq	5 min	<−1.5 and >1.5	588	412
				30 min		≈ 865	≈ 505
				1HPI		82	0
				6HPI		≈ 1060	≈ 690
				12HPI		≈ 1260	≈ 1260
				21HPI		≈ 1430	≈ 1620
Poimala et al. (2022) ¹³⁸	<i>Phytophthora cactorum</i>	PcBV1 & 2	RNAseq	− ^a	<−2 and >2	23	10
			Nanoflow reverse-phase LC-MS	− ^a	<−1.19 and >1.37	17	36
Provenzano et al. (1997) ¹⁷¹	<i>T. vaginalis</i>	TVV	2D	−	−	41	47
He et al. (2017) ¹⁷⁴	<i>T. vaginalis</i>	TVV	iTRAQ labeling	−	<−1 and >1	21	29
Rada et al. (2022) ¹⁷⁵	<i>T. vaginalis</i> (exosome vesicle)	TVV	LFQ-MS	−	<−2 and >2	55	20

"FoldChange" means threshold utilized by each study to determine when a protein or transcript presents increased expression (upregulated - "up") or decreased expression (downregulated - "down"). Other protists are not included on this table since information on the number of deregulated proteins or transcripts was not the focus of the publication or this data was not provided.

^aThe virus was removed from the protist strain, thus there was no time post-infection.

trapping the virus inside⁹⁰. Nuclear alterations of infected *V. vermiformis* include loss of the rounded shape, decrease in surface area, or nucleus disappearance^{12,13,90–94}. Furthermore, clandestinovirus formed its VF within the nucleus⁸⁸. *V. vermiformis*' mitochondrias were also reported to be recruited around VFs^{12,13,90,91,93,94}. Investigating the unique cellular responses in amoebae to various GV is essential for establishing correlations in host-virus interaction and comparing potential CPEs to other types of host cells.

Other protists' morphology and behavior in response to virus inoculation

When exploring protist susceptibility to Tupanvirus beyond amoebae, it was found that the ciliate *Tetrahymena* sp. experience cytotoxic effects in response to the viral presence. The morphological changes included gradual vacuolization, nuclear degradation, loss of motility, and formation of vesicles, similarly to the observed in *A. castellanii*. However, the ciliate does not support viral replication⁶¹.

In contrast to the extensive use of amoeba to study viral infections, there are relatively fewer reports focusing on interactions between viruses and other protists. Among algae protists, the viral infection of *Aureococcus anophagefferens*, a component of ocean blooms, can trigger nucleus and organelle degradation (such as chloroplast) in some infected cells, cell lysis, and loss of the outer polysaccharide glycocalyx during viral infection, even in environmental samples^{95–99}. In addition, viral infection of another bloom-forming protist, *Heterosigma akashiwo*, pointed to nucleus degradation and damaged cell wall¹⁰⁰. When infected by HaNIV virus, cell nucleus presented margination of heterochromatin, but no morphological changes¹⁰¹. Conversely, the infection by HaV01 and HaV triggered cell rounding, and loss of cell motility, together with chloroplast degradation for HaV infection^{102,103}. Interestingly, chloroplasts have been reported to remain intact until cell lysis¹⁰⁴. Vacuolation, disintegrated cytoplasm and ER swelling were reported during HaRNAV infection in *H. akashiwo*¹⁰⁵. Furthermore, in the haptophyte algae *Chrysochromulina parva* and *Prymnesium parvum* the infection by their respective viruses resulted in absence or degradation of cell nucleus^{106,107}.

In the diatoms *Chaetoceros* sp, *Guinardia delicatula* and *Rhizosolenia setigera* the infection by their respective viruses resulted in common CPEs: degradation of chloroplasts and/or photosynthetic pigments^{98,108–113}. Few *Chaetoceros* species additionally exhibited cytoplasm degradation^{108–111,114}, and nucleus or nucleolus degradation was reported in *Chaetoceros salsgineum* and *Guinardia delicatula*^{98,115}. Similarly, chloroplast shrinkage and chlorophyll degradation were suggested during the infection of the phytoplankton coccolithophore *E. huxleyi* by its lytic virus^{116,117}. Nucleus degradation was identified in infected *E. huxleyi* cells, as well as cellular aggregation, likely as a defense strategy to sink infected aggregated cells^{116–118}. Furthermore, mitochondrial damage with vacuolar acidification was described during the infection by EhV99B1¹¹⁹.

Regarding flagellates, the protist *Giardia* sp. has two life cycle stages, a trophozoite and a cyst form^{120,121}. Only its trophozoite stage is susceptible to viral infection, and few morphological changes have been reported for *Giardia canis* and *Giardia lamblia* infected by GCV and GLV, respectively. During GCV infection, the protists presented vacuolization, enlargement of ER, and its cytoplasm became very loose, and during GLV only cell adherence was impaired upon infection^{122,123}. Although, no cell death was reported for both studies, corroborating to the suggestion that giardiaviruses are released from the cell without triggering cell lysis¹²¹. *G. duodenalis*, however, experiences reduction in cell growth without apparent morphological changes¹²⁴. In the free-living flagellate *Bodo saltans* infected by the virus BsV, morphological changes included degraded nucleus and intracellular structures, with lipid vesicles migrating through the VF, while kinetoplast remained intact for longer⁹⁶.

The bloom-forming dinoflagellate *Heterocapsa circularisquama*, associated with red tides and bivalve mortality, can be infected by RNA and DNA viruses. During the DNA virus infection, a VF is formed with granular or fibrous material inside, in addition to organelles disruption^{125–127}. However, the RNA virus infection triggered organelles, nuclear and chlorophyll degradation, and loss of cellular motility^{128–130}. Interestingly, some of the cells were resistant to the RNA virus infection, but not the DNA one, and the non-resistant cells underwent cell lysis^{129,131}. Furthermore, the dsDNA virus

infection triggered damage to *Symbiodinium* cell nucleus initially, and later to organelles¹³². The dinoflagellate *Gymnodinium mikimotoi* shows nuclear degradation and swollen chloroplasts upon viral infection¹³³. Nuclear degradation is also reported to the nonphotosynthetic stramenopile from the genus *Sicyodochytrium* upon viral infection¹³⁴. Overall, relating viral responses across different protist organisms and viruses remains challenging, due to the limited number of studies in this field and the scarcity of available isolates for research.

Metabolic changes in protist cells during virus infection

To investigate cell-virus interactions at the molecular level, different approaches, including transcriptomics and proteomics, have been employed, revealing that each virus modulates their host in a unique manner^{135,136}.

Changes in *Acanthamoeba* sp

Proteomic analysis of *A. castellanii* infected by Mollivirus sibericum identified a few upregulated proteins related to histones, autophagy, DNA synthesis and packaging; and a few downregulated proteins with no clear functional relationship⁶². Together with the fact that this virus is released through exocytosis rather than cell lysis, an overall cellular integrity is suggested throughout the replication cycle⁶². Likewise, few genes were deregulated at the transcriptional level during Marseillevirus infection¹³⁷. Transcripts expression across early, intermediate, and late time periods showed only 48 genes deregulated among these periods (Table 1), with the upregulation of proteins mainly linked to exosome secretion, transfer RNA biogenesis and genetic information processing; and the downregulation of proteins linked to translation apparatus (rRNA related proteins), tRNA encoding genes, carbohydrate metabolism and lipid biosynthesis¹³⁷. For Marseillevirus, amoebal and mitochondrial transcript levels decreased after 1 h post-infection (HPI)¹³⁷.

Contrastingly, Medusavirus infection in *A. castellanii* resulted in thousands of differentially expressed host genes, although mitochondrial gene modulation resembled the pattern of Marseillevirus infection. Only 25% of nuclear host genes, mainly related to ribosome and proteasome pathways, increased its expression between 8 to 16 HPI, suggesting the host to be suffering protein degradation, along with increased viral protein synthesis. The remaining 75% experienced decreased expression during that same period. Gene ontology suggested a reduction in transport activity, although at 48HPI many of these genes were downregulated, in addition to an upregulation of encystment-mediated genes¹³⁸.

Furthermore, Tupanvirus inoculation triggered rRNA shutdown only by the presence of viral particles, unrelated to viral replication⁶¹. Genomic analysis of tupanviruses revealed the involvement of the ribonuclease T2, an enzyme related to the reduction of the physiological activity and phagocytosis capacity in protist hosts¹³⁹.

Bloom-forming and dinoflagellate protists

Infected by lytic and non-lytic viruses, the coccolithophore *E. huxleyi*, is ecologically crucial for marine carbon flux^{140,141}. The correlation between transcriptomics and metabolomics analysis during viral infection indicated that early during infection (1–4 HPI), there was an upregulation of sphingolipid biosynthesis and glycolysis shuffling energy to fatty acids biosynthesis. Later (24HPI), a shift to the activation of the pentose phosphate pathway to produce nucleotides occurred, while glycolysis and fatty acids became downregulated¹¹⁷. Similar results were found for genes related to sphingolipid metabolism, such as the upregulation of dihydroceramide desaturase (DCD) at 6HPI and its downregulation at 45HPI¹⁴². *E. huxleyi* DCD transcript, together with serine palmitoyl transferase (SPT) transcript (another enzyme from sphingolipid biosynthesis pathway) level results pointed to the same tendency of a decrease in these host transcripts production in the environment¹⁴³. Other studies also found lipid modulation in *E. huxleyi* during viral infection, including fatty acids and highly saturated triacylglycerols, and lipid metabolism was suggested to be regulated by modulation of the PI3K-Akt-TOR signaling pathway^{144–152}.

Furthermore, a single cell transcriptomics analysis of *E. huxleyi* during infection revealed an early shutdown of nuclear transcripts, with mitochondrial and chloroplast transcripts initially higher but gradually declined¹⁵³. Indeed, there was a decrease in genes involved in photosynthesis at 6, 12 and 24 HPI, although some cells seemed to be intact and photosynthesizing¹⁴⁸. Enriched functions associated with *E. huxleyi* responding to viral infection included modified amino acid, lipid binding, porin activity, calcium channel activity, pore complex, cell outer membrane, bacterial-type flagellum functions, among others¹⁵⁴. High expression of glycolysis and nucleotides biosynthesis related genes were reported when studying single protist cells from a coccolithophore bloom¹⁵¹. In addition, there was an upregulation of autophagy related genes, with decreased expression of negative regulatory factors, such as PI3K, and an increase of reactive oxygen species (ROS), which is all related to the programmed cell death (PCD) triggered in *E. huxleyi* after infection¹¹⁹. In fact, 20 ROS scavenging genes were impacted after viral infection, and ROS-related genes were found to be increased through transcriptomic analysis^{155,156}. Cell death induced by viral presence could occur before viral particles release, which can be associated with cell autophagy as a defense strategy to avoid viral dissemination, although autophagy is also essential for viral propagation^{117,157}. *E. huxleyi* cell cycle can be affected by virus infection through the modulation of cyclin expression, and host life cycle genes have been found to be impacted during viral takeover of the protist in environment^{149,158,159}. Altogether, these findings suggest substantial cellular changes in *E. huxleyi* after viral encounter, influencing factors related to cell death, modulation of energy, and specific lipids production, alongside a decline in nuclear activity, potentially playing a role in the modulation of oceans blooms¹⁵³.

Another bloom forming protist, *Aureococcus anophagefferens*, exhibited transcripts downregulated at six different time points during viral infection. A massive deregulation started as early as 5 min post-infection and continued after 6HPI. Downregulated proteins indicated the suppression of pathways related to host cytoskeleton formation, photosynthesis, fatty acid metabolism, and carbohydrate biosynthesis. The upregulated group of proteins indicated the activation of host cellular respiration, transcription, protein synthesis, polyamine biosynthesis, and RNA processing. Transcripts related to host defense mechanism were likely suppressed. This dramatic cell modulation indicates the virus–host interaction in a time dependent manner until cell lysis occurs, correlating with the virus's role in brown tide blooms¹⁶⁰.

For *Phaeocystis globosa*, viral infection prevented the accumulation of polyunsaturated fatty acids, decreased protist photosynthetic performance and upregulated mitochondrial respiration, triggered the fragmentation of DNA and activation of caspases, and prevented *P. globosa* to release star-like structures, which in turn affects host carbon assimilation^{161–164}. The correlation among such changes is speculative, suggesting an association with the lysis mechanism triggered in this organism post-viral infection.

Comparing the transcriptomic responses of the dinoflagellates *Symbiodinium tridacnidorum* and *Symbiodinium* C3 under or not UV light exposure (with and without viral replication, respectively), both exhibited upregulated viral transcription and related terms, but only *S. tridacnidorum* triggered downregulation of transcripts related to host-response to viral infection¹⁶⁵.

Protists responsible for causing human or plant disease

After the discovery of *Trichomonas vaginalis* virus (TVV), few studies investigated *Trichomonas vaginalis* cell responses to infection^{166,167}. Initial findings described that most clinical isolates had the presence of a dsRNA icosahedral virus, and that the expression of a major immunogen (P270) on cell surface, and the expression of Ig-degrading proteinases, were directly correlated with its presence^{168–170}. An overview of the protein expression pattern of *T. vaginalis* after TVV infection found at least 47 expressed and 41 suppressed proteins linked to the virus¹⁷¹. Isolates of *T. vaginalis* might be infected by up to 4 different strains of TVV, and only the cells infected by TVV2 and TVV3 are capable of inducing P270 protein expression^{172,173}.

Proteomic studies comparing infected and uninfected cells detected the upregulation of adhesin proteins related to the pathogenicity of *T. vaginalis* when infecting humans¹⁷⁴. Further, metabolic enzymes, ribosomal and heat shock proteins were also differentially impacted¹⁷⁵. Extracellular vesicles of *T. vaginalis* infected by TVV exhibit differences in protein content, such as a protein responsible for increasing adherence, suggesting enhanced exosome binding and pathogenicity¹⁷⁵. Overall, TVV plays an important role in *T. vaginalis* dynamics by modulating different genes that aid the protist to successfully establish in its host.

After the discovery of endosymbiotic viruses in *Leishmania*'s cytoplasm, most studies focused on the relation *Leishmania*'s pathogenicity to humans when infected or not by one of its two viruses, LRV and LBV^{176–178}. The metastasizing *L. Viannia* had a higher amount of LRV1, suggesting that the virus triggers parasitic resistance¹⁷⁹. A study looked for transcriptional changes in *L. guyanensis* and *L. major* when removing their respective viruses LRV1 and LRV2¹⁸⁰. As a result, *L. guyanensis* did not suffer significant changes, with only 2 differentially expressed genes. On the other hand, proliferation rate of *L. major* cells decreased, with 67 upregulated and 20 downregulated transcripts after its virus removal. Among the downregulated group, there was a cyclin related to growth kinetics; and the membrane-bound acid phosphatase 2 enzyme, which has been already suggested to affect the biology of *Leishmania*. The upregulated group presented transcripts related to autophagy, cell response to various stimuli, and nucleosome assembly¹⁸⁰.

Oomycetes from the genus *Phytophthora* are responsible for causing great damage to agriculture¹⁸¹. A metagenomic study of *Phytophthora condilina* resulted in 15 different putative viruses identified¹⁸². Moreover, GV sequences have been identified in the genome of *Phytophthora parasitica*¹⁸³. *Phytophthora* endornavirus 2 (PEV2) and *Phytophthora* endornavirus 3 (PEV3), are suggested to stimulate zoosporangium development and inhibit hyphal growth, also reducing the host oomycete sensitivity to the antifungal metalaxyl¹⁸⁴. In addition, seven different viruses were able to replicate in *P. infestans*, and viral infection apparently did not affect the cell host morphology but stimulated the growth of the mycelium mass¹⁸⁵. Notably, the infection of *P. infestans* by PiRV-2 is suggested to increase the protist sporulation, which in turn is associated with a likely hypervirulent factor against its plant host¹⁸⁶. However, sporulation of *P. cactorum* infected by PcBV1 & 2, seemed deeply affected when compared to *P. cactorum* with PcBV1 & 2 ablation. Its sporangia production and size decreased during viral infection, together with reduced hyphal growth¹⁸⁷. This same research used *P. cactorum* as a model to explore the transcriptomic and proteomic changes after removing PcBV1 & 2 from the protist, and identified 10 up- and 23 downregulated transcripts, as well as 36 up- and 17 downregulated proteins¹⁸⁷. The excreted protein elicitor was found upregulated in the infected cells¹⁸⁷. This protein is related to the suppression of the plant immune response but can also be recognized by the plant and become a factor that reduces protist pathogenicity^{188,189}. Overall, *Phytophthora* can be pathogenic or not to its plant host depending on many factors, including the infecting virus type.

Cellular resistance strategies after stress exposure

All living organisms are subjected to the exposure of extrinsic factors that can trigger a stress response at the cellular level. Stress refers to any environmental condition that can be harmful to cells and induce physiological changes that disturb homeostasis¹⁹⁰. Biotic stress is caused by interaction with other living beings that will act as stressors, particularly parasitic relationships involving viruses, bacteria, and fungi. Abiotic stress arises from physicochemical factors that can affect cell physiology, such as pH, temperature, radiation, osmolarity and chemical molecules^{190,191}. Despite its disruption of homeostasis, environmental stress is an important factor when it comes to evolution. Cells capable of restoring homeostasis after stress, called acclimatized cells, are favored through natural selection¹⁹⁰. Protist cells employ various responses to stressors programmed cell death (PCD). One example is the response of *Peridinium gatunense* to oxidative stress. At the end of the algal blooms caused by this dinoflagellate, CO₂ becomes

significantly limited, leading to the production of reactive oxygen species which trigger PCD¹⁹². *Aureococcus anophagefferens*, responds to stress conditions such as low concentrations of inorganic nitrogen and inorganic phosphorus through a transcriptional shift: transcripts related to nitrogen transport and metabolism, and transcripts encoding enzymes that hydrolyze organic phosphorus or alleviate arsenic toxicity are upregulated in these scenarios. In addition, in the context of low light levels, which acts as a stress factor, transcripts encoding enzymes that catabolize organic compounds, restructure lipid membranes, or are involved in the biosynthesis of sulfolipids are upregulated to restructure lipids and renovate the photosynthetic apparatus. The cell undergoes several physiological changes to maintain the survival and ecological success of the species¹⁹³.

Spore formation as a response to infection

The marine diatom *Chaetoceros socialis* responds to stress caused by adverse conditions of temperature, light and lack of nutrients through spore formation. Viruses are widely distributed and persistent biological stressors, and some of them are also capable of inducing spore formation¹⁹⁴. For instance, CsfRNAV01 can effectively induce significant spore formation in *C. socialis*. Viral infection acts as a biotic trigger that induces a substantial formation of heavily silicified spores. Interestingly, their spores do not produce infectious viral particles, making this shift in the life cycle an effective defense strategy against viruses and preventing the loss of a portion of the protist population¹⁹⁴.

Emiliania huxleyi life cycle shift as a defense strategy

Another protist that relies on a life cycle shift is *E. huxleyi*, a photosynthetic organism widely distributed throughout the oceans. It has two independent phases, haploid and diploid, each with distinct morphologies¹⁹⁵. The haploid form (N) consists of biflagellated cells covered by thin, organic, non-mineralized scales. In contrast, the diploid (2N) is nonmotile and contains minute calcite platelets, named coccoliths¹⁹⁶. The latter is responsible for large natural algal blooms, an important environmental problem¹⁹⁷. It is known that calcium plays an important role during the viral entry process into the host cell in different infection models^{198–201}. One significant difference between the N and 2N phases of *E. huxleyi* is the calcium metabolism being considerably greater in the diploid phase than in the haploid phase, affecting virus-host interactions¹⁹⁷. Haploid cells are resistant to infection and subsequent lysis by *Emiliania huxleyi* virus, whereas this infection contributes to the decline of diploid populations during algal blooms¹⁹⁷. Evidence suggests that those lytic viruses are responsible for demising the algae population and terminating algae blooms. Thus, *Emiliania huxleyi* viruses are responsible for regulating the population of this coccolithophore both in abundance of cells in the environment and in composition^{195,197,202,203}. Such protist-virus interaction can directly activate the life cycle transition, since the oxidative stress caused by viral infection can trigger the diploid-to-haploid shift¹⁹⁷. Viral infections are known to increase the production of reactive oxygen species in 2N cells, which induces the activation of metacaspases and, thus PCD is triggered²⁰⁴. However, oxidative stress response leads not only to cell death but also to induction of the aforementioned shift¹⁹⁶. Although studies conducted using 1N *E. huxleyi* strains found viral RNA and small amounts of EhV glycosphingolipids within the cells^{146,205}, a later study using the host resistant strains have not found any EhV genetic material, which suggests that there is no resistant strain bearing any form of the virus¹⁹⁵. Haploid virus-resistant cells are produced as a response to viral infection¹⁹⁵. This is referred to as the 'Cheshire Cat' strategy, in which the organism escapes the parasite by shifting to a life stage that is infection-resistant¹⁹⁷.

Encystment as amoebae' resistance form against viral infections

The 'Cheshire Cat' theory extends to describe a relationship between a giant virus and an amoeba as well. Some amoeba hosts are also capable of shifting to an infection-resistant life form²⁰⁶. Amoebae have two life cycle stages, trophozoites and cysts. The trophozoite stage is the one that predominates when environmental conditions are favorable, such as nutrient supply and




Fig. 3 | Overview of the strategies employed by different protist life forms to resist stress factors, such as virus infection. **a** Upon interaction with viruses, the cocolithophore *E. huxleyi* shifts to a life form that is resistant to infection¹⁹⁵. **b** Amoebae encyst under unfavorable conditions, and the cysts formed are not able to be infected

by a virus²⁰⁶. **c** Under unfavorable conditions *Chaetoceros socialis* cells transform into resting spores that are unable to produce infectious viral particles^{114,194,225}. Red text: Cellular resistance strategies after stress exposure.

temperature, while the cyst is the form of resistance^{191,206}. Under unfavorable conditions such as starvation, or osmotic stress, amoeba trophozoites undergo both cellular and molecular regulatory processes that lead to encystment^{206–208}. Amoeba cysts are more resistant to adverse conditions. For instance, *A. castellanii* cysts can survive five cycles of freeze–thawing, exposure to high doses of UV or gamma radiation²⁰⁹. Furthermore, giant viruses, such as mimiviruses and marseilleviruses, can only infect amoeba when they are in the trophozoite form. The shift to a non-permissive cell stage is an astounding stratagem for escaping giant viruses²⁰⁶. Some viruses, however, have developed strategies to circumvent this resistance mechanism. Mimivirus isolates and Tupanvirus, can prevent the encystment of amoebae, by reducing the expression of amoebal encystment-mediating subtilisin-like serine proteinase, a proteinase involved in trophozoite encystment. By doing so, the virus can effectively block the formation of the host's form of resistance and continue its cycle^{90,206,210}.

The tripartite systems

The tripartite systems are an interesting dynamic in which the giant virus–host relationship is accompanied by a further member: a virophage or a bacteria. Such tripartite systems are found, for instance in amoebae. Virophages are viruses capable of infecting giant viruses, they multiply inside the VFs, meaning that they do not replicate if giant viruses are absent. Therefore, the replication of the virophages interferes with the cycle of the giant virus, interrupting it and thus reducing the amoeba mortality related to the viral infection^{79,211}. The first virophage discovered was sputnik, a dsDNA virus associated with mamavirus. Some types of virophages increase the formation of abnormal giant virus particles, reduce infectivity and the ability to lyse their host cell²¹¹. The second virophage discovered was the mavirus associated with the Cafeteria roenbergensis virus (CroV). Like the sputnik, this virophage inhibits the cycle of the giant virus. Interestingly, mavirus can integrate its genome into the protist's genome, but its genes are only expressed when CroV infection occurs. This reactivation is not enough to prevent cell lysis; however, it causes mavirus particles to be released into the environment and protect neighboring cells^{79,212,213}. Despite this, it has been shown that not all virophages negatively affect the cycle of giant viruses; zamilon, for instance, does not affect the ability of the giant virus to lyse the amoeba²¹⁴. There are also bacterial endosymbionts capable of protecting

amoebae from being infected by giant viruses, such as *Parachlamydia acanthamoebae*, which manages to suppress the viral replication of sympatric Viennavirus, APMV and Tupanvirus by inhibiting the maturation of the VF, which results ultimately in the survival of the infected amoebae²¹⁵. Such complex systems are also present in other protists, such as *Cryptomonas* sp. SAG25.80, which is a quadripartite system²¹⁶. A single cell harbors a phage (MAnkyphage) and two bacterial endosymbionts (*Grellia numerosa* and *Megaira polyxenophila*), and a complex community of organelles and selfish elements²¹⁶. As discussed above, protist organisms have developed various strategies throughout their evolutionary history to survive unfavorable environmental conditions and escape viral infection. A graphical representation of the main strategies discussed is shown by Fig. 3.

Summary and concluding remarks

Early eukaryotes are believed to have emerged more than a billion years ago in the Proterozoic oceans²¹⁷. Despite the ongoing efforts to characterize and explore the diversity of viruses that infect protists, few isolates have been described to date, and the majority of the known diversity was discovered through metagenomic studies. One of the most notable groups of viruses, with the higher number of viral isolates, are the representatives of the phylum *Nucleocytoviricota*, an ancient component of the eukaryotic virome^{32,218}. Each of these organisms have undergone extensive co-evolution, resulting in their present-day diversity, which encompasses a broad array of eukaryotes as hosts, and different taxa of viruses as parasites⁴⁸. Even though certain viruses of this phylum possess a highly diverse set of genes involved in different metabolic pathways, their obligated intracellular parasitic nature necessitates an intricate relationship with their hosts.

Viruses often disrupt normal cellular processes to acquire the necessary elements for viral replication, or as a result of dysbiosis originated from the infection process. The intensification of studies on viral–protists host interactions is enhancing our understanding of the molecular mechanisms underlying their interactions, shedding light on viral replication and host defense mechanisms. Following viral infection, host cells often go through significant alterations and begin to exhibit a distinct nature⁴⁹. Some of the alterations can be observed microscopically and are considered as an indicative of viral infection. These particular alterations, known as cytopathic effects, typically comprise morphological changes that may manifest as

noticeable variations in cell shape, size and integrity. In the case of protists, it has been observed that these changes can include increased cellular motility, rounding of cells, reduced adhesion, or increased adhesion to other cells, and cell lysis^{12,80–83,85–92}. Furthermore, morphological modifications are accompanied by intracellular alterations. During a viral infection, intracellular changes in protists cells include structural transformations of the cytoskeleton, degradation of chloroplasts or photosynthetic pigments (if present), recruitment of mitochondria, membranes, vesicles, etc. Even if the VF is exclusively formed in the cytoplasm and the nucleus remains intact throughout viral replication, it can still go through changes, albeit temporary, during the course of the infection^{12,66,109,117,128,133,219,220}. In addition, viral infections induce functional changes in host cells through widespread genetic and metabolic reprogramming. Apart from inducing the expression of viral genes, viruses can elevate the expression of host genes associated with processes such as DNA synthesis, energy generation, and the modification of lipid, protein, and nucleic acid biosynthesis pathways. These alterations collectively contribute to the complex interplay between viruses and host cells during infection^{62,117,135–138,140,144,153,154,167}. Still, deeper exploration of virus-host dynamics is much needed, and can help elucidate host factors influencing susceptibility, pathogenicity, resistance, and protists basic cytology.

As aforementioned, protists are a diverse group of eukaryotic organisms that exhibit a remarkable adaptability when exposed to harsh conditions and invasive encounters. To thrive in ever-changing environments and fend off predators and parasites, protists have evolved a repertoire of sophisticated cellular responses to stress, often including the formation of resistant life forms. Moreover, as they occupy a wide range of ecological niches, protists have a dynamic interplay with other organisms with which they form communities, including viruses. They can act as regulators of population diversity and density, as in the case of *Emiliania huxleyi* viruses, which are responsible for ending coccolithophorid blooms^{195,197}. Furthermore, the interaction between viruses and protists can also be very complex, as in tripartite systems, in which the virus-host relationship involves another member which may interfere in the replication cycle of the giant virus and protect amoebae from infection^{79,212,213}. Competitive interactions are one of the main driving forces that lead to the diversity and complexity of life on our planet, and the investigation into how viruses influence protists communities, and their ecological interactions can help understand how virus-driven evolution shapes the diversity, dynamics and the impacts in ecosystems, including their role in nutrient cycling and energy flow.

Many are the taxonomic groups of viruses that infect protists, and together, they are present in the majority of Earth's ecosystems. One outcome of such interactions is horizontal gene transfer. Viruses might act as agents for transferring genetic material across species boundaries, including protists, while certain species are hot spots for horizontal gene transfer (HGT) among viruses, eukaryotes, prokaryotes and mobile elements^{160,221}. This process has the potential to introduce novel genes or traits into protist genomes, and potentially influence their biology, adaptation, and evolution²²². Furthermore, it contributes to studies on the evolution of protists and its viruses, providing insights into the coevolutionary dynamics, and as these organisms often have unique genomic features with a high percentage of ORFan genes⁶⁰, the study of the viral genomes can uncover novel genes and genetic features that may be noteworthy in protist hosts. This can expand our understanding of genetic diversity, the impact of viral genes on protists' biology, ecology and evolution, as well as potential functions within the protists.

Protists and viruses play a crucial role in Earth's ecosystems. Understanding how viruses impact protist biology can provide valuable insights into several fields of science, shedding light on evolutionary biology, ecosystem dynamics, nutrient cycling, and biomedical research. However, our knowledge of unicellular organisms and microorganisms known to humankind remains anthropocentric, with little known about the true diversity of these beings. Therefore, it is important that prospecting and characterization studies of these organisms are carried out to better elucidate the complete picture. In summary, viruses can serve as valuable tools and

subjects of study in the field of protist biology. Their interactions with protist hosts, unique genomes, and ecological roles can offer new information that enhances our understanding of protist biology and its broader implications for ecosystems and biotechnology.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

No datasets were generated or analysed for this review paper. Literature cited is shown in the reference list and in the Supplementary Data File.

Received: 28 October 2023; Accepted: 29 February 2024;

Published online: 11 March 2024

References

- Owen, R. Kingdom *Protozoa in Paleontology*. 1–17 (Edinburgh, Colin Macfarquhar & Andrew Bell, 1858).
- Cavalier-Smith, T. *Kingdom Protozoa and Its 18 Phyla. Microbiol. Rev.* **57**, 953–994 (1993).
- Jahn, T. L. & Votta, J. J. Locomotion of protozoa. *Annu. Rev. Fluid Mech.* **4**, 93–116 (1972).
- Adl, S. M. et al. The new higher level classification of eukaryotes with emphasis on the taxonomy of protists. *J. Eukaryotic Microbiol.* **52**, 399–451 (2005).
- Caron, D. A., Worden, A. Z., Countway, P. D., Demir, E., & Heidelberg, K. B. Protists are microbes too: a perspective. *ISME J.* **3**, 4–12 (2009).
- Adl, S. M. et al. Diversity, nomenclature, and taxonomy of protists. *Syst. Biol.* **56**, 684–689 (2007).
- Brusca, R., Shuster, S. & Moore, W. *Invertebrates* (2016).
- da Silva Ferreira, M., de Souza Gonçalves, D., Medeiros, E. G., Peralta, J. M. & Guimarães, A. J. “Feast-Fit-Fist-Feat”: overview of free-living amoeba interactions with fungi and virulence as a foundation for success in battle. *Curr. Trop. Med. Rep.* **8**, 18–31 (2021).
- Mungroo, M. R., Siddiqui, R. & Khan, N. A. War of the microbial world: *Acanthamoeba* spp. interactions with microorganisms. *Folia Microbiologica* **66**, 689–699 (2021).
- Queiroz, V. F. et al. Amoebae: hiding in plain sight: unappreciated hosts for the very large viruses. *Annu. Rev. Virol.* **9**, 79–98 (2022).
- Korn, E. D. & Weisman, R. A. Phagocytosis of latex beads by *Acanthamoeba*. II. Electron microscopic study of the initial events. *J. Cell Biol.* **34**, 219–227 (1967).
- Souza, F. et al. In-depth analysis of the replication cycle of Orpheovirus. *Virol. J.* **16**, 158 (2019).
- Silva, L. C. F. et al. Microscopic analysis of the tupanvirus cycle in *Vermamoeba vermiformis*. *Front. Microbiol.* **10**, 671 (2019).
- Andrade, A.C.S.P. et al. Filling knowledge gaps for mimivirus entry, uncoating, and morphogenesis. *J. Virol.* **91**, e01335-17 (2017).
- Arantes, T. S. et al. The large marseillevirus explores different entry pathways by forming giant infectious vesicles. *J. Virol.* **90**, 5246–5255 (2016).
- Shi, Y. et al. The ecology and evolution of amoeba-bacterium interactions. *Appl. Environ. Microbiol.* **87**, e01866-20 (2021).
- Scola, B. L. et al. A giant virus in amoebae. *Science* **299**, 2033 (2003).
- Horn, M. Bacterial endosymbionts of free-living amoebae. *J. Eukaryot. Microbiol.* **51**, 509–514 (2004).
- Samba-Louaka, A., Delafont, V., Rodier, M. H., Cateau, E. & Héchar, Y. Free-living amoebae and squatters in the wild: ecological and molecular features. *FEMS Microbiol. Rev.* **43**, 415–434 (2019).
- de Souza, G. A. P., Queiroz, V. F., Coelho, L. F. L. & Abrahão, J. S. Alohomora! What the entry mechanisms tell us about the evolution

- and diversification of giant viruses and their hosts. *Curr. Opin. Virol.* **47**, 79–85 (2021).
21. Kao, S., Kao, C.-F., Chang, W. & Ku, C. Widespread distribution and evolution of poxviral entry-fusion complex proteins in giant viruses. *Microbiol. Spectr.* **11**, e0494422 (2023).
22. De la Fuente, I. M. et al. The nucleus does not significantly affect the migratory trajectories of amoeba in two-dimensional environments. *Sci. Rep.* **9**, 16369 (2019).
23. DuBose, J. G., Robeson, M. S., Hoogshagen, M., Olsen, H. & Haselkorn, T. S. Complexities of inferring symbiont function: paraburkholderia symbiont dynamics in social amoeba populations and their impacts on the amoeba microbiota. *Appl. Environ. Microbiol.* **88**, e0128522 (2022).
24. De la Fuente, I. M. & López, J. I. Cell motility and cancer. *Cancers* **12**, 2177 (2020).
25. Jeon, K. W. Nuclear control of cell movement in amoebae: nuclear transplantation study. *Exp. Cell Res.* **50**, 467–471 (1967).
26. Lhee, D. et al. Amoeba genome reveals dominant host contribution to plastid endosymbiosis. *Mol. Biol. Evol.* **38**, 344–357 (2021).
27. Schulz, F. et al. Life in an unusual intracellular niche: a bacterial symbiont infecting the nucleus of amoebae. *ISME J.* **8**, 1634–1644 (2014).
28. Shu, L. et al. Symbiont location, host fitness, and possible coadaptation in a symbiosis between social amoebae and bacteria. *Elife* **7**, 1–25 (2018).
29. Thomas, V. & Greub, G. Amoeba/amoebal symbiont genetic transfers: lessons from giant virus neighbours. *Intervirology* **53**, 254–267 (2010).
30. Enquist, L. W. Virology in the 21st century. *J. Virol.* **83**, 5296–5308 (2009).
31. Wang, A. L. & Wang, C. C. Viruses of parasitic protozoa. *Parasitol. Today* **7**, 76–80 (1991).
32. Krupovic, M., Dolja, V. V. & Koonin, E. V. The virome of the last eukaryotic common ancestor and eukaryogenesis. *Nat. Microbiol.* **8**, 1008–1017 (2023).
33. Oldstone, M. B. A. & Levine, A. J. Virology in the next millennium minireview achievements led to the eradication of viral diseases. *Cell* **100**, 139–142 (2000).
34. Hershey, A. D. & Chase, M. Independent functions of viral protein and nucleic acid in growth of bacteriophage*. *J. Gen. Physiol.* **36**, 39–56 (1952).
35. Hendrix, R. W. There 1031 virus particles on earth, or more, or fewer? *Proc. Natl Acad. Sci. USA* **96**, 2192–2197 (1999).
36. Li, T. et al. CRISPR/Cas9 therapeutics: progress and prospects. *Sig. Transduct. Target. Ther.* **8**, 36 (2023).
37. Rodrigues, R. A. L. et al. An anthropocentric view of the virosphere-host relationship. *Front. Microbiol.* **8**, 1673 (2017). **Demonstration that the known virosphere is extremely anthropocentric and that viruses that infect protists are among the least described ones.**
38. Suttle, C. A. Viruses in the sea. *Nature* **437**, 356–361 (2005).
39. Miller, J. H. & Swartzwelder, J. C. Virus-like particles in an *Entamoeba histolytica* Trophozoite. *Source: J. Parasitol.* **46**, 523 (1960).
40. Attoui, H. et al. Micromonas pusilla reovirus: a new member of the family *Reoviridae* assigned to a novel proposed genus (Mimoreovirus). *J. General Virol.* **87**, 1375–1383 (2006).
41. Coy, S., Gann, E., Pound, H., Short, S. & Wilhelm, S. Viruses of eukaryotic algae: diversity, methods for detection, and future directions. *Viruses* **10**, 487 (2018).
42. Diamond, L. S. & Mattern, C. F. T. Protozoal viruses. *Adv. Virus Res.* **20**, 87–112 (1976).
43. Fukuhara, T. Endornaviruses: persistent dsRNA viruses with symbiotic properties in diverse eukaryotes. *Virus Genes* **55**, 165–173 (2019).
44. Khramtsov, N. V. & Upton, S. J. Association of RNA polymerase complexes of the parasitic protozoan *Cryptosporidium parvum* with virus-like particles: heterogeneous system †. *J. Virol.* **74**, 5788–5795 (2000).
45. Malathi, V. G. & Renuka Devi, P. ssDNA viruses: key players in global virome. *Virus Dis.* **30**, 3–12 (2019).
46. Nibert, M. L., Woods, K. M., Upton, S. J. & Ghabrial, S. A. Cryspovirus: a new genus of protozoan viruses in the family *Partitiviridae*. *Arch. Virol.* **154**, 1959–1965 (2009).
47. Short, S. M., Staniewski, M. A., Chaban, Y. V., Long, A. M. & Wang, D. Diversity of viruses infecting eukaryotic algae. *Curr. Issues Mol. Biol.* **39**, 29–62 (2020).
48. Koonin, E. V. et al. Global organization and proposed megataxonomy of the virus world. *Microbiol. Mol. Biol. Rev.* **84**, e00061–19 (2020).
49. Forterre, P. The virocell concept and environmental microbiology are the great virus comeback. *ISME J.* **7**, 233–236 (2013). **Emphasizes the dynamics and importance of viral infections in nature.**
50. De Castro, I. F., Volonté, L. & Risco, C. Virus factories: biogenesis and structural design. *Cell. Microbiol.* **15**, 24–34 (2013).
51. Bisio, H. et al. Evolution of giant pandoravirus revealed by CRISPR/Cas9. *Nat Commun* **14**, 428 (2023).
52. La Scola, B. Looking at protists as a source of pathogenic viruses. *Microb. Pathog.* **77**, 131–135 (2014).
53. Diesend, J., Kruse, J., Hagedorn, M. & Hammann, C. Amoebae, giant viruses, and virophages make up a complex, multilayered threesome. *Front. Cell. Infect. Microbiol.* **7**, 527 (2018).
54. Machado, T. B., de Aquino, I. L. M. & Abrahão, J. S. Isolation of giant viruses of *Acanthamoeba castellanii*. *Curr. Protoc.* **2**, e455 (2022).
55. Philippe, N. et al. Pandoraviruses: Amoeba viruses with genomes up to 2.5 Mb reaching that of parasitic eukaryotes. *Science (1979)* **341**, 281–286 (2013).
56. Alempic, J. M. et al. An update on eukaryotic viruses revived from ancient permafrost. *Viruses* **15**, 564 (2023). **Isolation and characterization of 13 viruses from siberian permafrost samples.**
57. Akashi, M. & Takemura, M. Co-isolation and characterization of two pandoraviruses and a mimivirus from a riverbank in Japan. *Viruses* **11**, 1123 (2019).
58. Yoshikawa, G. et al. Medusavirus, a novel large DNA virus discovered from hot spring water. *J. Virol.* **93**, e02130–18 (2019).
59. Fabre, E. et al. Noumeavirus replication relies on a transient remote control of the host nucleus. *Nat. Commun.* **8**, 15087 (2017).
60. Boratto, P. V. M. et al. Yaravirus: a novel 80-nm virus infecting *Acanthamoeba castellanii*. *PNAS* **117**, 16579–16586 (2020).
61. Abrahão, J. et al. Tailed giant Tupanvirus possesses the most complete translational apparatus of the known virosphere. *Nat. Commun.* **9**, 749 (2018).
62. Legendre, M. et al. In-depth study of Mollivirus sibericum, a new 30,000-yold giant virus infecting *Acanthamoeba*. *Proc. Natl Acad. Sci. USA* **112**, E5327–E5335 (2015).
63. Queminn, E. R. et al. Complex membrane remodeling during virion assembly of the 30,000-year-old mollivirus sibericum. *J. Virol.* **93**, e00388–19 (2019).
64. Christo-Foroux, E. et al. Characterization of Mollivirus kamchatka, the first modern representative of the proposed Molliviridae family of giant viruses. *J. Virol.* **94**, e01997–19 (2020).
65. Pereira Andrade, A.C.S. et al. New isolates of pandoraviruses: contribution to the study of replication cycle steps. *J. Virol.* **93**, e01942–18 (2019).
66. Fukaya, S., Masuda, L. & Takemura, M. Analysis of morphological changes in the nucleus and vacuoles of *Acanthamoeba castellanii* following giant virus infection. *Microbiol. Spectr.* **11**, e0418222 (2023). **Changes on the nucleus, vacuoles and A. castellanii cell shape are triggered during four different giant virus infection.**

67. Goyal, N., Barai, A., Sen, S. & Kondabagil, K. Amoebal tubulin cleavage late during infection is a characteristic feature of mimivirus but not of marseillevirus. *Microbiol. Spectr.* **10**, e0275322 (2022).
68. Legendre, M. et al. Thirty-thousand-year-old distant relative of giant icosahedral DNA viruses with a pandoravirus morphology. *PNAS* **111**, 4274–4279 (2014).
69. Legendre, M. et al. Diversity and evolution of the emerging *Pandoraviridae* family. *Nat. Commun.* **9**, 2285 (2018).
70. Takemura, M. Morphological and Taxonomic properties of Tokyovirus, the first Marseilleviridae member isolated from Japan. *Microbes Environ.* **31**, 442–448 (2016).
71. Bertelli, C. et al. *Cedratvirus lausannensis* – digging into *Pithoviridae* diversity. *Environ. Microbiol.* **19**, 4022–4034 (2017).
72. Doutre, G., Philippe, N., Abergel, C. & Claverie, J.-M. Genome analysis of the first Marseilleviridae representative from Australia indicates that most of its genes contribute to virus fitness. *J. Virol.* **88**, 14340–14349 (2014).
73. Silva, L. K. S. et al. Cedratvirus getuliensis replication cycle: an in-depth morphological analysis. *Sci. Rep.* **8**, 4000 (2018).
74. Thomas, V. et al. Lausannevirus, a giant amoebal virus encoding histone doublets. *Environ. Microbiol.* **13**, 1454–1466 (2011).
75. Campos, R. K. et al. Samba virus: a novel mimivirus from a giant rain forest, the Brazilian Amazon. *Virol. J.* **11**, 95 (2014).
76. Arslan, D., Legendre, M., Seltzer, V., Abergel, C. & Claverie, J. M. Distant Mimivirus relative with a larger genome highlights the fundamental features of *Megaviridae*. *Proc. Natl Acad. Sci. USA* **108**, 17486–17491 (2011).
77. Aoki, K. et al. *Marseilleviridae* lineage b diversity and bunch formation inhibited by galactose. *Microbes Environ.* **36**, n/a (2021).
78. Scheid, P. L. New insights into the interaction of free-living amoebae and pandoravirus inopinatum: investigations of the host range and the role of multilamellar bodies. *Open Parasitol. J.* **6**, 63–74 (2018).
79. Oliveira, G., La Scola, B. & Abrahão, J. Giant virus vs amoeba: fight for supremacy. *Virol. J.* **16**, 126 (2019).
80. Fukaya, S. & Takemura, M. Kinetic analysis of *Acanthamoeba castellanii* infected with giant viruses quantitatively revealed process of morphological and behavioral changes in host cells. *Microbiol. Spectr.* **9**, e0036821 (2021). **Time lapse videos identified cell changes, such as rounding, and behaviour, such as rotation, in A. castellanii after infection of different giant viruses.**
81. Oliveira, G. et al. Tupanvirus-infected amoebas are induced to aggregate with uninfected cells promoting viral dissemination. *Sci. Rep.* **9**, 183 (2019).
82. Aoki, K. et al. Fifteen marseilleviruses newly isolated from three water samples in Japan reveal local diversity of marseilleviridae. *Front. Microbiol.* **10**, 1152 (2019).
83. Suzan-Monti, M., Scola, B. L., Barrassi, L., Espinosa, L. & Raoult, D. Ultrastructural characterization of the giant volcano-like virus factory of *Acanthamoeba polyphaga* Mimivirus. *PLoS One* **2**, e328 (2007).
84. Mutsafi, Y., Shimon, E., Shimon, A. & Minsky, A. Membrane assembly during the infection cycle of the giant mimivirus. *PLoS Pathog.* **9**, e1003367 (2013).
85. Yaakov, L. B., Mutsafi, Y., Porat, Z., Dadosh, T. & Minsky, A. Kinetics of mimivirus infection stages quantified using image flow cytometry. *Cytometry Part A* **95**, 534–548 (2019).
86. Boyer, M. et al. Giant Marseillevirus highlights the role of amoebae as a melting pot in emergence of chimeric microorganisms. *PNAS* **106**, 21848–21853 (2009).
87. Dos Santos, R. N. et al. A new marseillevirus isolated in Southern Brazil from *Limnoperna fortunei*. *Sci. Rep.* **6**, 35237 (2016).
88. Rolland, C. et al. Clandestinovirus: a giant virus with chromatin proteins and a potential to manipulate the cell cycle of its host *Vermamoeba vermiformis*. *Front. Microbiol.* **12**, 715608 (2021).
89. Bajrai, L. et al. Kaumobavirus, a new virus that clusters with Faustoviruses and Asfarviridae. *Viruses* **8**, 278 (2016).
90. Borges, I. et al. Trapping the Enemy: *Vermamoeba vermiformis* circumvents faustovirus mariensis dissemination by enclosing viral progeny inside cysts. *J. Virol.* **93**, e00312–19 (2019).
91. Andreani, J. et al. Orpheovirus IHUMI-LCC2: a new virus among the giant viruses. *Front. Microbiol.* **8**, 2643 (2018).
92. Andreani, J. et al. Morphological and genomic features of the new Klosneuvirinae isolate fadovirus IHUMI-VV54. *Front. Microbiol.* **12**, 719703 (2021).
93. Bajrai, L. H. et al. Isolation of Yasminevirus, the first member of Klosneuvirinae isolated in coculture with *Vermamoeba vermiformis*, demonstrates an extended arsenal of translational apparatus components. *J. Virol.* **94**, e01534–19 (2019).
94. Reteno, D. G. et al. Faustovirus, an Asfarvirus-related new lineage of giant viruses infecting amoebae. *J. Virol.* **89**, 6585–6594 (2015).
95. Popovic, M., Samgadharan, M. G., Read, E. & Gallo, R. C. Isolation of virus capable of lysing the brown tide microalga, *Aureococcus Anophagefferens*. *Biophys. Res. Commun.* **224**, 497–500 (1984).
96. Gastrich, M. D. et al. Ultrastructural analysis of viral infection in the brown-tide alga, *Aureococcus Anophagefferens* (Pelagophyceae). *Phycologia* **37**, 300–306 (1998).
97. Gastrich, M. D. et al. Viruses as potential regulators of regional brown tide blooms caused by the alga, *Aureococcus anophagefferens*. *Estuaries* **27**, 112–119 (2004).
98. Gastrich, M. D., Anderson, O. R. & Cosper, E. M. Viral-like particles (VLPs) in the alga, *Aureococcus anophagefferens* (pelagophyceae), during 1999–2000 brown tide blooms in Little Egg Harbor, New Jersey. *Estuarine Res. Federation Estuaries* **25**, 938–943 (2002).
99. Gobler, C., Anderson, O., Gastrich, M., Mary Downes, G. & Wilhelm, S. Ecological aspects of viral infection and lysis in the harmful brown tide alga *Aureococcus anophagefferens*. *Aquatic Microbial Ecol.* **47**, 25–36 (2007).
100. Nagasaki, K., Ando, M., Itakura, S., Imai, I. & Ishida, Y. Viral mortality in the final stage of *Heterosigma Akashiwo* (Raphidophyceae) red tide. *J. Plankton Res.* **16**, 1595–1599 (1994).
101. Lawrence, J. E., Chan, A. M. & Suttle, C. A. A novel virus (HaNIV) causes lysis of the toxic bloom-forming alga *Heterosigma akashiwo* (Raphidophyceae) 1. *J. Phycol.* **37**, 216–222 (2001).
102. Nagasaki, K., Tarutani, K. & Yamaguchi, M. Growth characteristics of heterosigma Akashiwo virus and its possible use as a microbiological agent for red tide control. *Appl. Environ. Microbiol.* **65**, 898–902 (1999).
103. Nagasaki, K. & Yamaguchi, M. Isolation of a virus infectious to the harmful bloom causing microalga *Heterosigma akashiwo* (Raphidophyceae). *Aquatic Microbial Ecol.* **13**, 135–140 (1997).
104. Juneau, P., Lawrence, J., Suttle, C., Curtis, P. & Harrison, P. J. Effects of viral infection on photosynthetic processes in the bloom-forming alga *Heterosigma akashiwo*. *Aquatic Microbial Ecol.* **31**, 9–17 (2003).
105. Tai, V. et al. Characterization of HaRNAV, a single-stranded RNA virus causing lysis of *Heterosigma akashiwo* (Raphidophyceae) 1. *J. Phycol.* **39**, 343–352 (2003).
106. Mirza, S. F. et al. Isolation and characterization of a virus infecting the freshwater algae *Chrysochromulina parva*. *Virology* **486**, 105–115 (2015).
107. Wagstaff, B. et al. Isolation and characterization of a double stranded DNA megavirus infecting the toxin-producing haptophyte *Prymnesium parvum*. *Viruses* **9**, 40 (2017).
108. Tomaru, Y. et al. New single-stranded DNA virus with a unique genomic structure that infects marine diatom *Chaetoceros setoensis*. *Sci. Rep.* **3**, 3337 (2013).
109. Shirai, Y. et al. Isolation and characterization of a single-stranded RNA virus infecting the marine planktonic diatom *Chaetoceros tenuissimus* meunier. *Appl. Environ. Microbiol.* **74**, 4022–4027 (2008).
110. Kimura, K. & Tomaru, Y. Isolation and characterization of a single-stranded DNA virus infecting the marine diatom *Chaetoceros* sp.

- Strain SS628-11 isolated from western Japan. *PLoS One* **8**, e82013 (2013).
111. Kimura, K. & Tomaru, Y. Discovery of two novel viruses expands the diversity of single-stranded DNA and single-stranded RNA viruses infecting a cosmopolitan marine diatom. *Appl. Environ. Microbiol.* **81**, 1120–1131 (2015).
112. Nagasaki, K. et al. Isolation and characterization of a novel single-stranded RNA virus infecting the bloom-forming diatom *Rhizosolenia Setigera*. *Appl. Environ. Microbiol.* **70**, 704–711 (2004).
113. Tomaru, Y. et al. Isolation and characterization of a single-stranded RNA virus that infects the marine planktonic diatom *Chaetoceros* sp. (SS08-C03). *Phycolog. Res.* **61**, 27–36 (2013).
114. Tomaru, Y., Takao, Y., Suzuki, H., Nagumo, T. & Nagasaki, K. Isolation and characterization of a single-stranded RNA virus infecting the bloom-forming diatom *Chaetoceros socialis*. *Appl. Environ. Microbiol.* **75**, 2375–2381 (2009a).
115. Nagasaki, K. et al. Previously unknown virus infects marine diatom. *Appl. Environ. Microbiol.* **71**, 3528–3535 (2005).
116. Vincent, F., Sheyn, U., Porat, Z., Schatz, D. & Vardi, A. Visualizing active viral infection reveals diverse cell fates in synchronized algal bloom demise. *PNAS* **118**, e2021586118 (2021).
117. Rosenwasser, S. et al. Rewiring host lipid metabolism by large viruses determines the fate of *Emiliania huxleyi*, a bloom-forming alga in the ocean. *Plant Cell* **26**, 2689–2707 (2014).
118. Nissimov, J. I. et al. Dynamics of transparent exopolymer particle production and aggregation during viral infection of the coccolithophore, *Emiliania huxleyi*. *Environ. Microbiol.* **20**, 2880–2897 (2018).
119. Yuxuan, X. et al. Virus-induced autophagy in the marine coccolithophorid *Emiliania Huxleyi*. *Haiyang Xuebao* **45**, 143–154 (2023).
120. Visvesvara, G. S. Giardiasis: an overview. *IMJ. Illinois Med. J.* **164**, 34–39 (1983).
121. Lagunas-Rangel, F. A., Kameyama-Kawabe, L. Y. & Bermúdez-Cruz, R. M. Giardavirus: an update. *Parasitol Res.* **120**, 1943–1948 (2021).
122. Cao, L. et al. *Giardia canis*: ultrastructural analysis of *G. canis* trophozoites transfected with full length *G. canis* virus cDNA transcripts. *Exp. Parasitol.* **123**, 212–217 (2009).
123. Miller, R. L., Wang, A. L. & Wang, C. C. Purification and characterization of the *Giardia Lamblia* double-stranded RNA virus. *Mol. Biochem. Parasitol.* **28**, 189–195 (1988).
124. Marucci, G. et al. Re-discovery of giardavirus: genomic and functional analysis of viruses from *Giardia duodenalis* isolates. *Biomedicines* **9**, 654 (2021).
125. Horiguchi, T. *Heterocapsa circularisquama* sp. nov. (*Peridinales*, *Dinophyceae*): a new marine dinoflagellate causing mass mortality of bivalves in Japan. *Phycokogica Res.* **43**, 129–136 (1995).
126. Tarutani, K., Nagasaki, K., Itakura, S. & Yamaguchi, M. Isolation of a virus infecting the novel shellfish-killing dinoflagellate *Heterocapsa circularisquama*. *Aquatic Microbial Ecol.* **23**, 103–111 (2001).
127. Takano, Y., Tomaru, Y. & Nagasaki, K. Visualization of a dinoflagellate-infecting virus HcDNAV and its infection process. *Viruses* **10**, 554 (2018).
128. Coy, S. R. et al. Visualization of RNA virus infection in a marine protist with a universal biomarker. *Sci. Rep.* **13**, 5813 (2023).
129. Tomaru, Y., Mizumoto, H. & Nagasaki, K. Virus resistance in the toxic bloom-forming dinoflagellate *Heterocapsa circularisquama* to single-stranded RNA virus infection. *Environ. Microbiol.* **11**, 2915–2923 (2009b).
130. Tomaru, Y. et al. Isolation and characterization of two distinct types of HcRNAV, a single-stranded RNA virus infecting the bivalve-killing microalga *Heterocapsa circularisquama*. *Aquatic Microbial Ecol.* **34**, 207–218 (2004).
131. Mizumoto, H., Tomaru, Y., Takao, Y., Shirai, Y. & Nagasaki, K. Diverse responses of the bivalve-killing dinoflagellate *Heterocapsa circularisquama* to infection by a single-stranded RNA virus. *Appl. Environ. Microbiol.* **74**, 3105–3111 (2008).
132. Weynberg, K. D. et al. Prevalent and persistent viral infection in cultures of the coral algal endosymbiont *Symbiodinium*. *Coral Reefs* **36**, 773–784 (2017).
133. Nakano, S., Suzuki, S. I. & Onji, M. Virus-like particles suppress growth of the red-tide-forming marine Dinoflagellate *Gymnodinium mikimotoi*. *Mar. Biotechnol.* **5**, 435–442 (2003).
134. Takao, Y., Nagasaki, K. & Honda, D. Squashed ball-like dsDNA virus infecting a marine fungoid protist *Sicyodochytrium minutum* (*Thraustochytriaceae*, *Labyrinthulomycetes*). *Aquatic Microbial Ecol.* **49**, 101–108 (2007).
135. Westermann, A. J., Gorski, S. A. & Vogel, J. Dual RNA-seq of pathogen and host. *Nat. Rev. Microbiol.* **10**, 618–630 (2012).
136. Lum, K. K. & Cristea, I. M. Proteomic approaches to uncovering virus-host protein interactions during the progression of viral infection. *Expert Rev. Proteom.* **13**, 325–340 (2016).
137. Rodrigues, R. A. L. et al. Analysis of a Marseillevirus transcriptome reveals temporal gene expression profile and host transcriptional shift. *Front. Microbiol.* **11**, 651 (2020). **Giant virus infection rapidly triggers the downregulation of genes related to host transcriptional machinery.**
138. Zhang, R., Endo, H., Takemura, M. & Ogata, H. RNA sequencing of medusavirus suggests remodeling of the host nuclear environment at an early infection stage. *Microbiol. Spectr.* **9**, e0006421 (2021).
139. Rolland, C., La Scola, B. & Levasseur, A. How tupanvirus degrades the ribosomal RNA of its amoebal host? The ribonuclease T2 track. *Front. Microbiol.* **11**, 1691 (2020). **Tupanvirus soda lake virus infection triggers rRNA shutdown potentially through RNase T2 enzyme gene modulation.**
140. Jones, B. M., Edwards, R. J., Skipp, P. J., O'Connor, C. D. & Iglesias-Rodriguez, M. D. Shotgun proteomic analysis of *Emiliania huxleyi*, a marine phytoplankton species of major biogeochemical importance. *Mar. Biotechnol.* **13**, 496–504 (2011).
141. Schroeder, D. C., Oke, J., Malin, G. & Wilson, W. H. Coccolithovirus (Phycodnaviridae): characterisation of a new large dsDNA algal virus that infects *Emiliania huxleyi*. *Arch. Virol.* **147**, 1685–1698 (2002).
142. Liu, J. et al. Characterization of the sphingolipid profiling of *Emiliania huxleyi* against virus infection. *J. Oceanol. Limnol.* **41**, 1547–1557 (2023).
143. Pagarete, A., Allen, M. J., Wilson, W. H., Kimmance, S. A. & De Vargas, C. Host-virus shift of the sphingolipid pathway along an *Emiliania huxleyi* bloom: survival of the fittest. *Environ. Microbiol.* **11**, 2840–2848 (2009).
144. Schleyer, G. et al. In plaque-mass spectrometry imaging of a bloom-forming alga during viral infection reveals a metabolic shift towards odd-chain fatty acid lipids. *Nat. Microbiol.* **4**, 527–538 (2019).
145. Evans, C., Pond, D. W. & Wilson, W. H. Changes in *Emiliania huxleyi* fatty acid profiles during infection with *E. huxleyi* virus 86: Physiological and ecological implications. *Aquatic Microb. Ecol.* **55**, 219–228 (2009).
146. Hunter, J. E., Frada, M. J., Fredricks, H. F., Vardi, A. & Van Mooy, B. A. S. Targeted and untargeted lipidomics of *Emiliania huxleyi* viral infection and life cycle phases highlights molecular biomarkers of infection, susceptibility, and ploidy. *Front. Mar. Sci.* **2**, 81 (2015).
147. Malitsky, S. et al. Viral infection of the marine alga *Emiliania huxleyi* triggers lipidome remodeling and induces the production of highly saturated triacylglycerol. *New Phytol.* **210**, 88–96 (2016).
148. Kegel, J. U. et al. Transcriptional host-virus interaction of *Emiliania huxleyi* (*Haptophyceae*) and EhV-86 deduced from combined analysis of expressed sequence tags and microarrays. *Eur. J. Phycol.* **45**, 1–12 (2010).

149. Sheyn, U. et al. Expression profiling of host and virus during a coccolithophore bloom provides insights into the role of viral infection in promoting carbon export. *ISME J.* **12**, 704–713 (2018).
150. Pagarete, A. et al. Unveiling the transcriptional features associated with coccolithovirus infection of natural *Emiliania huxleyi* blooms. *FEMS Microb. Ecol.* **78**, 555–564 (2011).
151. Hevroni, G., Vincent, F., Sheyn, U., Vardi, A., & Ku, C. Daily turnover of active giant virus infection during algal blooms revealed by single-cell transcriptomics. *Sci. Adv.* **9**, 41 (2023).
152. Zhang, E. et al. MicroRNA-mediated regulation of lipid metabolism in virus-infected *Emiliania huxleyi*. *ISME J.* **16**, 2457–2466 (2022).
153. Ku, C. et al. A single-cell view on alga-virus interactions reveals sequential transcriptional programs and infection states. *Sci. Adv.* **6**, 1–10 (2020).
154. Feldmesser, E., Ben-Dor, S. & Vardi, A. An *Emiliania huxleyi* pan-transcriptome reveals basal strain specificity in gene expression patterns. *Sci. Rep.* **11**, 20795 (2021).
155. Xue, T. et al. Transcriptome analysis of marine microalga *Emiliania huxleyi* in response to virus infection[J]. *Haiyang Xuebao* **41**, 103–112 (2019).
156. Sheyn, U., Rosenwasser, S., Ben-Dor, S., Porat, Z. & Vardi, A. Modulation of host ROS metabolism is essential for viral infection of a bloom-forming coccolithophore in the ocean. *ISME J.* **10**, 1742–1754 (2016).
157. Schatz, D. et al. Hijacking of an autophagy-like process is critical for the life cycle of aDNA virus infecting oceanic algal blooms. *New Phytol.* **204**, 854–863 (2014).
158. Liu, J., Bratbak, G., Zheng, T. & Thyrhaug, R. Effects of virus infection on expression of cell cycle regulatory proteins in the unicellular marine algae *Emiliania huxleyi*. *Acta Oceanol. Sin.* **30**, 89–95 (2011).
159. Jingwen Liu. Virus infection disturbs cyclin expression, leading to cell cycle arrest in the unicellular marine algae *Emiliania huxleyi* and *Chrysochromulina ericina*. *Afr. J. Microbiol. Res.* **5**, 1801–1807 (2011).
160. Moniruzzaman, M., Gann, E. R. & Wilhelm, S. W. Infection by a giant virus (AaV) induces widespread physiological reprogramming in *Aureococcus anophagefferens* CCMP1984-A harmful bloom algae. *Front. Microbiol.* **9**, 752 (2018). **Giant virus infection triggers fast host transcriptional alterations, such as the suppression of transcripts related to photosynthesis, and activation of protein synthesis transcripts.**
161. Bale, N. J. et al. Fatty acid dynamics during viral infection of *Phaeocystis globosa*. *Aquatic Microb. Ecol.* **74**, 85–94 (2015).
162. Chen, S., Gao, K. & Beardall, J. Viral attack exacerbates the susceptibility of a bloom-forming alga to ocean acidification. *Glob. Chang. Biol.* **21**, 629–636 (2015).
163. Ray, J. L. et al. Virus infection of *Haptolina ericina* and *Phaeocystis pouchetii* implicates evolutionary conservation of programmed cell death induction in marine haptophyte-virus interactions. *J. Plankton Res.* **36**, 943–955 (2014).
164. Sheik, A. R. et al. Viral infection of *Phaeocystis globosa* impedes release of chitinous star-like structures: quantification using single cell approaches. *Environ. Microbiol.* **15**, 1441–1451 (2013).
165. Lawrence, S. A., Floge, S. A., Davy, J. E., Davy, S. K. & Wilson, W. H. Exploratory analysis of *Symbiodinium* transcriptomes reveals potential latent infection by large dsDNA viruses. *Environ. Microbiol.* **19**, 3909–3919 (2017).
166. Wang, A. L. & Wang, C. C. A linear double-stranded RNA in *Trichomonas vaginalis*. *J. Biol. Chem.* **260**, 3697–3702 (1985).
167. Graves, K. J., Ghosh, A. P., Kissinger, P. J. & Muzny, C. A. *Trichomonas vaginalis* virus: a review of the literature. *Int. J. STD AIDS* **30**, 496–504 (2019).
168. Wang, A., Wang, C. C. & Alderete, J. F. “*Trichomonas vaginalis* phenotypic variation occurs only among trichomonads infected with the double-stranded RNA virus.”. *J. Exp. Med.* **166**, 1 (1987).
169. Khoshnan, A. & Alderete, J. F. “*Trichomonas vaginalis* with a double-stranded RNA virus has upregulated levels of phenotypically variable immunogen mRNA.”. *J. Virol.* **68**, 6 (1994).
170. Provenzano, D. & Alderete, J. F. Analysis of human immunoglobulin-degrading cysteine proteinases of *Trichomonas Vaginalis*. *Infect. Immun.* **63**, 3388–3395 (1995).
171. Provenzano, D., Khoshnan, A. & Alderete, J. F. Involvement of dsRNA virus in the protein composition and growth kinetics of host *Trichomonas vaginalis*. *Arch. Virol.* **142**, 939–952 (1997).
172. Goodman, R. P. et al. Clinical isolates of *Trichomonas vaginalis* concurrently infected by strains of up to four trichomonasvirus species (Family Totiviridae). *J. Virol.* **85**, 4258–4270 (2011).
173. Bessarab, I. N., Nakajima, R., Liu, H. W. & Tai, J. H. Identification and characterization of a type III *Trichomonas vaginalis* virus in the protozoan pathogen *Trichomonas vaginalis*. *Arch. Virol.* **156**, 285–294 (2011).
174. He, D. et al. Differential protein expressions in virus-infected and uninfected *Trichomonas vaginalis*. *Korean J. Parasitol.* **55**, 121–128 (2017).
175. Rada, P. et al. Double-stranded RNA viruses are released from *Trichomonas vaginalis* inside small extracellular vesicles and modulate the exosomal cargo. *Front. Microbiol.* **13**, 893692 (2022).
176. Tarr, P. I. et al. LR1: a candidate RNA virus of *Leishmania* (Leishmania *Braziliensis* Guyanensis/Single-Stranded RNA/Viral Particle/ Electron Microscopy). *Proc. Natl. Acad. Sci. USA* **85**, 9572–9575 (1988).
177. Gupta, V. & Deep, A. An insight into the Leishmania RNA virus. *Indian J. Med. Microbiol.* **25**, 7–9 (2007).
178. Cantanhêde, L. M. et al. The maze pathway of coevolution: a critical review over the *Leishmania* and its endosymbiotic history. *Genes* **12**, 657 (2021).
179. Ives, A. et al. Leishmania RNA virus controls the severity of mucocutaneous leishmaniasis. *Science* **331**, 775–778 (2011).
180. Saura, A. et al. Elimination of LRVs elicits different responses in *Leishmania* spp. *mSphere* **7**, e0033522 (2022).
181. Kamoun, S. et al. The Top 10 oomycete pathogens in molecular plant pathology. *Mol. Plant Pathol.* **16**, 413–434 (2015).
182. Botella, L. & Jung, T. Multiple viral infections detected in phytophthora condilina by total and small rna sequencing. *Viruses* **13**, 620 (2021).
183. Hannat, S. et al. Diverse trajectories drive the expression of a giant virus in the oomycete plant pathogen *Phytophthora Parasitica*. *Front. Microbiol.* **12**, 662762 (2021).
184. Uchida, K. et al. Two novel endornaviruses co-infecting a phytophthora pathogen of *Asparagus officinalis* modulate the developmental stages and fungicide sensitivities of the host oomycete. *Front. Microbiol.* **12**, 633502 (2021).
185. Mascia, T. et al. Infection of *Colletotrichum acutatum* and *Phytophthora infestans* by taxonomically different plant viruses. *Eur. J. Plant Pathol.* **153**, 1001–1017 (2019).
186. Cai, G., Fry, W. E. & Hillman, B. I. PiRV-2 stimulates sporulation in *Phytophthora infestans*. *Virus Res.* **271**, 197674 (2019).
187. Poimala, A. et al. Bunyaviruses affect growth, sporulation, and Elicitin production in *Phytophthora cactorum*. *Viruses* **14**, 2596 (2022).
188. Kamoun, S. A catalogue of the effector secretome of plant pathogenic oomycetes. *Ann. Rev. Phytopathol.* **44**, 41–60 (2006).
189. Kharel, A., Islam, M. T., Rookes, J. & Cahill, D. How to unravel the key functions of cryptic oomycete elicitor proteins and their role in plant disease. *Plants* **10**, 1201 (2021).
190. Slaveykova, V., Sonntag, B. & Gutiérrez, J. C. Stress and protists: no life without stress. *Eur. J. Protistol.* **55**, 39–49 (2016).
191. Khan, N. A. Life Cycle and Ecological Significance in *Acanthamoeba: biology and pathogenesis*. 70–73 (Caister Academic Press, 2009).

192. Vardi, A. et al. Programmed cell death of the dinoflagellate *Peridinium gatunense* is mediated by CO₂ limitation and oxidative stress. *Curr. Biol.* **9**, 1061–1064 (1999).
193. Frischkorn, K. R., Harke, M. J., Gobler, C. J. & Dyhrman, S. T. De novo assembly of *Aureococcus anophagefferens* transcriptomes reveals diverse responses to the low nutrient and low light conditions present during blooms. *Front. Microbiol.* **5**, 375 (2014).
194. Pelusi, A. et al. Virus-induced spore formation as a defense mechanism in marine diatoms. *New Phytol.* **229**, 2251–2259 (2021).
195. Frada, M. J. et al. Morphological switch to a resistant subpopulation in response to viral infection in the bloom-forming coccolithophore *Emiliania huxleyi*. *PLoS Pathog.* **13**, e1006775 (2017). **Show that the haploid phase of *E. huxleyi* is resistant to *E. huxleyi* viruses, unlike the diploid cell.**
196. De Vargas, C., Aubry, M. P., Probert, I. A. N., & Young, J. Origin and evolution of coccolithophores: from coastal hunters to oceanic farmers in *Evolution of Primary Producers in the Sea*. 251–285 (Academic Press, 2007).
197. Frada, M., Probert, I., Allen, M. J., Wilson, W. H. & de Vargas, C. The ‘Cheshire Cat’ escape strategy of the coccolithophore *Emiliania huxleyi* in response to viral infection. *PNAS* **105**, 15944–15949 (2008).
198. Zajac, A. J., Amphlett, E. M., Rowlands, D. J. & Sangar, D. V. Parameters influencing the attachment of Hepatitis A virus to a variety of continuous cell lines. *J. General Virol.* **72**, 1667–1675 (1991).
199. Stapleton, J. T., Frederick, J. & Meyer, B. Hepatitis A. Virus attachment to cultured cell lines. *J. Infect. Dis.* **164**, 1098–1103 (1991).
200. Cheshenko, N., Liu, W., Satlin, L. M. & Herold, B. C. Multiple receptor interactions trigger release of membrane and intracellular calcium stores critical for herpes simplex virus entry. *Mol. Biol. Cell* **18**, 3119–3130 (2007).
201. Bishop, N. E. & Anderson, D. A. Early interactions of Hepatitis A virus with cultured cells: viral elution and the effect of PH and calcium ions. *Arch Virol.* **142**, 2161–2178 (1997).
202. Wilson, W. H. et al. Isolation of viruses responsible for the demise of an *Emiliania huxleyi* bloom in the English. *Channel. J. Mar. Biol.* **82**, 369–377 (2002).
203. Bratbak, G., Egge, J. K. & Heldal, M. Viral mortality of the marine alga *Emiliania huxleyi* (Haptophyceae) and termination of algal blooms. *Mar. Ecol. Prog. Ser.* **93**, 39–48 (1993).
204. Evans, C., Malin, G., Mills, G. P. & Wilson, W. H. Viral infection of *Emiliania huxleyi* (Prymnesiophyceae) leads to elevated production of reactive oxygen species. *J. Phycol.* **42**, 1040–1047 (2006).
205. Mordecai, G. J., Verret, F., Highfield, A. & Schroeder, D. C. Schrödinger’s cheshire cat: are haploid *Emiliania huxleyi* cells resistant to viral infection or not? *Viruses* **9**, 51 (2017).
206. Silva, L. K., dos, S., Boratto, P. V. M., La Scola, B., Bonjardim, C. A. & Abrahão, J. S. *Acanthamoeba* and mimivirus interactions: the role of amoebal encystment and the expansion of the ‘Cheshire Cat’ theory. *Curr. Opin. Microbiol.* **31**, 9–15 (2016). **Highlights interactions between amoebae and giant viruses concerning encystment of the protist cells.**
207. Neff, R. J. S. A. R. W. F. B. & Wilborn, M. Induction of synchronous encystment (differentiation) in *Acanthamoeba* sp. *Methods Cell Biol.* **1**, 55–83 (1964).
208. Cordingley, J. S., Wills, R. A. & Vilemez, C. L. Osmolarity is an independent trigger of *Acanthamoeba Castellanii* differentiation. *J. Cell Biochem.* **61**, 167–171 (1996).
209. Aksozek, A., McClellan, K. & Howard, K. Resistance of *Acanthamoeba Castellanii* cysts to physical, chemical, and radiological conditions. *J. Parasitol.* **88**, 621–623 (2002).
210. Moon, E. K., Chung, D., Il, Hong, Y. C. & Kong, H. H. Characterization of a serine proteinase mediating encystation of *Acanthamoeba*. *Eukaryot Cell* **7**, 1513–1517 (2008).
211. La Scola, B. et al. The virophage as a unique parasite of the giant mimivirus. *Nature* **455**, 100–104 (2008). **Describes a small virus, a virophage, that is associated with a new strain of mimivirus.**
212. Fischer, M. G. & Hackl, T. Host genome integration and giant virus-induced reactivation of the virophage mavirus. *Nature* **540**, 288–291 (2016).
213. Fischer, M. G. & Suttle, C. A. A virophage at the origin of large DNA transposons. *Science* **332**, 231–234 (2011).
214. Gaia, M. et al. Zamilon, a novel virophage with Mimiviridae host specificity. *PLoS One* **9**, e94923 (2014).
215. Arthofer, P., Delafont, V., Willemssen, A., Panholzl, F. & Horn, M. Defensive symbiosis against giant viruses in amoebae. *PNAS* **119**, e2205856119 (2022).
216. George, E. E. et al. A single cryptomonad cell harbors a complex community of organelles, bacteria, a phage, and selfish elements. *Curr. Biol.* **33**, 1982–1996.e4 (2023).
217. Javaux, E. J., Knoll, A. H. & Walter, M. R. Morphological and ecological complexity in early eukaryotic ecosystems. *Nature* **412**, 66–69 (2001).
218. Guglielmini, J., Woo, A. C., Krupovic, M., Forterre, P. & Gaia, M. Diversification of giant and large eukaryotic dsDNA viruses predated the origin of modern eukaryotes. *Proc Natl Acad Sci USA* **116**, 19585–19592 (2019).
219. Arsenieff, L. et al. First viruses infecting the marine diatom *Guinardia delicatula*. *Front. Microbiol.* **10**, 3235 (2019).
220. Bidle, K. D., Haramaty, L., Barcelos Ramos, J. & Falkowski, P. Viral activation and recruitment of metacaspases in the unicellular coccolithophore, *Emiliania huxleyi*. *PNAS* **104**, 6049–6054 (2007).
221. Irwin, N. A. T., Pittis, A. A., Richards, T. A. & Keeling, P. J. Systematic evaluation of horizontal gene transfer between eukaryotes and viruses. *Nat. Microbiol.* **7**, 327–336 (2022). **Highlight the large number of horizontal gene transfer events that nucleocytoviruses have had mainly with protist hosts throughout natural history.**
222. Moliner, C., Fournier, P. E. & Raoult, D. Genome analysis of microorganisms living in amoebae reveals a melting pot of evolution. *FEMS Microbiol. Rev.* **34**, 281–294 (2010).
223. Keeling, P. J. & Eglit, Y. Openly available illustrations as tools to describe eukaryotic microbial diversity. *PLoS Biol.* **21**, e3002395 (2023).
224. Mihara, T. et al. Linking virus genomes with host taxonomy. *Viruses* **8**, 66 (2016).
225. Tomaru, Y., Shirai, Y., Suzuki, H., Nagumo, T. & Nagasaki, K. Isolation and characterization of a new single-stranded DNA virus infecting the cosmopolitan marine diatom *Chaetoceros debilis*. *Aquatic Microbial Ecol.* **50**, 103–112 (2008).

Acknowledgements

We thank our colleagues from Laboratório de Vírus—UFMG for their technical support. We acknowledge financial support from Rede Vírus—Ministério da Ciência, Tecnologia e Inovações (MCTI), Câmara Pox—405249/2022-5. We thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), grant number 88882.348380/2010-1, Fundação de Amparo à Pesquisa do estado de Minas Gerais (FAPEMIG), Programas Institutos Nacionais de Ciência e Tecnologia (INCT), grant number 406441/2022-7, chamada 58/2022, and Pró-Reitorias de Pesquisa e Pós-Graduação de UFMG, and the Centre for New Antibacterial Strategies (CANS) of the Arctic University of Norway (project ID #2520855). R.A.L.R. and J.S.A. are CNPq researchers.

Author contributions

V.F.Q., J.M.T. and B.B.B. performed the literature review and prepared the figures and tables. R.A.L.R., G.M.F.A. and J.S.A. designed the text structure and revised the manuscript. All authors contributed to the writing.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s42003-024-06001-2>.

Correspondence and requests for materials should be addressed to Gabriel Magno de Freitas Almeida.

Peer review information *Communications Biology* thanks Chuan Ku, Guifré Torruella and the other, anonymous, reviewer(s) for their contribution to the peer review of this work. Primary Handling Editor: Manuel Breuer. A peer review file is available.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024