


# Galactic planetary science (Introduction)

Giovanna Tinetti

*University College London & Royal Society*


# The Exoplanet Revolution

9 to 2000 in 20 years!



# The Exoplanet Revolution

9 to 2000 in 20 years!



# Planets before 1995...

Small rocky planets close to the Sun  
Gas-giant planets more distant from the star



# The first extrasolar planet

Sun

Mercury



Venus




Earth



Solar system:  
small rocky planets close to the Sun  
gas-giant planets more distant from the star → →

51 Peg



51 Peg b

51 Pegasus: a gas-giant very close to its parent star (hot-Jupiter)


# Hot Jupiters – migration



**Hot-Jupiters are Gas-Giant planets, orbiting VERY close to their parent star. They are probably tidally locked, i.e. one face is always illuminated and the other is in perpetual darkness.**

**They easily reach Temperatures 1000-2000 K**

# ~2000 Exoplanets!



# Exoplanet today

## The lightest

Mercury's mass (KOI-1843 b)

## The heaviest

30 Jupiter masses (HD 284149 b)

## The shortest year

4.5 hours (KOI-1843 b)

## The longest year


730000 days (Oph 11 b)

## The closest to the Earth

4.4 light years (Alpha Cent Bb)

## The farthest to the Earth


22000 light years (OGLE-390 b)







# Sub- and Super-Earths



# Exoplanets are common...

On average, every star host a planet




Thousands of planetary candidates discovered by Kepler!


# Locations of Kepler Planet Candidates

*As of January 7, 2013*

- Earth-size
- Super-Earth size  
1.25 - 2.0 Earth-size
- Neptune-size  
2.0 - 6.0 Earth-size
- Giant-planet size  
6.0 - 22 Earth-size



As of January 7, 2013



# Pulsar timing

---

In 1992 Aleksander Wolszczan and Dale Frail used this method to discover planets around the pulsar PSR 1257+12.

Their discovery was quickly confirmed, making it the first confirmation of planets outside our Solar System.

*A pulsar is a neutron star: the small, ultradense remnant of a star that has exploded as a supernova. Pulsars emit radio waves extremely regularly as they rotate. Slight anomalies in the timing of its observed radio pulses can be used to track the pulsar's motion. Like an ordinary star, a pulsar will move in its own small orbit if it has a planet.*


# Radial velocity & astrometry

---



# Astrometry

Motion of the Solar System barycenter relative to the Sun



# Detections of exoplanets using astrometry


---

- So far no planets detected
- Gaia space mission will measure stellar parallaxes with unprecedented precision
- Gaia space mission will feature accuracy of  $\sim 30$  micro arc-sec
- Gaia is predicted to discover 10,000 planets using astrometry in our local surrounding alone





# Radial velocity




The star motion is detectable with the Doppler Effect.

The light coming from a star moving toward the Earth will be Doppler shifted to bluer (shorter) wavelengths, while a star receding from the Earth will emit light shifted to redder (longer) wavelengths.

The effect is very small. For instance Jupiter induces a 12 m/s velocity change on the Sun for an external observer whereas the Saturn effect is only 2.7 m/s.

# Doppler spectroscopy

A tiny portion of the spectrum in the near-IR



# Doppler spectroscopy

---


Look at the line which appears in the Sun at about 882.4 nm.  
In the spectrum of Arcturus, it appears at about 882.55 nm.

$$\begin{aligned}\text{shift (Arcturus - Sun)} &= 882.55 \text{ nm} - 882.4 \text{ nm} \\ &= 0.15 \text{ nm}\end{aligned}$$

$$\begin{aligned}\text{radial velocity} &= \frac{0.15 \text{ nm}}{882.4 \text{ nm}} * (300,000 \text{ km/s}) \\ &= 50 \text{ km/s}\end{aligned}$$

So Arcturus was moving AWAY from us at about 50 km/s  
when the spectrum was taken.

# Measuring the radial velocity parameters



# Summary of equations

---

Kepler's third law:

$$a^3 = \frac{GM_*}{4\pi^2} P^2$$

Conservation of momentum:

$$M_p v_p = M_* v_*$$

Radial velocity curve amplitude:

$$K_* \simeq v_* \sin i = \frac{M_p \sin i}{M_*} v_p$$

Planetary speed:

$$v_p = \frac{2\pi a}{P}$$

# Which parameters can we observe directly?

---

Kepler's third law:

$$a^3 = \frac{GM_*}{4\pi^2} P^2$$

Conservation of momentum:

$$M_p v_p = M_* v_*$$

Radial velocity curve amplitude:

$$K_* \simeq v_* \sin i = \frac{M_p \sin i}{M_*} v_p$$

Planetary speed:

$$v_p = \frac{2\pi a}{P}$$

# Which parameters need to be observed otherwise?

Kepler's third law:

$$\overset{\text{derived}}{a^3} = \frac{GM_*}{4\pi^2} P^2$$

Conservation of momentum:

$$\overset{\text{derived}}{M_p v_p} = M_* v_*$$

Radial velocity curve amplitude:

$$\overset{\text{derived}}{K_*} \simeq v_* \sin i = \frac{\overset{\text{derived}}{M_p \sin i}}{M_*} v_p$$

Planetary speed:

$$\overset{\text{derived}}{v_p} = \frac{2\pi a}{P}$$

# Examples of RV of measurements


Planet Mass	Distance AU	Radial velocity
Jupiter	1	28.4 m/s
Jupiter	5	12.7 m/s
Neptune	0.1	4.8 m/s
Neptune	1	1.5 m/s
Super-Earth (5 M <sub>⊕</sub> )	0.1	1.4 m/s
Super-Earth (5 M <sub>⊕</sub> )	1	0.45 m/s
Earth	1	9 cm/s

For MK-type stars with planets in the habitable zone

Stellar Mass (M <sub>☉</sub> )	Planet Mass (M <sub>⊕</sub> )	Lum. (L <sub>☉</sub> )	Type	RHAB. (AU)	RV (cm/s)	Period (days)
0.10	1.0	8e-4	M8	0.028	168	6
0.21	1.0	7.9e-3	M5	0.089	65	21
0.47	1.0	6.3e-2	M0	0.25	26	67
0.65	1.0	1.6e-1	K5	0.40	18	115
0.78	2.0	4.0e-1	K0	0.63	25	209




# Effect of orientation




# The effect of ellipticity

- A different ellipticity will also change the variation of the line-of-sight velocity with time
- For simplicity, let us assume that the plane of the orbit is aligned parallel to the line of sight



# Upsilon Andromeda<sub>e</sub>

A complex system




This system shows a much more complex velocity structure revealing the presence of a multiple planetary system.

Host star F8V ( $1.3M_{\odot}$ )

Distance: 44 lyr


# Radial velocity instruments

- HARPS-South and HARPS-North (High Accuracy Radial velocity Planet Searcher) are the most accurate spectrographs ever built.
- It can detect the wobble of a star by with a precision of 30cm per second. As a guide, 51 Peg b has a radial velocity amplitude  $\sim 16$  times larger.
- Multiple Super-Earths and complex multi-planet systems were discovered
- Current sites: 3.6m telescope on La Silla (Chile), 3.58m TNG on La Palma (Spain)




# Transit of an exoplanet

Even a Jupiter size planet will only obscure ~ 1% of the stellar light.





The transit depth is proportional to the surface ratios of the planet and the star:

$$\Delta F = \left( \frac{R_{planet}}{R_{star}} \right)^2$$





# Transit of an exoplanet




$$R_{\text{planet}}^2 / R_{\text{star}}^2 \sim 1\%$$

# You can do it from London, with 35 cm telescope...



*HUBBLE SPACE TELESCOPE TIME-SERIES PHOTOMETRY OF THE TRANSITING PLANET  
OF HD 209458*<sup>1</sup>


TIMOTHY M. BROWN,<sup>2</sup> DAVID CHARBONNEAU,<sup>2,3</sup> RONALD L. GILLILAND,<sup>4</sup> ROBERT W. NOYES,<sup>3</sup> AND ADAM BURROWS<sup>5</sup>  
*Received 2000 November 21; accepted 2001 January 18*



**FIG. 3.**—Phased light curve for all four transits, assuming a planetary orbital period of 3.52474 days. The time series for each transit has been scaled to have the same average intensity over the second and fifth (out-of-transit) orbits.



# Solving the star-planet system



# Solving the star-planet system

From Kepler's third law and radial velocity we have:


$$P^2 = \frac{4\pi^2 a^3}{G(M_* + M_p)}$$

From the transit depth measurement we have the planet/star ratio:

$$\frac{R_p}{R_*} = \sqrt{\Delta F}$$

We can now calculate the orbital inclination:

$$i = \cos^{-1} \left( b \frac{R_*}{a} \right)$$



# Solving the star-planet system

We can now calculate the orbital inclination:


$$i = \cos^{-1} \left( b \frac{R_*}{a} \right)$$

What is b?? - it's called the impact parameter:

$$b = \left[ \frac{\left(1 - \sqrt{\Delta F}\right)^2 - (t_F/t_T)^2 \left(1 + \sqrt{\Delta F}\right)^2}{1 - (t_F/t_T)^2} \right]^{1/2}$$



We can even calculate the stellar density:

$$\rho_* = \frac{32}{G\pi} P \frac{\Delta F^{3/4}}{(t_T^2 - t_F^2)^{3/2}}$$




# Transit hunters

Earth-size planet transiting Sun-type star ~ 0.01 %



Kepler (NASA)  
2009 - present



COROT (CNES/ESA)  
2006-2013



# Transits from the ground



MEarth




Hat-NET



Super-WASP

# Future transit missions/surveys

- ESA-Cheops
- NASA-TESS
- ESA-PLATO
- NGTS



# From Jupiter's moons...




Galileo (the mission), 1989



Galileo « Sidereus Nuncius » 1610


# Search for Exomoons

---







# Transit duration variation (TDV)




# Transit duration variation (TDV)




# Transit duration variation (TDV)




# Transit duration variation (TDV)



# Transit duration variation (TDV)




# Transit duration variation (TDV)




# Gravitational Lensing

- Einstein's General Theory of Relativity: light rays can be bent by a sufficiently strong gravitational field.
- Take 1 background (A) and 1 foreground star (B)
- Light rays from A will be bent by the gravitational field of B → more light from the distant star reaches the Earth than would otherwise be the case



# Gravitational Lensing

- This amount of bending is very small and is related to the size of a black hole with the mass of the lensing star (B): Schwarzschild radius  $R_{\text{SCH}} = 2GM/c^2$




$$\alpha = 4GM/c^2 b = 2R_{\text{SCH}}/b$$



For the Sun:  
 $R_{\text{SCH}} = 2.95\text{km}$   
 $\alpha = 2R_{\text{SCH}}/R_{\odot} = 1.74''$




# Gravitational Lensing




# Gravitational Lensing

---



# Microlensing and exoplanets


- If the lensing star has a planet, it could be seen as a small glitch in the light curve



# Direct detection

---






The image shows a star at the center, marked with a white cross. Three planets are visible as bright spots, each circled in green. The planets are labeled 'b' (top left), 'c' (top right), and 'd' (bottom right). The background is a dark, noisy field of stars.

**Direct detection:  
Gemini + Keck  
Planetary system HR 8799**




# Direct detection: VLT-NACO $\beta$ -Pictoris b (~8 AU)



# New generation of DI instruments:


GPI, SPHERE, SUBARU



# Formalhaut b (HST)

Fomalhaut System

Hubble Space Telescope • STIS




NASA and ESA

STScI-PRC13-01a






# Formalhaut b (Alma)

---



## Formalhaut b

- Hot Jupiter first discovered in 2008
- Orbital parameters:  $a = 115\text{au}$ ,  $e = 0.11$ ,  $i = 66$  degrees
- Orbital period of 872 years
- Some debates on its existence in the literature but is now thought to be confirmed.



# Where are these planets?

