

# LaeA, a Regulator of Secondary Metabolism in *Aspergillus* spp.

Jin Woo Bok and Nancy P. Keller\*

Department of Plant Pathology, University of Wisconsin—Madison, Madison, Wisconsin 53706

Received 5 November 2003/Accepted 16 December 2003

**Secondary metabolites, or biochemical indicators of fungal development, are of intense interest to human-kind due to their pharmaceutical and/or toxic properties. We present here a novel *Aspergillus* nuclear protein, LaeA, as a global regulator of secondary metabolism in this genus. Deletion of *laeA* ( $\Delta laeA$ ) blocks the expression of metabolic gene clusters, including the sterigmatocystin (carcinogen), penicillin (antibiotic), and lovastatin (antihypercholesterolemic agent) gene clusters. Conversely, overexpression of *laeA* triggers increased penicillin and lovastatin gene transcription and subsequent product formation. *laeA* expression is negatively regulated by AflR, a sterigmatocystin Zn<sub>2</sub>Cys<sub>6</sub> transcription factor, in a unique feedback loop, as well as by two signal transduction elements, protein kinase A and RasA. Although these last two proteins also negatively regulate sporulation,  $\Delta laeA$  strains show little difference in spore production compared to the wild type, indicating that the primary role of LaeA is to regulate metabolic gene clusters.**

A complex and fascinating aspect of fungal development is the production of secondary metabolites. These compounds, frequently associated with sporulation processes, are considered part of the chemical arsenal required for niche specialization and have garnered intense interest by virtue of their biotechnological and pharmaceutical applications (9, 11). Many of them display a broad range of useful antibiotic, antiviral, antitumor, antihypercholesterolemic, and immunosuppressant activities as well as less desirable phyto- and mycotoxic activities. Hawksworth's studies of fungal biodiversity led to the conclusion that nearly 1.5 million fungal species exist on Earth, with only 5% identified thus far (16). Thus, the potential for fungal secondary metabolite discovery is vast. Furthermore, the discovery of global regulators for fungal secondary metabolite production is critical, as it would allow for universal manipulation of secondary metabolite production.

A large number of known fungal secondary metabolites have been ascribed to the Ascomycete genus *Aspergillus*. Studies of *Aspergillus nidulans* have demonstrated the power of using a model system to elucidate the biochemistry and molecular genetics of fungal secondary metabolism, principally penicillin (PN, an antibiotic) and sterigmatocystin (ST, a carcinogen biochemically related to the agricultural contaminant aflatoxin) biosynthesis (6, 17). These studies have established several characteristics of fungal secondary metabolism, including the clustering of biosynthetic and regulatory genes and a genetic connection linking secondary metabolite biosynthesis with sporulation through a shared signal transduction pathway (9).

The discovery of the G protein-cyclic AMP (cAMP)-protein kinase A regulation of ST, aflatoxin, and other fungal secondary metabolites (4, 17, 29, 34) has been helpful for establishing a model of global regulation of secondary metabolism. However, all of the signal transduction mutants described in the literature have pleiotropic effects on fungi, the most notable

effect being the gross impact on spore production and vegetative hyphal growth (1, 9, 18, 29). Similarly, mutations in other major regulators of ST biosynthesis, such as RcoA, a WD protein (19), and SpdA, or spermidine synthase (20), also have gross effects on fungal morphology. Thus, currently available *Aspergillus* mutants are so impaired in fungal development that further elucidation of genes that are specific for the regulation of secondary metabolism is difficult.

In a previous mutagenesis hunt, 23 *A. nidulans* mutants that displayed a phenotype of loss of ST production but had normal sporulation were isolated (8). Here we describe the identification of a gene called *laeA* that complements one of these mutants. *laeA* encodes a nuclear protein that is required for the expression of secondary metabolite genes. We propose that LaeA is a regulator of secondary metabolism in *Aspergillus*, as it is required not only for ST biosynthesis but also for PN biosynthesis and the biosynthesis of mycelial pigments in *A. nidulans* and gliotoxin and mycelial pigments in *Aspergillus fumigatus*. Furthermore, this protein is required for expression of the heterologous lovastatin (LOV) gene cluster in *A. nidulans* as well as for native LOV expression in *Aspergillus terreus*. Interestingly, the protein appears to be conserved in filamentous fungi, but it is not present in *Saccharomyces cerevisiae*, a fungus devoid of secondary metabolites. Unlike other genes that regulate secondary metabolism, the loss of *laeA* has a negligible impact on morphological developmental processes.

## MATERIALS AND METHODS

**Fungal strains and growth conditions.** Table 1 lists all of the fungal strains used for this study. Some strains are not discussed in the text but were used for sexual crosses to obtain the strains of interest. Sexual crosses of *A. nidulans* strains were conducted according to the method of Pontecorvo et al. (26). All strains were maintained as glycerol stocks and were grown at 37°C for *A. nidulans* and *A. fumigatus* or 32°C for *A. terreus* on glucose minimal medium (GMM) (29), threonine minimal medium (TMM) (29), or lactose minimal medium (LMM) (21) amended with 30 mM cyclopentanone. Threonine and cyclopentanone both induce *alcAp*, which was used to promote *laeA* expression. All media contained appropriate supplements to maintain auxotrophs (2).

**Cloning and sequence of *A. nidulans* and *A. fumigatus laeA* genes.** The *A. nidulans aflR* expression mutant, RYJ8 (derived from strain MRB300), was transformed with an *A. nidulans* genomic cosmid library. Norsolorinic acid-producing transformants were purified, and a cosmid, pCOSJW3, that comple-

\* Corresponding author. Mailing address: Department of Plant Pathology, University of Wisconsin, 1630 Linden Dr., Madison, WI 53706. Phone: (608) 262-9795. Fax: (608) 263-2626. E-mail: npk@plantpath.wisc.edu.

TABLE 1. Fungal strains used for this study

| Strains or category                | Genotype  | Source or reference              |
|------------------------------------|---|----------------------------------|
| <b>Wild type and controls</b>      |   |                                  |
| <i>A. nidulans</i> strains         |   |                                  |
| FGSC 26                            | <i>biA1 veA1</i>  | Fungal Genetics Stock Center     |
| RDIT 2.1                           | <i>methG1</i>   | D. Tsitsigiannis                 |
| RAMB38                             | <i>biA1 methG1 ΔaflR::argB trpC801 veA1</i>                                       | A. M. Bergh                      |
| RDIT 2.3                           | <i>veA1</i>   | D. Tsitsigiannis                 |
| RDIT 7.24                          | <i>methG1 veA1</i>  | D. Tsitsigiannis                 |
| RDIT 30.34                         | <i>methG1 trpC801 pyrG89 veA1</i>   | D. Tsitsigiannis                 |
| RJH26                              | <i>biA1 wA3 argB2 ΔstcE::argB veA1 trpC801</i>                                    | This study                       |
| RJW3                               | <i>pyrG89 wA3 argB2 pyroA4 ΔstcE::argB veA1 trpC801</i>                           | This study                       |
| RJW51                              | <i>alcAp::lovB::pyr4 hygB::lov gene cluster</i>                                   | This study                       |
| RKIS 1                             | <i>pabaA1 yA2 veA1</i>  | 29                               |
| RMFV2                              | <i>pabaA1 yA2 veA1 argB2 ΔaflR::argB</i>  | 12                               |
| TJH3.40                            | <i>biA1 wA3 argB2 methG1 ΔstcE::argB2 veA1</i>                                    | 8                                |
| TJH34.10                           | <i>pabaA1 yA2 veA1 alcA p::aflR::trpC</i>   | 29                               |
| TPK1.1                             | <i>biA1 methG1 veA1</i>   | N. Keller                        |
| WMH1739                            | <i>pabaA1 yA2 alcAp::lovB::pyr4 hygB::lov gene cluster</i>                        | 21                               |
| <i>A. fumigatus</i> strains        |   |                                  |
| AF293                              |   | G. May                           |
| AF293.1                            | <i>pyrG<sup>-</sup></i>   | G. May                           |
| TJW55.2                            | <i>pyrG<sup>-</sup>, A. parasiticus pyrG</i>                                      | This study                       |
| <i>A. terreus</i> strains          |   |                                  |
| ATCC 20542                         |   | American Type Culture Collection |
| TJW58.9                            | <i>hygB</i>   | This study                       |
| <i>laeA</i> mutants                |   |                                  |
| <i>A. nidulans</i> strains         |   |                                  |
| MRB300                             | <i>biA1wA3 methG1 ΔstcE::argB2 veA1 laeA1</i>                                     | 8                                |
| RJW32                              | <i>biA1 wA3 argB2 methG1 ΔstcE::argB veA1 trpC801 ΔlaeA::methG</i>                | This study                       |
| RJW33.2                            | <i>wA3 argB2 methG1 pyroA4ΔstcE::argB veA1 trpC801 ΔlaeA::methG</i>               | This study                       |
| RJW 40.4                           | <i>biA1 methG1 veA1 ΔlaeA::methG</i>  | This study                       |
| RJW44.2                            | <i>biA1 methG1 alcAp::laeA::trpC veA1 ΔlaeA::methG</i>                            | This study                       |
| RJW 46.4                           | <i>methG1 veA1ΔlaeA::methG</i>  | This study                       |
| RYJ 8                              | <i>biA1 wA3ΔstcE::argB veA1 trpC801 laeA1</i>                                     | This study                       |
| RJW 52                             | <i>alcAp::laeA::trpC alcAp::lovB::pyr4 hygB::lov gene cluster</i>                 | This study                       |
| RJW 53                             | <i>ΔlaeA::methG alcAp::lovB::pyr4 hygB::lov gene cluster</i>                      | This study                       |
| TJW 46.16                          | <i>biA1 wA3 argB2 methG1 ΔstcE::argB veA1 alcAp::gfp::laeA::trpC ΔlaeA::methG</i> | This study                       |
| TJW 57.9                           | <i>wA3 argB2 methG1 pyroA4 ΔstcE::argB veA1 aflR::trpCΔlaeA::methG</i>            | This study                       |
| <i>A. fumigatus</i> strain         |   |                                  |
| TJW 54.2                           | <i>ΔlaeA::A. parasiticus pyrG pyrG<sup>-</sup></i>                                | This study                       |
| <i>A. terreus</i> strains          |   |                                  |
| TJW 58.2                           | <i>hygB alcAp::laeA</i>   | This study                       |
| TJW 58.4                           | <i>hygB alcAp::laeA</i>   | This study                       |
| TJW 58.7                           | <i>hygB alcAp::laeA</i>   | This study                       |
| TJW 58.8                           | <i>hygB alcAp::laeA</i>   | This study                       |
| TJW 58.14                          | <i>hygB alcAp::laeA</i>   | This study                       |
| <b>Signal transduction mutants</b> |   |                                  |
| HIFAD4                             | <i>biA1 veA1 fadA<sup>G42R</sup></i>  | 40                               |
| RKIS 11.1                          | <i>pabaA1 yA2 veA1 argB2 ΔfadA::argB</i>  | 29                               |
| RKIS 28.5                          | <i>pabaA1 yA2 veA1 alcAp::rasA17V::argB</i>                                       | 29                               |
| TBN 39.5                           | <i>biA1 methG1 argB2 ΔflbA::argB veA1</i>   | 23                               |
| TJH 34.10                          | <i>pabaA1 yA2 trpC801 trpC::alcA::aflR veA1</i>                                   | J. Hicks                         |
| TKIS 18.11                         | <i>pabaA1 yA2 ΔargB::trpC trpC801 veA1 ΔpkaA::argB</i>                            | 29                               |
| TKIS 20.1                          | <i>pabaA1 yA2 veA1 alcP::pkaA::trpC</i>   | 29                               |
| TSRB 1.38                          | <i>biA1 methG1 argB2 ΔsfaD::argB veA1</i>   | 27                               |

mented the mutation was rescued from one transformant. Norsolorinic acid is an orange pigmented precursor in the ST biosynthetic pathway and is commonly used as an indicator of ST production (8). pJW15, a 4.5-kb KpnI-EcoRI subclone of pCOSJW3, also complemented the mutation and was sequenced by use of synthetic primers and an ABI PRISM DNA sequencing kit (Perkin-Elmer Life Science). The mutant allele *laeA1* was sequenced after subcloning of a 3-kb PCR fragment from RYJ8 genomic DNA amplified with primers LAE1 and LAE2 (Table 2) into the Zero Blunt TOPO vector (Invitrogen) to produce pJW31. Rapid amplification of cDNA ends technology using a Gene Racer kit (Invitro-

gen) was employed to clone *laeA* cDNA according to the manufacturer's instructions. The cloned cDNA was then sequenced. The Institute for Genomic Research (TIGR) database contains a partial *A. fumigatus* genome sequence (<http://www.tigr.org/tdb/e2k1/afu1/>). A putative *A. fumigatus laeA* homolog was obtained by comparing the *A. fumigatus* genomic data with the *A. nidulans laeA* sequence.

**Nucleic acid analysis.** The extraction of DNAs from fungi and bacteria, restriction enzyme digestion, gel electrophoresis, blotting, hybridization, and probe preparation were performed by standard methods (28, 29). Total RNAs were

TABLE 2. Primers

| Primer | Sequence <sup>a</sup>            | Restriction site |
|--------|----------------------------------|------------------|
| LAE1   | ATCTACCTTCTGGGCTCCTGG            |                  |
| LAE2   | CGTGAAGAACTTGGCGTTGTAG           |                  |
| LA2    | GACGAGCTCGTGAACAGTGAAGAAC        | SacI             |
| LA3    | GCGAAGCTTATGAACCGCATCAACCGA      | HindIII          |
| OEF    | GCTGTGAAGCTTTGTACCTGTTTCGCC      | HindIII          |
| OER    | GATTGAAGCTTTGCTGGCATGGAACGG      | HindIII          |
| MT1    | ATGCTGAAGCTTGGAAACTGGGAAAGGGGTC  | HindIII          |
| GFP2   | TGACGAATTCCTTAATGGTTTCTAGCCTG    | EcoRI            |
| GFP31  | TGCGGAATTCATGAGCAAGGGCGAGGAA     | EcoRI            |
| GFP4   | GGATGCCTCGAGTTTGTACAGCTCGTCCATGC | XhoI             |
| GFP5   | AAGCAGCTCGAGTAAGAGCAAAGGGCGACCAC | XhoI             |
| GF1    | CTAGCGAAGCTTGGCCACCATGAGCAAGGGCG | HindIII          |
| GF2    | CGGCGAATTCCTTGTACAGCTCGTCCATGC   | EcoRI            |
| GF3    | TTTGGAAATTCGTTTCGCCGCTGATGTTTGG  | EcoRI            |
| FUM1   | GCGCACTTCTTTGTTTCCCTT            |                  |
| FUM2   | CATCGGAATTCCTTTCTTGAGCGGGC       | EcoRI            |
| FUM3   | TACCAGGATCCAAAACCTCTCGCCA        | BamHI            |
| FUM4   | CATGACGGTAACTAAGGATTTGG          |                  |

<sup>a</sup> Underlined sequences show the placement of restriction sites.

extracted from *Aspergillus* strains by use of Trizol reagent (Invitrogen) according to the manufacturer's instructions. RNA blots were hybridized with a 0.7-kb SacII-KpnI fragment from pRB7 containing the *stcU* coding region (18), a 1.3-kb EcoRV-XhoI fragment from pJW19 containing the *afIR* coding region, a 3-kb HindIII fragment from pJW45.4 containing the *laeA* coding region, a 1.1-kb EcoRI-HindIII fragment from pUCHH(458) containing the *ipnA* coding region (36), a 5-kb BamHI fragment from pWHM1401 containing the *lovE* coding region (21), and a 1.3-kb PCR product from pWHM1263 containing the *lovC* coding region (21). Also, *A. nidulans* cosmids pW07H03, pL11C09, and pL24B03 were used as probes. pL11C09 contains most of the ST gene cluster, whereas pW07H03 and pL24B03 primarily contain genes located upstream and downstream of the ST gene cluster, respectively (7).

**Construction of transformation vectors and strains.** Plasmids were generated by standard techniques, and the primers used for this study are listed in Table 2. *Pfu* Turbo (Stratagene) was used for PCRs. The *A. nidulans* disruption plasmid pJW34 was constructed by ligating a 1.2-kb DNA fragment upstream of the *laeA* start codon (primers LAE1 and LA2) and a 1.2-kb DNA fragment downstream of the *laeA* stop codon (primers LA3 and LAE2) to either side of the *methG* gene in the pUG11-41 vector (31). The 5'-end PCR product and 3'-end PCR product were inserted into the SacI site and HindIII site of pUG11-41, respectively, by blunt end ligation. pJW34 was used to disrupt the *laeA* gene ( $\Delta laeA$ ) in strain TJH3.40 to create TJW35.5. TJW35.5 was subsequently sexually crossed to RDIT2.1 to create RJW46.4. Plasmid pJW47.4 was constructed to overexpress *laeA* from the *alcA* promoter. The 2.5-kb coding sequence of *laeA* was amplified with primers OEF and OER and ligated into the HindIII site of pCN2, which contains the 5' half of the *trpC* gene and the *alcA* promoter (19). This resulted in an *alcAp::laeA* fusion, referred to as *OE::laeA* hereafter. pJW47.4 was used to transform RJW32 to tryptophan auxotrophy to yield the strain TJW44.39. TJW44.39 was subsequently sexually crossed to RDIT2.1 to create RJW47.3. pJW47.4 and a hygromycin B (*hygB*) resistance gene in plasmid pUCH2-8 (3) were used for cotransformation to introduce the overexpression *laeA* construct into *A. terreus* ATCC 20542. Transformants were selected in hygromycin B (500  $\mu$ g/ml)-containing medium and confirmed by PCR and Southern hybridization. Five transformants, namely TJW58.2, TJW58.4, TJW58.7, TJW58.8, and TJW58.14, containing *hygB* and *OE::laeA*, were examined for LOV production, and TJW58.9 containing *hygB* alone was used as a control (Table 1). pJW45.4, containing a wild-type copy of the *laeA* gene, was used to complement the  $\Delta laeA$  strain RJW33.2. pJW45.4 was created by ligating the 3-kb *laeA* gene (primers MT1 and OER) into the HindIII site of pSH96. pSH96 contains the 5' half of the *trpC* gene (39). RJW33.2 is a sexual progeny of a cross between strains TJW35.5 and RJW3. pJW45.4 was used to transform RJW33.2 to produce TJW42.7. TJW42.7 was crossed with RDIT7.24 sexually to create RJW49.1. Plasmids pJW48 and pJW49 were created to visualize *LaeA* by fusing of the green fluorescent protein (GFP) gene (10, 13) to the N-terminal and C-terminal ends, respectively, of *LaeA*. pJW48 was made by ligating the 0.7-kb *gfp* gene (primers GF1 and GF2) to the 5' end of the 2.5-kb encoding region of the *laeA* gene (primers GF3 and OER) and then inserting the ligated fragment into the pCN2 HindIII site to yield an *alcAp::gfp::laeA* chimera. pJW49 was constructed by

consecutively ligating a 2-kb *laeA* coding region (primers OEF and GFP2), a 0.7-kb *gfp* gene (primers GFP31 and GFP4), and a 0.5-kb *laeA* termination cassette (primers GFP5 and OER) into the HindIII site of pCN2 to yield an *alcAp::laeA::gfp::laeAterm* chimera. pJW48 and pJW49 were used to transform RJW32 to yield transformants TJW46.16 (5' GFP) and TJW47.9 (3' GFP), respectively. The *A. fumigatus laeA* gene disruption vector, pJW58, was constructed by insertion of a 0.9-kb DNA fragment upstream of the *laeA* start codon (primers FUM1 and FUM2) and a 1.0-kb DNA fragment downstream of the *laeA* stop codon (primers FUM3 and FUM4) on either side of the *Aspergillus parasiticus pyrG* marker gene obtained from pBZ5 (32). pJW58 was used to disrupt the *A. fumigatus laeA* gene in strain AF293.1 to create strain TJW54.2.

**Fungal transformation procedures.** Fungal transformation was done essentially as described by Miller et al. (24), with the modification of embedding the protoplasts in top agar (0.75%) rather than spreading them by a glass rod on solid medium.

**Secondary metabolite analysis.** Published procedures were used to extract and analyze ST (12), gliotoxin (5), LOV (21), and monocolin J (MONJ) (21). All metabolites were extracted with chloroform from both mycelial and culture filtrates. MONJ and LOV concentrations in each strain were estimated by comparisons to standard spots on thin-layer chromatography (TLC) plates by dilution spotting. Pictures of TLC plates were taken at 254 nm. ST was extracted from either 50-ml shake cultures in GMM inoculated with  $10^7$  spores/ml and grown for 60 h or solid medium cultures spread with  $10^6$  spores/plate and grown for 5 days. Dried ST extracts were resuspended in 100  $\mu$ l of chloroform, and 10  $\mu$ l was separated in chloroform-acetone (8:2) on TLC plates. ST (Sigma) was spotted as a standard. MONJ was extracted from 50-ml GMM shake cultures inoculated with  $10^7$  spores/ml and grown for 72 h. MONJ from WT/*lov+* and *OE::laeA/lov+* strains was extracted from cultures grown in 50 ml of shaking liquid GMM for 14 h at 37°C and then transferred to shaking liquid TMM for 24 h. Dried MONJ extracts were resuspended in 100  $\mu$ l of methanol, and 10  $\mu$ l was separated in methanol-0.1% phosphoric acid (9:1) on C<sub>18</sub> reversed-phase TLC plates. The MONJ standard was extracted from *A. nidulans* strain WMH1739 (Table 1). All experiments were performed in triplicate. Gliotoxin production in *A. fumigatus* was analyzed by modification of the TLC method of Belkacemi et al. (5). Gliotoxin was extracted from 50-ml GMM shake cultures inoculated with  $10^7$  spores/ml and grown for 3 days. Dried chloroform extracts were resuspended in 100  $\mu$ l of methanol, and 10  $\mu$ l was separated in chloroform-methanol (9:1). Gliotoxin (Sigma) was spotted as a standard. All experiments were performed in triplicate. For the assessment of PN production, *Micrococcus luteus* ATCC 9341 was grown in TBS (17 g of Bacto Tryptone, 3 g of Bacto Soytone, 5 g of NaCl, 2.5 g of K<sub>2</sub>HPO<sub>4</sub>, and 2.5 g of glucose in a 1-liter total volume) at 37°C at 180 rpm until it reached an optical density of 1. Three and one-third milliliters of *M. luteus* culture was mixed with 40 ml of TSA (15 g of Bacto Tryptone, 5 g of Bacto Soytone, 5 g of NaCl, and 10 g of agar in a 1-liter total volume) and poured into 150-cm-diameter plates to solidify. Fifty-milliliter cultures of the wild-type (WT),  $\Delta laeA$ , and *OE::laeA* strains ( $10^7$  spores/ml) were grown in GMM with shaking for 14 h at 37°C and then transferred to LMM shake cultures amended with 30 mM cyclopentanone for 24 h. For each strain, 6 ml was removed, lyophilized, and resuspended in 1 ml of distilled water. One-hundred-microliter samples, with or without 6 U of  $\beta$ -lactamase, were placed in 10-cm-diameter wells of the *M. luteus* plates. Plates were placed for 2 h at 4°C and then incubated overnight at 37°C to evaluate PN inhibition zones. All experiments were duplicated. LOV was extracted from *A. terreus* cultures grown in 50-ml GMM shake flasks for 18 h at 32°C and then transferred to LMM shake cultures with 30 mM cyclopentanone for 36 h at 32°C. Extraction and identification on TLC were followed by the previously described method of MONJ examination. LOV (Merck Co.) was spotted onto TLC plates as a control. All experiments were duplicated.

**Nucleotide sequence accession numbers.** GenBank numbers for the *A. nidulans* and *A. fumigatus laeA* genes are AY394722 and AY422723, respectively.

## RESULTS

**Cloning and characterization of *A. nidulans* and *A. fumigatus laeA*.** A mutagenesis screen previously led to the isolation of 23 mutants displaying a loss of ST production but normal sporulation in *A. nidulans* (8). Three of the mutants were unable to express *afIR*, which encodes an ST cluster Zn<sub>2</sub>Cys<sub>6</sub> transcription factor regulating ST biosynthetic gene expression (12). We were able to complement one of these three mutants, RYJ8, with an *A. nidulans trpC* genomic cosmid library. Sequencing of




FIG. 1. Overview of *LaeA*. (A) Schematic of *laeA* gene. Although *LaeA* contains the exact SAM motif found in histone methyltransferases and arginine methyltransferases, it lacks other conserved domains (e.g., a SET domain and a double E loop) typically found in these proteins. In addition, likely histone methyltransferase and arginine methyltransferase candidates are found in the *Aspergillus* database ( $1e^{-42}$  and  $1e^{-94}$ ). Therefore, *LaeA* appears to be a unique protein methyltransferase. (B) Amino acid comparison of *A. nidulans*, *A. fumigatus*, *N. crassa*, *Magnaporthe grisea*, *C. immitis*, and *F. sporotrichioides* *LaeA* proteins showing conserved protein methyltransferase SAM binding sites in red. (C) *A. nidulans* *LaeA* protein localizes to the nucleus. GFP was fused to the N-terminal end of *LaeA*. Nuclei were stained with the DNA-specific dye 4,6-diamidino-2-phenylindole (DAPI).

a 4.5-kb subclone (pJW15) of the complementing cosmid pCOSJW3 revealed a 3-kb open reading frame designated *laeA* (for loss of *affR* expression). Sequencing of the mutant allele, *laeA1*, from RJY8 showed that it has a base pair transversion (at position 1455; C→G) and a 1-bp deletion (at position 1453) in the gene. The deletion resulted in a premature stop codon. An examination of genomic and cDNA sequences revealed that *laeA* has one intron and three putative *AflR* binding sites (12), one in the promoter (−607) and two in the encoding region (positions 607 and 1487) (Fig. 1A). cDNA analysis showed that *laeA* possesses a 5' untranslated region (642 bp) (Fig. 1A). Analyses of available genomic databases indicated that only filamentous fungi, including *A. fumigatus* (human pathogen causing aspergillosis; TIGR [http://www.tigr.org/tdb/e2k1/afu1/]), *Neurospora crassa* (model fungus; GenBank), *Magnaporthe grisea* (plant pathogen causing rice blast fungus [http://www-genome.wi.mit.edu/annotation/fungi/magnaporthe]), *Coccidioides immitis* (human pathogen causing coccidioidomycosis; GenBank), and *Fusarium sporotrichioides* (plant pathogen producing trichothecene mycotoxin [http://www.genome.ou.edu/fsporo.html]), have possible *LaeA* homologs (Fig. 1B). An examination of the *LaeA* amino acid sequence (375 amino acids) revealed a conserved *S*-adenosylmethionine (SAM) binding site found in nuclear protein methyltransferases (Fig. 1B) (15). Although the amino acid sequence of *LaeA* did not show the presence of a nuclear localization motif, GFP tagging

of either the 5' or 3' end of *A. nidulans laeA* showed *LaeA* to be primarily localized in the nucleus (Fig. 1C).

***LaeA* is required for secondary metabolite production.** *laeA* null mutants ( $\Delta laeA$ ) were created by replacing *laeA* with *methG* and *pyrG* in *A. nidulans* TJH3.40 (a *methG1* auxotroph) and *A. fumigatus* AF293.1 (a *pyrG* auxotroph), respectively. Southern blot and PCR analyses were carried out to confirm single gene replacement events in several transformants, including *A. nidulans* TJW35.5 and *A. fumigatus* TJW54.2. Prototroph RJW46.4 was obtained from TJW35.5 by a sexual cross, as described in Materials and Methods. Strains RJW46.4 and TJW54.2 were used for our study. For both species,  $\Delta laeA$  strains were visually detectable due to the loss of mycelial pigment from the backside of the colonies (Fig. 2A and data not shown). A TLC examination of chloroform extracts of *A. nidulans* and *A. fumigatus*  $\Delta laeA$  strains showed that the production of several metabolites, including ST in *A. nidulans* (Fig. 2B) and the immunotoxin gliotoxin in *A. fumigatus*, was reduced (Fig. 2C). The identities of the other metabolites are not known. Interestingly, the levels of two of the *A. nidulans* metabolites were not reduced and even appeared to be somewhat increased in the  $\Delta laeA$  strain (Fig. 2B). Verification that these defects were caused by the loss of *laeA* was obtained by the transformation of *A. nidulans*  $\Delta laeA$  with wild-type *laeA* and the observed remediation of metabolite production (data not shown).




FIG. 2. Phenotypes of *laeA*. (A) Asexual sporulation (top) and mycelial pigmentation (bottom) patterns of *A. nidulans* wild-type (RDIT2.3) (WT) and  $\Delta laeA$  (RJW46.4) strains after 5 days of cultivation on GMM. *A. fumigatus*  $\Delta laeA$  presented a similar loss of mycelial pigmentation (data not shown). (B) TLC analysis of chloroform extracts of RDIT2.3 and RJW46.4 after 5 days of cultivation on solid GMM. (C) TLC analysis of chloroform extracts of *A. fumigatus* AF293 (WT) and TJW54.2 ( $\Delta laeA$ ) grown in liquid shaking GMM for 3 days. The experiment was performed in triplicate. Std, gliotoxin standard.

### Transcriptional regulation of secondary metabolism gene clusters by *LaeA*.

(i) **Native cluster regulation.** To confirm our initial observation that *laeA* is required for ST gene regulation, we assessed *aflR* and *stcU* (a gene encoding a biosynthetic enzyme required for ST production) (17) expression in the  $\Delta laeA$  background. Neither gene was expressed (Fig. 3A). A transcriptional profile of the entire ST gene cluster, which covers ca. 60 kb and contains ca. 26 genes (7), suggested that *LaeA* transcriptional control is ST cluster specific, as transcription of the upstream and downstream genes of the ST cluster was unaffected (data not shown). Because many uncharacterized metabolites were reduced in the  $\Delta laeA$  strains (Fig. 2B and C), we thought it possible that *LaeA* is a global regulator for secondary metabolite gene expression. To address this hypothesis, we examined PN gene expression in the *A. nidulans*  $\Delta laeA$  strain. Figure 3A shows that *ipnA* (encoding isopenicillin N synthetase, a biosynthetic enzyme required for PN biosynthesis) (6) expression was greatly reduced in the  $\Delta laeA$  strain.

(ii) **Heterologous cluster regulation.** Our results for ST and PN gene expression suggested a role for *LaeA* in secondary metabolite gene cluster regulation. To further address this potential role, we examined the expression of the heterologous LOV gene cluster in the *A. nidulans*  $\Delta laeA$  background. The partial LOV cluster, derived from *A. terreus*, was originally transformed into *A. nidulans* to study aspects of LOV biosynthesis (21). This strain was used to cross the LOV cluster (LOV<sup>+</sup>) into appropriate mutant *laeA* backgrounds. Figure 3B shows that the  $\Delta laeA$ /LOV<sup>+</sup> strain displayed very diminished levels of both *lovE* (encoding a LOV-specific Zn<sub>2</sub>Cys<sub>6</sub> transcription factor) and *lovC* (a LOV biosynthetic gene) transcripts. Chloroform extracts of this strain also showed diminished production of MONJ (Fig. 3B), the LOV intermediate produced by the partial LOV cluster (21).

**Overexpression of *laeA* upregulates PN and LOV gene expression but not ST gene expression.** We next constructed *laeA* overexpression strains (*OE::laeA*) of both *A. nidulans* and *A. terreus* to examine secondary metabolite gene expression and product formation. As shown in Fig. 3C, *ipnA*, *lovE*, and *lovC*, but not *stcU*, expression levels were remarkably elevated in the *A. nidulans* *OE::laeA* background. Secondary metabolite production was correlated with transcript levels. MONJ pro-

duction was increased ~400%, and high levels of PN were produced during times when the wild type showed no PN activity (Fig. 3D). ST levels remained the same as that of the wild type in the *OE::laeA* background (Fig. 3C). Overexpression of the *A. nidulans* *laeA* gene in the LOV-producing fungus *A. terreus* led to 400 to 700% increases in LOV (Fig. 3E).

**Feedback regulation of *laeA* by *aflR*.** An examination of the *laeA* transcript in wild-type strains showed that it is an inducible, low-expression-level gene that is observed in Northern blots before and after *aflR* transcripts are observed (data not shown). Due to the presence of three *AflR* binding sites in the gene (Fig. 1A), we thought it possible that *AflR* regulates *laeA* expression. As shown in Fig. 4, overexpression of *aflR* (*OE::aflR*) downregulates *laeA* expression, although elimination of *aflR* ( $\Delta aflR$ ) does not affect the *laeA* transcript level (Fig. 5). This indicates that there are both negative (*laeA*) and positive (*stc* genes) regulatory effects of *AflR* on gene transcription.

**Protein kinase A and *RasA* negatively regulate *laeA* expression.** ST biosynthesis is regulated in *A. nidulans* via a signal transduction pathway, and many of the genes involved in this signaling pathway are known (9). Therefore, we looked at the possible interactions with *laeA* of five signaling genes, encoding two members of a heterotrimeric G protein (*fadA* and *sfadD*) (27, 40), a regulator of G-protein signaling protein regulating *FadA* activity (*flbA*) (23), a cAMP-dependent kinase (*pkaA*) (29), and a Ras protein (*rasA*) (33). *laeA* expression was examined in the wild type and in strains carrying the following alleles:  $\Delta flbA$ , *fadA*<sup>G42R</sup>,  $\Delta sfadD$ ,  $\Delta sfadD$ ,  $\Delta pkaA$ , *OE::pkaA*, and *OE::rasA*<sup>G17A</sup> (Table 1). mRNA analyses of these mutants showed that *OE::pkaA* and *OE::rasA*<sup>G17A</sup> completely inhibited *laeA* expression (Fig. 4), whereas *laeA* transcription was not repressed in any of the other strains (Fig. 5 and data not shown). Interestingly, the *laeA* transcript level was elevated in the  $\Delta sfadD$  strain (Fig. 5). The presence of *laeA* transcripts in these mutants (Fig. 5 and data not shown) shows that *laeA* is not sufficient for *aflR* expression, as *aflR* was not expressed in these strains (18). Figure 6 depicts our current understanding of *LaeA* involvement in secondary metabolite regulation.




FIG. 3. *laeA* regulation of secondary metabolism. (A) *aflR*, *stcU*, and *ipnA* gene expression in *A. nidulans* wild-type (WT) (RDIT2.3) and  $\Delta laeA$  (RJW46.4) strains grown in liquid shaking GMM for 12, 24, 48, and 72 h at 37°C. Ethidium bromide-stained rRNA is indicated as a loading control. (B) *laeA*, *lovE*, and *lovC* gene expression in *A. nidulans* WT/*lov+* (RJW51) and  $\Delta laeA$ /*lov+* (RJW53) strains grown in liquid shaking GMM for 12, 24, 48, and 72 h at 37°C. Ethidium bromide-stained rRNA is indicated as a loading control. MONJ was extracted from WT/*lov+* (RJW51) and  $\Delta laeA$ /*lov+* (RJW53) *A. nidulans* strains grown in liquid shaking GMM for 3 days. The experiment was performed in triplicate. (C) *laeA*, *ipnA*, *stcU*, *lovE*, and *lovC* gene expression in *A. nidulans* wild-type (WT) (RDIT2.3), OE::*laeA* (RJW47.3), WT/*lov+* (RJW51), and OE::*laeA*/*lov+* (RJW52) strains grown in liquid shaking GMM for 14 h at 37°C and then transferred to liquid shaking TMM for the induction of *laeA* expression. Time points were 0, 6, 12, and 24 h after transfer. ST and MONJ were extracted from *A. nidulans* WT (RDIT2.3), OE::*laeA* (RJW47.3), WT/*lov+* (RJW51), and  $\Delta laeA$ /*lov+* (RJW52) strains grown in liquid shaking GMM for 14 h at 37°C and then transferred to liquid shaking TMM for 24 h. ST, ST standard. The MONJ standard was extracted from *A. nidulans* strain WMH1739. The experiment was performed in triplicate. (D) PN bioassay. Wild-type (FGSC26),  $\Delta laeA$  (RJW40.4), and OE::*laeA* (RJW44.2) strains were grown in liquid shaking GMM for 14 h at 37°C and then transferred to LMM amended with 30 mM cyclopentanone for the induction of *laeA* for 24 h at 37°C. (E) TLC examination of LOV production in *A. terreus laeA* overexpression strains. The wild type (ATCC 20542; lane 1), TJW58.9 (*hygB* resistance gene-containing transformant used as a control; lane 2), and OE::*laeA* strains containing *hygB* (TJW58.2, TJW58.4, TJW58.7, TJW58.8, and TJW58.14, in lanes 3 to 7, respectively) were grown in liquid shaking GMM for 18 h at 32°C and then transferred to LMM with 30 mM cyclopentanone for the induction of *laeA* for 36 h at 32°C. Std, LOV standard. The experiment was performed in duplicate.




FIG. 4. Regulation of *laeA*. Effects of overexpression of *afIR*, *pkaA*, and *ras<sup>G17V</sup>* on *laeA* expression. Wild-type (RKIS1), *OE::afIR* (TJH34.10), *OE::pkaA* (TKIS20.1), and *OE::ras<sup>G17V</sup>* (RKIS28.5) strains were grown in liquid shaking GMM for 14 h at 37°C and then transferred to TMM. Time points were 0, 6, 12, and 24 h after transfer.

## DISCUSSION

Secondary metabolite biosynthesis is often associated with the advent of sporulation and cellular development in filamentous fungi (9) and filamentous *Streptomyces* bacteria (38). These developmental processes reflect the need to access multiple nutrients and to optimize cellular morphology and metabolic differentiation for effective competition in complex environments. We are interested in the identification of secondary metabolism-specific global regulators that can uncouple sporulation and secondary metabolism. Such regulatory elements are extremely desirable because they would possess broad specificity for the activation and/or repression of entire families of secondary metabolite gene clusters while providing strains that are capable of otherwise normal or near-normal development and growth. The identification of such regulatory elements would enable the increased production of secondary metabolites by providing improved strains of engineered organisms and would also contribute to a broader understanding of molecular mechanisms by which secondary metabolites are produced. Through complementation of an ST developmental mutant, we have identified such a protein, called LaeA, which is an archetypal global regulator of secondary metabolism in fungi.

LaeA regulation of metabolite production is transcriptional, as assessed by the effects of  $\Delta laeA$  and *OE::laeA* alleles on ST, PN, and LOV gene expression in *A. nidulans* and *A. terreus*. In all cases, gene transcripts were reduced or eliminated in  $\Delta laeA$  strains. However, although overexpression of *laeA* increased PN and LOV gene transcript and concomitant production formation (Fig. 3C, D, and E), this was not the case for ST gene transcription or production (Fig. 3A). The steady-state level of ST transcripts and product formation in the *OE::laeA* background, in contrast to the increased PN and LOV transcripts

and product formation, suggested a unique interaction between *laeA* and ST gene regulation.

Due to the presence of three potential AfIR binding sites in the *A. nidulans* gene (Fig. 1A) and the lack of ST cluster gene upregulation in the *OE::laeA* background, we thought it possible that AfIR negatively regulates *laeA* expression. As shown in Fig. 4, overexpression of *afIR* (*OE::afIR*) downregulates *laeA* expression, although elimination of *afIR* ( $\Delta afIR$ ) does not affect *laeA* transcript levels (Fig. 5). This indicates that there are both negative (*laeA*) and positive (*stc* genes) (12) regulatory effects of AfIR on gene transcription. To our knowledge, this is the first description of a putative secondary metabolite feedback mechanism. As overproduction of ST negatively affects fungal growth (N. P. Keller, unpublished data), we speculate that this feedback loop may have evolved as a fitness trait. In contrast, neither the promoter nor the encoding region of *A. fumigatus laeA* contained AfIR binding sites, and no *afIR* ortholog was found in the genome. Some *A. fumigatus* strains are reported to produce ST (14); it would be interesting to see if *laeA* genes from those isolates contained AfIR binding sites. Initial examinations of the *Aspergillus*  $\Delta laeA$  strains showed them to be more susceptible to killing by neutrophils in vitro than the wild type (S. Balajee, L. Delbridge, J. Bok, N. Keller, and K. Marr, unpublished data). Presumably this is due to a loss of toxin secondary metabolites or melanins, known virulence factors in several fungal systems (5, 22).

*laeA* expression is also negatively regulated by two signal transduction molecules, PkaA and RasA. Both proteins have been shown to transcriptionally and posttranscriptionally regulate *afIR* (29, 30). It appears that LaeA mediates PkaA transcriptional regulation of *afIR*, since overexpression of *laeA* in a *pkaA* overexpression background, a condition that normally suppresses *stc* expression, partially restored *stc* gene expression




FIG. 5. *laeA* expression is not affected in  $\Delta flbA$ ,  $\Delta sfaD$ , and  $\Delta afIR$  strains. *laeA*, *afIR*, and *stcU* gene expression was examined in *A. nidulans* wild-type (TPK1.1 and RKIS1),  $\Delta flbA$  (TBN39.5),  $\Delta sfaD$  (TSRB1.38), and  $\Delta afIR$  (RMFV2) strains grown in liquid shaking GMM for 12, 24, 48, and 72 h at 37°C. Ethidium bromide-stained rRNA is indicated as a loading control.




FIG. 6. Proposed model of secondary metabolite regulation by LaeA. Gliotoxin is formed from serine and alanine and is proposed to be produced by a nonribosomal peptide synthase (NRPS). Fungal pigments belong to several chemical classes, including polyketides, terpenes, and DOPA melanins.

(29; data not shown). With regard to PkaA regulation of *laeA*, the lack of conventional PkaA phosphorylation consensus sequences in LaeA indicates that PkaA regulation of LaeA is not direct. Alternatively, LaeA may contain unconventional PkaA phosphorylation sites. RasA regulation of *laeA* gene expression may occur through PkaA and/or another pathway(s).

The requirement of a kinase for *laeA* function is reminiscent—to a degree—of a *Streptomyces* global regulatory system involving the protein AfsR. AfsR is a transcription factor that regulates secondary metabolism in *Streptomyces coelicolor* but regulates morphogenesis in *Streptomyces griseus* (contrast this to the similar role LaeA has in the three *Aspergillus* spp. examined here). Phosphorylation of AfsR enhances its activity (37). Like the case for AfsR, LaeA regulation occurs at the transcriptional level, but it shows no homology to transcription factors. Its nuclear location, its role in transcriptional regulation, and the presence of a SAM motif in LaeA suggest that it may be a protein methyltransferase. Well-known protein methyltransferases include histone and arginine methyltransferases that play important roles in the regulation of gene expression in eukaryotes, in part through modification of the chromatin structure (15, 25, 35).

Regardless of the mechanism, these findings with LaeA present an advance toward understanding the complex regu-

lation of secondary metabolite production and provide a means for the discovery of new metabolites. Indeed, initial comparative microarray studies between  $\Delta laeA$  and the wild type have identified putative secondary metabolism gene clusters in the *Aspergillus* genome (L. Maggio-Hall, J. Bok, and N. Keller, unpublished data). The manipulation of LaeA in filamentous fungi may enable the increased production of pharmaceuticals or the elimination of fungal toxins by providing improved strains of engineered organisms and may also contribute to the broader understanding of molecular mechanisms by which secondary metabolites are produced.

#### ACKNOWLEDGMENT

This research was funded by NSF grant MCB-9874646 to N.P.K.

#### REFERENCES

- Adams, T. H., and J. H. Yu. 1998. Coordinate control of secondary metabolite production and asexual sporulation in *Aspergillus nidulans*. *Curr. Opin. Microbiol.* **1**:674–677.
- Adams, T. H., and W. E. Timberlake. 1990. Developmental repression of growth and gene expression in *Aspergillus*. *Proc. Natl. Acad. Sci. USA* **87**: 5405–5409.
- Alexander, N. J., T. H. Hohn, and S. P. McCormick. 1998. The TR111 gene of *Fusarium sporotrichioides* encodes a cytochrome P-450 monooxygenase required for C-15 hydroxylation in trichothecene biosynthesis. *Appl. Environ. Microbiol.* **64**:221–225.
- Alspaugh, J. A., J. R. Perfect, and J. Heitman. 1997. *Cryptococcus neoformans*.



- mans* mating and virulence are regulated by the G-protein alpha subunit GPA1 and cAMP. *Genes Dev.* **11**:3206–3217.
5. **Belkacemi, L., R. C. Barton, V. Hopwood, and E. G. V. Evans.** 1999. Determination of optimum growth conditions for gliotoxin production by *Aspergillus fumigatus* and development of a novel method for gliotoxin detection. *Med. Mycol.* **37**:227–233.
  6. **Brakhage, A. A.** 1998. Molecular regulation of beta-lactam biosynthesis in filamentous fungi. *Microbiol. Mol. Biol. Rev.* **62**:547–585.
  7. **Brown, D. W., J. H. Yu, H. S. Kelkar, M. Fernandes, T. C. Nesbitt, N. P. Keller, T. H. Adams, and T. J. Leonard.** 1996. Twenty-five coregulated transcripts define a sterigmatocystin gene cluster in *Aspergillus nidulans*. *Proc. Natl. Acad. Sci. USA* **93**:1418–1422.
  8. **Butchko, R. A., T. H. Adams, and N. P. Keller.** 1999. *Aspergillus nidulans* mutants defective in *stc* gene cluster regulation. *Genetics* **153**:715–720.
  9. **Calvo, A. M., R. A. Wilson, J. W. Bok, and N. P. Keller.** 2002. Relationship between secondary metabolism and fungal development. *Microbiol. Mol. Biol. Rev.* **66**:447–459.
  10. **Chalfie, M., Y. Tu, G. Euskirchen, W. W. Ward, and D. C. Prasher.** 1994. Green fluorescent protein as a marker for gene expression. *Science* **263**:802–805.
  11. **Demain, A., and A. Fang.** 2000. The natural functions of secondary metabolites. *Adv. Biochem. Eng. Biotechnol.* **69**:1–39.
  12. **Fernandes, M., N. P. Keller, and T. H. Adams.** 1998. Sequence-specific binding by *Aspergillus nidulans* AfIR, a C6 zinc cluster protein regulating mycotoxin biosynthesis. *Mol. Microbiol.* **28**:1355–1365.
  13. **Fernandez-Abalos, J. M., H. Fox, C. Pitt, B. Wells, and J. H. Doonan.** 1998. Plant-adapted green fluorescent protein is a versatile vital reporter for gene expression, protein localization and mitosis in the filamentous fungus *Aspergillus nidulans*. *Mol. Microbiol.* **27**:121–130.
  14. **Hah, H.** 1993. Differential action of cercoran and topsin-M on sensitive and tolerant strains of toxic fungi. *Cryptogamie Mycologie* **14**:61–68.
  15. **Hamahata, A., Y. Takata, T. Gomi, and M. Fujioka.** 1996. Probing the S-adenosylmethionine-binding site of rat guanidinoacetate methyltransferase. Effect of site-directed mutagenesis of residues that are conserved across mammalian non-nucleic acid methyltransferases. *Biochem. J.* **317**:141–145.
  16. **Hawksworth, D. L.** 2001. The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycol. Res.* **105**:1422–1432.
  17. **Hicks, J. K., K. Shimizu, and N. P. Keller.** 2002. Genetics and biosynthesis of aflatoxins and sterigmatocystin, p. 55–69. *In* F. Kempken and J. W. Bennett (ed.), *The mycota*, vol. XI. Springer-Verlag, Berlin, Germany.
  18. **Hicks, J. K., J.-H. Yu, N. P. Keller, and T. H. Adams.** 1997. *Aspergillus* sporulation and mycotoxin production both require inactivation of the FadA G alpha protein-dependent signaling pathway. *EMBO J.* **16**:4916–4923.
  19. **Hicks, J., R. A. Lockington, J. Strauss, D. Dieringer, C. P. Kubicek, J. Kelly, and N. Keller.** 2001. RcoA has pleiotropic effects on *Aspergillus nidulans* cellular development. *Mol. Microbiol.* **39**:1482–1493.
  20. **Jin, Y., J. W. Bok, D. Guzman-de-Peña, and N. P. Keller.** 2002. Requirement of spermidine for developmental transitions in *Aspergillus nidulans*. *Mol. Microbiol.* **46**:801–812.
  21. **Kennedy, J., K. Auclair, S. G. Kendrew, C. Park, J. C. Vederas, and C. R. Hutchinson.** 1999. Modulation of polyketide synthase activity by accessory proteins during lovastatin biosynthesis. *Science* **284**:1368–1372.
  22. **Langfelder, K., M. Streibel, B. Jahn, G. Haase, and A. A. Brakhage.** 2003. Biosynthesis of fungal melanins and their importance for human pathogenic fungi. *Fungal Genet. Biol.* **38**:143–158.
  23. **Lee, B. N., and T. H. Adams.** 1994. Overexpression of *flbA*, an early regulator of *Aspergillus* asexual sporulation, leads to activation of *brlA* and premature initiation of development. *Mol. Microbiol.* **14**:323–334.
  24. **Miller, B. L., K. Y. Miller, and W. E. Timberlake.** 1985. Direct and indirect gene replacements in *Aspergillus nidulans*. *Mol. Cell. Biol.* **5**:1714–1721.
  25. **Mowen, K. A., J. Tang, W. Zhu, B. T. Schurter, K. Shuai, H. R. Herschman, and M. David.** 2001. Arginine methylation of STAT1 modulates IFN alpha/beta-induced transcription. *Cell* **104**:731–741.
  26. **Pontecorvo, G., J. A. Roper, L. M. Hemmons, K. P. Macdonald, and A. W. J. Bufton.** 1953. The genetics of *Aspergillus nidulans*. *Adv. Genet.* **5**:141–238.
  27. **Rosén, S., J.-H. Yu, and T. H. Adams.** 1999. The *Aspergillus nidulans* *sfaD* gene encodes a G protein beta subunit that is required for normal growth and repression of sporulation. *EMBO J.* **18**:5592–5600.
  28. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
  29. **Shimizu, K., and N. P. Keller.** 2001. Genetic involvement of a cAMP-dependent protein kinase in a G protein signaling pathway regulating morphological and chemical transitions in *Aspergillus nidulans*. *Genetics* **157**:591–600.
  30. **Shimizu, K., J.-K. Hicks, T.-P. Huang, and N. P. Keller.** 2003. Pka, Ras and RGS protein interactions regulate activity of AfIR, a Zn(II)<sub>2</sub>Cys<sub>6</sub> transcription factor in *Aspergillus nidulans*. *Genetics* **165**:1095–1104.
  31. **Sienko, M., and A. Paszewski.** 1999. The *metG* gene of *Aspergillus nidulans* encoding cystathionine beta-lyase: cloning and analysis. *Curr. Genet.* **35**:638–646.
  32. **Skory, C. D., P. K. Chang, J. Cary, and J. E. Linz.** 1992. Isolation and characterization of a gene from *Aspergillus parasiticus* associated with the conversion of versicolorin A to sterigmatocystin in aflatoxin biosynthesis. *Appl. Environ. Microbiol.* **58**:3527–3537.
  33. **Som, T., and V. S. Kolaparthi.** 1994. Developmental decisions in *Aspergillus nidulans* are modulated by Ras activity. *Mol. Cell. Biol.* **14**:5333–5348.
  34. **Tag, A., J. Hicks, G. Garifullina, M. Beremand, and N. Keller.** 2000. G-protein signalling mediates differential production of toxic secondary metabolites. *Mol. Microbiol.* **38**:658–665.
  35. **Tamaru, H., and E. U. Selker.** 2001. A histone H3 methyltransferase controls DNA methylation in *Neurospora crassa*. *Nature* **414**:277–283.
  36. **Tilburn, J., S. Sarkar, D. A. Widdick, E. A. Espeso, M. Orejas, J. Mungroo, M. A. Penalva, and H. N. Arst, Jr.** 1995. The *Aspergillus* PacC zinc finger transcription factor mediates regulation of both acid- and alkaline-expressed genes by ambient pH. *EMBO J.* **14**:779–790.
  37. **Umeyama, T., P. C. Lee, and S. Horinouchi.** 2002. Protein serine/threonine kinases in signal transduction for secondary metabolism and morphogenesis in *Streptomyces*. *Appl. Microbiol. Biotechnol.* **59**:419–425.
  38. **Wang, L., and L. C. Vining.** 2003. Control of growth, secondary metabolism and sporulation in *Streptomyces venezuelae* ISP5230 by *jadW* (1), a member of the *afsA* family of gamma-butyrolactone regulatory genes. *Microbiology* **149**:1991–2004.
  39. **Wieser, J., and T. H. Adams.** 1995. *flbD* encodes a Myb-like DNA-binding protein that coordinates initiation of *Aspergillus nidulans* conidiophore development. *Genes Dev.* **9**:491–502.
  40. **Yu, J.-H., J. Wieser, and T. H. Adams.** 1996. The *Aspergillus* FlbA RGS domain protein antagonizes G protein signaling to block proliferation and allow development. *EMBO J.* **15**:5184–5190.