



# Type III effector proteins from the plant pathogen *Xanthomonas* and their role in the interaction with the host plant

Doreen Gürlebeck\*, Frank Thieme, Ulla Bonas

Institute of Genetics, Martin-Luther University Halle-Wittenberg, Weinbergweg 10, 06120 Halle, Germany

Received 15 October 2005; accepted 15 November 2005

## KEYWORDS

*Xanthomonas*;  
Type III secretion;  
Effectors;  
AvrBs3;  
AvrRxv

## Summary

Pathogenicity of *Xanthomonas campestris* pathovar (pv.) *vesicatoria* and most other Gram-negative bacterial plant pathogens largely depends on a type III secretion (TTS) system which is encoded by hypersensitive response and pathogenicity (*hrp*) genes. These genes are induced in the plant and are essential for the bacterium to be virulent in susceptible hosts and for the induction of the hypersensitive response (HR) in resistant host and non-host plants. The TTS machinery secretes proteins into the extracellular milieu and effector proteins into the plant cell cytosol. In the plant, the effectors presumably interfere with cellular processes to the benefit of the pathogen or have an avirulence activity that betrays the bacterium to the plant surveillance system. Type III effectors were identified by their avirulence activity, co-regulation with the TTS system and homology to known effectors. A number of effector proteins are members of families, e.g., the AvrBs3 family in *Xanthomonas*. AvrBs3 localizes to the nucleus of the plant cell where it modulates plant gene expression. Another family that is also present in *Xanthomonas* is the YopJ/AvrRxv family. The latter proteins appear to act as SUMO cysteine proteases in the host. Here, we will present an overview about the regulation of the TTS system and its substrates and discuss the function of the AvrRxv and AvrBs3 family members in more detail.

© 2005 Elsevier GmbH. All rights reserved.

## Introduction

Plants have to cope not only with abiotic stresses but also have to defend themselves against pathogens. Among the phytopathogens are worms, fungi,

\*Corresponding author. Tel.: +0049 345 55 26 293;  
fax: +0049 345 55 27 151.

E-mail address: [doreen.guerlebeck@genetik.uni-halle.de](mailto:doreen.guerlebeck@genetik.uni-halle.de)  
(D. Gürlebeck).

oomycetes and bacteria which all try to acquire nutrients from the host plant for their own advantage. Although plants have basal defense mechanisms, some organisms have evolved mechanisms to suppress the basal host defense and therefore are pathogenic. Our laboratory studies *Xanthomonas campestris* pathovar (pv., indicates host range) *vesicatoria* (*Xcv*),<sup>1</sup> the causal agent of bacterial spot disease, and its host plants pepper and tomato. The bacteria enter the plant tissue via stomata or wounds and multiply in the intercellular spaces of the plant tissue (Stall, 1995). Bacterial pathogenicity depends on a specialized type III secretion (TTS) system, which secretes some helper and accessory proteins into the extracellular milieu that support the injection of the effector proteins, the major substrate class of the TTS system, into the host cell (Hueck, 1998; Cornelis and Van Gijsegem, 2000).

However, some host plant genotypes are able to specifically recognize a given effector protein which thus makes the corresponding pathogen strain no longer virulent on this plant. In many cases, a localized cell death response, the hypersensitive response (HR), is induced resulting in attenuation of bacterial growth (Klement, 1982; Keen, 1990). Bacterial growth inhibition is probably due to a decrease in the apoplastic water potential during the HR (Wright and Beattie, 2004). Specific recognition follows the gene-for-gene hypothesis (Flor, 1971): plants carry resistance (*R*) genes that recognize the product of corresponding effector genes in the pathogen, which were termed avirulence (*avr*) genes. To understand the molecular basis of gene-for-gene recognition and the biological function of effector proteins, both effector (or *avr*) and *R* genes are studied intensively. In this review, we will focus on the TTS apparatus and mainly discuss the effector proteins translocated by this machinery.

## The TTS system of *Xanthomonas*

Basic pathogenicity of *Xcv* in pepper and tomato is determined by the TTS system, which is highly conserved in most Gram-negative bacterial pathogens of plants and animals (Tampakaki et al., 2004). Type III-dependent protein secretion was first identified in the animal pathogen *Yersinia enterocolitica* (Heesemann et al., 1984), however, mutants in genes coding for components of the

secretion machinery were first discovered in the plant pathogen *Pseudomonas syringae* by the analysis of non-pathogenic mutants (Niepold et al., 1985; Lindgren et al., 1986). In phytopathogenic bacteria, the TTS system is encoded by large gene clusters termed *hrp* (hypersensitive response and pathogenicity; Alfano and Collmer, 2004; Tampakaki et al., 2004; Fig. 1). *hrp* mutants no longer cause disease in susceptible plants, barely grow in plant tissue and are unable to induce the HR in resistant host and non-host plants (Bonas et al., 1991; Büttner and Bonas, 2002). The *hrp* gene clusters of plant pathogens (except for *Ralstonia solanacearum*) show typical features of pathogenicity islands, i.e., genes for virulence factors, mobile genetic elements, a deviant G+C content and codon usage as compared to the rest of the genome, and flanking tRNA genes (Fig. 1), all indicative for acquisition by horizontal gene transfer (Dobrindt et al., 2004). The regions flanking the *hrp* region vary in their gene content and often contain type III effector genes (Fig. 1; Alfano et al., 2000; Noël et al., 2002). The *hrp* gene cluster of *Xcv* is part of a 35.3 kb chromosomal genomic island (Bonas et al., 1991; Fenselau et al., 1992; Fenselau and Bonas, 1995; Rossier et al., 2000; Noël et al., 2002; Thieme et al., 2005), in which 18 out of the 37 putative coding sequences encode Hrp proteins. Eleven proteins are highly conserved in plant and animal pathogenic bacteria and were, therefore, termed Hrc (*hrp* conserved; Bogdanove et al., 1996; Tampakaki et al., 2004). Most Hrc proteins, probably forming the core of the TTS apparatus, also share homology to components of the flagellar assembly apparatus which is thought to be the evolutionary ancestor of the TTS system (Aizawa, 2001; Tampakaki et al., 2004).

The role of a few non-conserved Hrp proteins, which are essential for pathogenicity, could recently be clarified for *Xcv*. For example, HrpB2 and HrpE were found to be secreted and essential for secretion of other proteins by the TTS system (Rossier et al., 2000). HrpB2 is likely an extracellular part of the secretion machinery (Rossier et al., 2000). HrpE is the major subunit of the Hrp pilus, an extracellular appendix that is thought to be associated with the TTS apparatus (Koebnik, 2001; Weber et al., 2005). The Hrp pilus most likely serves as a conduit for secreted proteins. In contrast, the type III-secreted protein HrpF is essential for pathogenicity but dispensable for secretion in *Xcv* (Rossier et al., 2000). However, *hrpF* mutants in *X. oryzae* pv. *oryzae* (*Xoo*) were not completely abolished in virulence (Sugio et al., 2005). HrpF has been suggested to be the major component of the *Xcv* TTS translocon, a predicted

<sup>1</sup>Also designated as *Xanthomonas axonopodis* pv. *vesicatoria* (Vauterin et al., 2000) or *Xanthomonas euvesicatoria* (Jones et al., 2004).

protein channel in the plant plasma membrane that mediates the transport of effector proteins into the plant cell cytosol (Büttner et al., 2002).

The analysis of non-polar mutants in the *hrp* gene cluster of *Xcv* also identified *hpa* (*hrp* associated) genes that contribute to, but are not essential for the pathogenic interaction with the plant (Huguet et al., 1998; Noël et al., 2002; Büttner et al., 2004; U. Bonas and D. Büttner, unpublished data). For example, *hpaA* encodes a protein that is needed for the HR and full pathogenicity. HpaA is secreted by the TTS system and localizes to the plant nucleus when expressed in planta (Huguet et al., 1998). In contrast, HpaB remains in the bacterium where it plays an important role in the control of type III protein export (Büttner et al., 2004). Interestingly, the type III-secreted protein XopA (*Xanthomonas* outer protein) also shows features of an Hpa protein: The *xopA* gene is located adjacent to the left border of the *hrp* region, its expression is HrpG- and HrpX-dependent (see below). XopA shows homology to Hpa1 from *Xoo* (Zhu et al., 2000). However, XopA shows no harpin HR eliciting activity as it was shown for Hpa1 from *Xoo* and the homolog HpaG from *X. axonopodis* pv. *glycines* (Kim et al., 2003, 2004). *xopA* mutants are affected in bacterial virulence, but not in type III-mediated secretion (Fig. 1; Noël et al., 2001, 2002). As XopA is not translocated into the host cell, we speculate that it might be part of the translocon (Büttner et al., 2004).

## Regulation of *hrp* gene expression in phytopathogens

Expression of the TTS system in phytopathogenic bacteria is induced in planta or in specific minimal media mimicking the plant apoplast (Schulte and Bonas, 1992a, b; Lindgren, 1997). The regulatory cascade controlling expression of *hrp* genes varies in different plant pathogens. Based on similarities in gene organization and regulation, the *hrp* regions of different pathogens were classified into two groups. Group I comprises species of *Erwinia*, *Pantoea* and *P. syringae*, whereas *R. solanacearum* and *Xanthomonas* spp. belong to group II (Fig. 1; Alfano and Collmer, 1996). Expression of group I *hrp* genes depends on the activator HrpL, a member of the extracytoplasmic function (ECF) family of alternate sigma factors, which bind to a conserved promoter sequence termed *hrp* box (Xiao and Hutcheson, 1994; Xiao et al., 1994; Wei and Beer, 1995). *hrpL* expression is controlled by HrpR and HrpS in *P. syringae* (Xiao et al., 1994), and by HrpS,


HrpX and HrpY in *Erwinia* (Wei et al., 2000b; Mor et al., 2001; Chatterjee et al., 2002) and *Pantoea* (Frederick et al., 1993; Merighi et al., 2003). HrpS and HrpR of *Pseudomonas* are enhancer-binding proteins of the NtrC-family, synergistically activating the expression of *hrpL* in response to plant signals or in *hrp*-inducing minimal medium (Grimm and Panopoulos, 1989; Xiao et al., 1994; Grimm et al., 1995; Hutcheson et al., 2001). It was shown that HrpS and HrpR physically interact (Hutcheson et al., 2001). In *P. syringae* pv. *tomato* DC3000, *hrpL* expression is also regulated by the  $\sigma^{54}$  factor RpoN which is controlled by the GacA master regulator (Chatterjee et al., 2003). Intriguingly, under *hrp*-suppressing conditions, the Lon protease degrades HrpR and thus suppresses expression of the *hrp* regulon (Bretz et al., 2002). In contrast to HrpR, the negative regulator HrpV acts upstream of HrpR and HrpS and can downregulate *hrp* genes under inducing conditions (Preston et al., 1998). Recently, it was shown that the chaperone-like protein HrpG interacts with HrpV and serves as a suppressor of this negative regulator (Wei et al., 2005). Interestingly, *hrp* gene expression is also downregulated in *P. syringae* pv. *tomato* DC3000, when *hrpA*, which encodes the major subunit of the Hrp pilus, is mutated (Wei et al., 2000a). Although *hrp* gene expression in *Erwinia* strains also depends on HrpL, its activation differs from *Pseudomonas*. In case of *Erwinia*, HrpX is an  $\text{l}_c\text{T}$ -type sensor (Parkinson and Kofoed, 1992). In *E. amylovora*, HrpX is probably activated by phosphorylation. In turn, HrpX phosphorylates HrpY, an  $\text{RO}_{\text{III}}$  response regulator (Parkinson and Kofoed, 1992). Phosphorylated HrpY, complexed with activated HrpS, binds to the *hrpL* promoter and activates *hrpL* expression in an RpoN-dependent manner (Wei et al., 2000b). In *E. herbicola* pv. *gypsophila*, the key regulatory step is the phosphorylation of HrpY. The latter activates *hrpS* expression, and HrpS and  $\sigma^{54}$  together activate *hrpL*, whereas in this strain HrpX seems to be dispensable (Nizan-Koren et al., 2003).

Expression of group II *hrp* gene clusters is best studied in *R. solanacearum*. Here, the inducing signal is sensed by the outer membrane receptor PrhA and transduced by the putative transmembrane protein PrhR and the ECF sigma factor PrhI. Activated PrhI leads to the expression of the regulators PrhJ, HrpG and HrpB, which is homologous to HrpX from *Xcv* (Genin et al., 1992; Marena et al., 1998; Brito et al., 2002; Fig. 1A). For HrpB-dependent activation of promoters in *R. solanacearum* the *hrp*<sub>II</sub> box (TTCCG-N<sub>16</sub>-TTCCG) is required. In addition, a *cis*-acting element (YANNRT) essential for transcriptional activation,

was identified in the  $-10$  promoter region (Cunnac et al., 2004a).

The key regulatory proteins in *Xanthomonas* are HrpG and HrpX, which are encoded outside of the *hrp* gene cluster. HrpG belongs to the OmpR-family of two-component system response regulators (Wengelnik et al., 1996) and controls the expression of the regulator HrpX (Wengelnik and Bonas,

1996; Noël et al., 2001, 2002, 2003). HrpX, an AraC-type transcriptional activator, regulates the expression of most genes of the HrpG regulon including the *hrp* operons *hrpB* to *hrpF* (Wengelnik and Bonas, 1996; Noël et al., 2001, 2002, 2003). Many HrpX-regulated genes contain a PIP box (plant-inducible promoter box; consensus TTCGC-N<sub>15</sub>-TTCGC) in their promoter region, which was



proposed to serve as a regulatory element (Fenselau and Bonas, 1995; Wengelnik and Bonas, 1996). However, this motif is not sufficient to confer HrpX inducibility, because there are HrpX-independent PIP-box containing promoters, e.g., *avrRxv* (Ciesiolka et al., 1999). It appears that additionally, as described above for HrpB a –10 box is needed for HrpX inducibility (Tsuge et al., 2005). Remarkably, this –10 box is absent in the promoter of the constitutively expressed *avrRxv* mentioned above. Whether HrpX or a still unknown protein binds to the PIP box remains to be elucidated.

### Signals and recognition of type III substrates

Functional analysis of the TTS system and the identification of secreted proteins required the establishment of an in vitro secretion assay. However, in vitro growth conditions that led to the induction of the *Xcv* TTS system were not suitable for detection of type III secretion. One breakthrough was the generation of the E44K point mutation in *hrpG* (designated *hrpG\**) which resulted in constitutive expression of *hrp* genes under different growth conditions (Wengelnik et al., 1999). This mutant allowed the development of an appropriate in vitro secretion medium. Incubation of bacteria in acidic minimal medium (pH 5.2–5.4) and the addition of BSA finally led to detectable amounts of secreted proteins (Rossier et al., 1999). Interestingly, the *Xcv* TTS system also secretes heterologous proteins, e.g., PopA from *R. solanacearum*, AvrB from *P. syringae* and YopE from *Yersinia pseudotuberculosis*, indicating that the properties of the secretion signal are conserved among plant and animal pathogenic bacteria (Rossier et al., 1999). However, the nature of the

TTS signal that resides in the N-terminal regions of type III-secreted proteins is still enigmatic. In contrast to *sec* signals, TTS signals show no common physicochemical properties. Furthermore, type III-secreted proteins are not processed during secretion (Holland, 2004; Tampakaki et al., 2004). Nevertheless, recent reports describe some common features of the N-terminal amino acid sequences of effectors, e.g., the presence of hydrophilic amino acids, absence of acidic residues in the first 12 amino acid residues, amphipathicity and a bias toward serine and glutamine in the first 50 residues (Lloyd et al., 2001a, 2002; Guttman et al., 2002; Petnicki-Ocwieja et al., 2002). An alternative model, which is discussed controversially, proposes an mRNA signal in *Yersinia* outer proteins (Yops; Anderson and Schneewind, 1999; Sorg et al., 2005). However, various studies identified the signal in the first 11–17 amino acid residues of Yops (Sory et al., 1995; Schesser et al., 1996; Lloyd et al., 2001b). Similarly, the secretion signal of the effector AvrBs2 from the plant pathogen *Xcv* resides in the first 28 amino acid residues, whereas the first 58 residues are necessary for its translocation (Mudgett et al., 2000).

In addition to the secretion and translocation signal, a number of type III-secreted proteins contain a TTS-chaperone-binding site (Alfano and Collmer, 2004; Tampakaki et al., 2004). TTS chaperones are typically small, acidic and leucine-rich and share only little sequence similarity (Wattiau et al., 1994; Feldman and Cornelis, 2003; Alfano and Collmer, 2004). Usually, type III chaperones bind their substrates to stabilize them, to connect them to the TTS system and to keep them partially unfolded for their transport through the TTS pilus (Stebbins and Galán, 2001; Page and Parsot, 2002; Alfano and Collmer, 2004). TTS chaperones can be divided into classes based on their interaction with one specific effector

**Figure 1.** *hrp* regions from different plant pathogenic bacteria. (A) Schematic overview of the *hrp* regions from sequenced plant pathogens. The *hrp* clusters of *X. campestris* pv. *vesicatoria* strain 85–10 (Thieme et al., 2005), *Ralstonia solanacearum* strain GMI1000 (Salanoubat et al., 2002), *Erwinia carotovora* subsp. *atroseptica* strain SCRI1043 (Bell et al., 2004) and *Pseudomonas syringae* pv. *tomato* strain DC3000 (Buell et al., 2003) are shown. (B) Comparison of the *hrp* gene clusters of the different sequenced xanthomonads. The following strains are shown: *X. campestris* pv. *vesicatoria* strain 85–10 (Thieme et al., 2005), *X. axonopodis* pv. *citri* strain 306 (da Silva et al., 2002), *X. axonopodis* pv. *glycines* strain 8ra (Kim et al., 2003), *X. oryzae* pv. *oryzae* strain KACC10331 and *X. campestris* pv. *campestris* strain ATCC3391 (da Silva et al., 2002). Comparative analyses were carried out using the Artemis comparison tool (<http://www.sanger.ac.uk/Software/Artemis/>). Red areas indicate DNA regions with at least 80% identity over 400bp and darker color indicates higher identity. Single-headed arrows represent CDS and the direction of transcription. Red arrows refer to *hrc* genes, blue arrows to *hrp* genes, green arrows to *hpa* genes, black arrows to putative/known type III effector genes, gray arrows to genes coding for proteins of unknown function, and white arrows to genes encoding regulators of the TTS system expression. Yellow boxes indicate IS elements. Arrows above the CDS indicate the operon structure and the black dots refer PIP boxes (CDS, coding sequences; *hrp*, hypersensitive response and pathogenicity; *hrc*, *hrp* conserved; *hpa*, *hrp* associated).

(class IA) or with multiple effectors (Cornelis and Van Gijsegem, 2000; Parsot et al., 2003). In *Pseudomonas*, genes encoding chaperones of class IA are often found next to the gene of the corresponding effector (Wattiau et al., 1996). In *Xcv*, so far only HpaB shows characteristic features of TTS chaperones (Büttner et al., 2004). HpaB is encoded within the *hrp* gene cluster and interacts with more than one effector and thus appears to be a class IB TTS chaperone. Interestingly, *hpaB* mutants were strongly reduced in in vitro secretion of effector proteins, but secretion of HrpF and XopA, presumably extracellular components of the TTS system, was not affected. Furthermore, the *hpaB* mutant was able to translocate the non-effector proteins HrpF and XopA into the plant cell. These results indicate that HpaB is not only a chaperone but also plays an essential role in exit control of the TTS system, i.e., translocation of effectors and inhibition of translocation of non-effectors (Büttner et al., 2004). Indeed, it was shown before in other bacteria that TTS chaperones can regulate secretion (Boyd et al., 2000; Francis et al., 2002; Wulff-Strobel et al., 2002; Thomas and Brett Finlay, 2003).

## How to identify and verify type III effectors?

### Identification of effectors based on avirulence activity

In plant pathogens, the first type III effectors have been identified as the products of avirulence genes that are recognized by corresponding plant disease resistance genes and their products, respectively. In most cases, recognition leads to the induction of the HR concomitant with arrest of bacterial growth (Klement, 1982; Staskawicz, 2001). *Xcv* strain 85–10 expresses three known avirulence proteins, AvrBs1, AvrBs2 and AvrRxv (Ronald and Staskawicz, 1988; Whalen et al., 1993; Swords et al., 1996). AvrBs1, which is encoded on the largest plasmid, and the chromosomal, highly conserved *avrBs2* gene contribute to bacterial fitness in the field and thus to the pathogen's virulence (Kearney and Staskawicz, 1990; Gassmann et al., 2000; Wichmann and Bergelson, 2004).

Based on avirulence protein translocation, in vivo screens for effectors were developed in *Xanthomonas* and *Pseudomonas* (Guttman et al., 2002; Roden et al., 2004b; Chang et al., 2005). In *Xcv*, the N-terminally deleted (amino acids 2–152) AvrBs3


derivative (see below) AvrBs3 $\Delta$ 2 was established as a reporter. AvrBs3 $\Delta$ 2 still induces the HR when transiently expressed in resistant plants mediated by *Agrobacterium* but cannot be secreted by *Xcv* (Szurek et al., 2001). This indicates that the N-terminal region of AvrBs3 contains a secretion and translocation signal and the C-terminus the effector domain. Translational fusions of 5' coding regions of effector candidate genes to *avrBs3\Delta*2 allowed the identification and verification of type III effectors (Noël et al., 2003). If a fusion protein induces the HR in AvrBs3-responsive resistant plants, the 5' region of the candidate gene restored type III-dependent delivery of AvrBs3 $\Delta$ 2 and thus contains a functional translocation signal (Fig. 2). Using this reporter assay, we could identify a number of type III effectors, e.g., XopC and XopJ (Noël et al., 2003). Similarly, AvrBs2 lacking its TTS signal was used as a reporter for effector translocation in a Tn5-based screen (Roden et al., 2004b). This screen led to the identification of a number of new effectors (Roden et al., 2004b; Table 1, see below).

### Identification of effectors based on co-regulation with the TTS system

Additional effector genes were identified by their co-regulation with the TTS system. While in *P. syringae* all known effector genes are co-regulated with the TTS system (Xiao and Hutcheson, 1994; Chang et al., 2005), this is not true for *Xanthomonas*. For example, the effector genes *avrBs1*, *avrBsT* (Escolar et al., 2001), *avrRxv* (Ciesiolka et al., 1999) and *avrBs3* (Knoop et al., 1991) are constitutively expressed. Nevertheless, a cDNA-AFLP screen for new members of the HrpG regulon in *Xcv* identified more than 20 genes, including four genes (*xopA*, *xopB*, *xopC* and *xopJ*) that were found to encode substrates of the TTS system (Noël et al., 2001, 2002, 2003).

### Identification of effectors by in silico prediction

In the era of high-throughput sequencing of bacterial genomes, effector candidates can also be uncovered based on homology to known effectors from other pathogens. In addition, the presence of eukaryotic motifs in the predicted gene products can indicate a function inside the host cell (Büttner and Bonas, 2003; Fig. 2). Furthermore, conserved sequence motifs such as PIP boxes in the promoter regions of *Xanthomonas* genes can be used to identify genes which might be co-regulated



**Figure 2.** Strategies for the prediction and verification of type III effector proteins. (A) Schematic overview of an effector locus. Effector genes can be predicted by the presence of promoter motifs and signs of acquisition by horizontal gene transfer (red). On the amino acid level, there might be homology to known type III effectors, presence of eukaryotic features and putative TTS signals. (B) In vitro and in vivo assays to confirm type III effectors. Effector candidate fusions to the AvrBs3 $\Delta$ 2 reporter expressed in wild-type (WT) and TTS mutant (TTS<sup>-</sup>) *Xcv* strains can be visualized by immunoblotting using an AvrBs3-specific antibody. Using these strains, in vitro type III secretion can be tested in secretion medium (left panel). The right panel shows a translocation assay in which the *Xcv* strains were inoculated into an AvrBs3-responsive resistant plant. The wild-type strain used is normally virulent on this plant (see text for details).

with the TTS system (Büttner et al., 2003; see above). In *P. syringae*, prediction algorithms for effectors have been developed based on a characteristic N-terminal amino acid composition (see above), which presumably serves as export signal (Guttman et al., 2002; Petnicki-Ocwieja et al., 2002). A number of effector genes show signs of horizontal gene transfer, e.g., low G+C content of the DNA, compared to the average of the genome, and flanking mobile elements or tRNA genes (Büttner et al., 2003; Noël et al., 2003; Thieme et al., 2005; Fig. 2 and Table 1). Based on the genome sequence, *Xcv* strain 85–10 contains 14 known effector genes (Table 1). Remarkably, *xopC* is the only specific effector gene unique for *Xcv*, because there is no homolog in the database and similarity to *R. solanacearum* proteins is only partial (Table 1). Interestingly, the *xopC* region shows signs of horizontal gene transfer and is associated with a putative effector integron consisting of genes encoding a transposase and coin-tegrate resolution proteins, flanked by inverted repeats (Noël et al., 2003). Similar integron elements are also present in *X. axonopodis* pv. *citri* (*Xac*). These elements are always associated with effector genes or candidates and could be involved in the horizontal transfer of virulence factors (Noël et al., 2003). In contrast to the unique XopC

protein, XopQ homologs are present in different pathogens. The effector belongs to the HopQ1-1 (formerly HopPtoQ) family (Rohmer et al., 2004) and is a putative inosine-uridine nucleoside *N*-ribohydrolase (Roden et al., 2004b; Table 1). Unique effectors like XopC could be important for a specific host interaction, whereas the wide distribution of an effector gene in different bacteria like XopQ suggests a more general role in virulence.

### What do we know about the function of type III effectors?

The first type III effector from plant pathogens was isolated as an avirulence protein from *P. syringae* pv. *phaseolicola* in 1984 (*avrA*; Staskawicz et al., 1984). Since then, more than 40 Avr proteins and many type III effectors, respectively, were identified in plant pathogens possessing a TTS system, e.g., *Ralstonia*, *Xanthomonas*, *Pantoea* and *Erwinia*. Although the first type III effectors were identified because of their activity in restricting the host range of the pathogen by inducing the HR (avirulence activity), it was assumed that there has to be a function of TTSS effector proteins for the producer's advantage

Table 1. Known type III-effector proteins of Xcv strain 85-10

Gene name	Predicted function/ homology*	HrpG-dep.†	G+C‡ (%)	Homologs§ in					Reference	
				Xac	Xcc	Xoo	Pst	Rs		Eca
<i>avrBs1</i>	Unknown function	–	42.23	–	+	–	–	–	–	Ronald and Staskawicz (1988) and Escobar et al. (2001)
<i>avrBs2</i>	Putative glycerophosphoryl-diester phosphodiesterase	nd	63.59	+	+	+	–	–	–	Swords et al. (1996)
<i>avrRxv</i>	YopJ/AvrRxv-family, putative cysteine protease	–	52.32	–	(+)	–	–	–	–	Whalen et al. (1993) and Ciesiolka et al. (1999)
<i>xopB</i>	Homology to HopD1 ( <i>P. syringae</i> pv. <i>tomato</i> )	+	55.54	–	–	–	+	–	–	Noël et al. (2001)
<i>xopC</i>	Unknown function	+	47.50	–	–	–	–	–	–	Noël et al. (2003)
<i>xopD</i>	SUMO cysteine protease; C48 family	+	54.76	–	+	–	–	(+)	–	Noël et al. (2002) and Hotson et al. (2003)
<i>xopF1</i>	Unknown function	nd	65.47	(+)	(+)	+	–	–	–	Roden et al. (2004b)
<i>xopF2</i>	Unknown function	nd	64.72	(+)	(+)	+	–	–	–	Roden et al. (2004b)
<i>xopJ</i>	AvrRxv/YopJ-family, putative cysteine protease	+	56.86	–	(+)	–	–	+	–	Roden et al. (2004b)
<i>xopN</i>	Unknown function	nd	63.44	+	+	+	–	–	–	Roden et al. (2004b)
<i>xopO</i>	Homology to HopK1 and AvrRps4 ( <i>P. syringae</i> )	nd	52.04	–	–	–	+	–	–	Roden et al. (2004b)
<i>xopP</i>	Unknown function	nd	61.66	+	+	+	–	+	–	Roden et al. (2004b)
<i>xopQ</i>	HopQ1-1 family protein, putative inosine-uridine nucleoside N-ribohydrolase	nd	68.88	+	+	+	+	+	–	Roden et al. (2004b)
<i>xopX</i>	Unknown function	nd	65.95	+	+	+	–	–	–	Metz et al. (2005)
<i>ecf</i>	Early chlorosis factor, homology to HopAE1 ( <i>P. syringae</i> pv. <i>syringae</i> )	–	64.88	–	–	+	–	–	–	Morales et al. (2005)

\* Predicted function and homology to known type III-effector proteins from *Pseudomonas syringae* or other *Xanthomonas* spp. For *Pseudomonas* effectors the unified nomenclature was used (Lindeberg et al., 2005).

† HrpG- and HrpX-dependent co-regulation with the TTS system (nd, not determined).

‡ G+C content of the DNA in the coding region.

§ Homologs were determined using BLAST algorithms. (+) indicates partial homology or a disrupted homolog in the corresponding genome. (Xac: *X. axonopodis* pv. *citri*; Xcc: *X. campestris* pv. *campestris*; Xoo: *X. oryzae* pv. *oryzae*; Pst: *P. syringae* pv. *tomato*; Rs: *R. solanacearum*; Eca: *E. carotovora* subsp. *atroseptica*).




(virulence activity). A contribution to virulence could indeed be observed for several Avr proteins and effectors, respectively (Kearney and Staskawicz, 1990; Swarup et al., 1991, 1992; Lorang et al., 1994; Yang et al., 1994, 2000; Ritter and Dangl, 1995; Bogdanove et al., 1998; Huguet et al., 1998; Jackson et al., 1999; Alfano et al., 2000; Chang et al., 2000; Chen et al., 2000; Gassmann et al., 2000; Shan et al., 2000; Tsiamis et al., 2000; Vera Cruz et al., 2000; Guttman and Greenberg, 2001; Badel et al., 2002; Guo et al., 2002; Lavie et al., 2002; Abramovitch et al., 2003; Deng et al., 2003; Holeva et al., 2004; Lopez-Solanilla et al., 2004; Losada et al., 2004; Yang and White, 2004; Kay et al., 2005; Lim and Kunkel, 2005; Lin and Martin, 2005; Metz et al., 2005). However, in many cases, a virulence function of type III effectors is difficult to assess, because the pathogens have several effectors which can be functionally redundant and often show no homologies to proteins with known function. It is interesting that effector proteins are quite diverse whereas the analysis of their counterparts, the known resistance gene products, revealed a large number of related proteins in different plant species (Dangl and Jones, 2001; Lahaye, 2002; Martin et al., 2003). This suggests that plants use similar mechanisms to detect a wide range of diverse effector proteins to defend themselves and avoid pathogen's growth. However, although effector proteins generally vary in their amino acid sequence, some of them are homologous and were therefore grouped into families. Does this mean that pathogens also pursue common mechanisms to manipulate the host, to defend themselves against plant's defense and to grow? Here, we will focus our discussion on two major families of type III effectors from *Xanthomonas* characterized by remarkable features (Fig. 3), which indicated a function for the proteins in virulence.

## A family of cysteine proteases


A growing number of type III effectors appear to act as cysteine proteases in plants. Based on primary sequence analysis, they were classified into four groups: the AvrRxx/YopJ (C55) family (Orth et al., 2000), the XopD (C48) family (Hotson et al., 2003), the HopAR1/YopT (C58) family (Shao et al., 2002) and the AvrRpt2 (C70) family (Mudgett and Staskawicz, 1999). Members of the AvrRxx/YopJ family have been identified in mammalian and plant pathogens suggesting that they play an important role in the interaction with the host. Examples include YopJ and YopP from *Y. pseudotuberculosis* and *Y. enterocolitica*, respectively,

AvrRxx, AvrXv4, AvrBsT and XopJ from *Xcv*; PopP1 and PopP2 from *R. solanacearum*; ORFB from *Erwinia*, HopZ1 (formerly known as HopPmaD), HopZ2 (formerly known as AvrPpiG1) and HopZ3 (formerly known as HopPsyV) from *P. syringae* (Whalen et al., 1993; Galyov et al., 1994; Ciesiolka et al., 1999; Alfano et al., 2000; Astua-Monge et al., 2000; Arnold et al., 2001; Noël et al., 2001, 2003; Lavie et al., 2002; Deng et al., 2003; Deslandes et al., 2003). Comparison of predicted secondary structures of these effectors revealed similarities to the cysteine protease from adenovirus (AVP; Orth et al., 2000; Orth, 2002), which shows similarity to yeast Ulp1, a SUMO protease of the C48 family (Orth, 2002). All members of this family contain a conserved catalytic triad necessary for the HR induction (Fig. 3A; Orth et al., 2000; Shao et al., 2002; Hotson et al., 2003; Roden et al., 2004a). The substrate of YopJ appears to be the small ubiquitin-like modifier protein SUMO-1. SUMO is post-translationally ligated to lysin residues of target proteins (Hochstrasser, 2000a,b) which, e.g., blocks ubiquitination and subsequent proteolysis and thus stabilizes the target proteins. Removal of SUMO from host proteins by a SUMO protease would allow binding of ubiquitin leading to degradation of the protein. Neither YopJ nor AvrXv4 showed protease activity in vitro (Orth et al., 2000; Roden et al., 2004a). This is in good agreement with the results for AVP, which requires an activator for catalytic activity (Orth, 2002). Overexpression of YopJ resulted in a decrease of SUMO bound to proteins (Orth et al., 2000, 2002). Furthermore, similar results were shown for AvrXv4 transiently expressed in planta (Roden et al., 2004a). For XopD, the ability to remove SUMO from plant proteins in plant extracts was demonstrated (Hotson et al., 2003). XopD, isolated from *Xcv* based on its position in the left *hrp* flanking region and homology to PsvA from *P. syringae* pv. *eriobotryae* (Noël et al., 2002), belongs to the C48 cysteine proteases, whereas YopJ, AvrBsT, AvrRxx, AvrXv4 and XopJ are C55 cysteine proteases (Fig. 3A, for review see Hotson and Mudgett, 2004). Interestingly, proteins of both cysteine protease families share another characteristic: XopD, PopP2, AvrRxx, AvrBsT and AvrXv4 contain putative nuclear localization signals (NLSs; Ciesiolka et al., 1999; Noël et al., 2002; Deslandes et al., 2003; Roden et al., 2004a). Indeed, PopP2 from *R. solanacearum* and XopD from *Xcv* localize to the plant cell nucleus where the latter accumulates in subnuclear foci (Deslandes et al., 2003; Hotson et al., 2003; L. Noël and U. Bonas, unpublished). In contrast, AvrXv4 and AvrRxx localize in the plant cell cytoplasm (Roden et al., 2004a; Bonshtien et al., 2005) suggesting

### A XopD, a C48 cysteine protease from *Xanthomonas*



### B AvrBs3 from *Xanthomonas*



**Figure 3.** Structural features of representatives of two type III effector families from *Xanthomonas*. (A) XopD as an example for a C48 cysteine protease. The characteristic features of cysteine protease effectors are presented: The reactive centre containing the catalytic triad (black box) is essential for the function of the family members and is shown for different members of the C48 and C55 cysteine protease families from *Xanthomonas* (XopD, AvrRxv, AvrXv4, AvrBsT, XopJ), *Yersinia* (YopJ), yeast (Ulp1) and adenovirus (AVP). The residues forming the catalytic triad are shown in bold. The nuclear localization signal is indicated (black bar). (B) Structural features of AvrBs3 from *Xcv*. The central repeat region consists of 17.5 nearly identical tandem repeats of 34 amino acids. The C-terminal region contains different eukaryotic motifs: imperfect heptad leucine zipper repeats (black box; Gabriel, 1999); two functional nuclear localization signals (NLS; black bars; Van den Ackerveken et al., 1996; Szurek et al., 2001); and an acidic activation domain (AAD; Szurek et al., 2001). The secretion signal and the HpaB binding site are indicated (gray hatched and gray, respectively; Szurek et al., 2002; Büttner et al., 2004).

that the plant target proteins of these structurally related effectors differ. For example, XopD might manipulate the transcriptome by deSUMOylating transcriptional regulators, which then will be degraded. The fact that a pathogen delivers a set of proteins with similar functions into the plant cell where they localize to different compartments is probably a very efficient way to attack a wide range of plant proteins by mimicking different plant SUMO proteases.

One major question concerns the proteins targeted by type III effectors. Using YopJ as a bait in a yeast two-hybrid approach, several MAPK kinases were isolated (Orth et al., 1999). A second example is PopP2 from *R. solanacearum*, which was shown to

interact with the corresponding R protein RRS1-R in *Arabidopsis* (Deslandes et al., 2003). Since RRS1-R recognizes PopP2, it is unlikely that RRS1-R is the virulence target of PopP2 (for review see; Alfano and Collmer, 2004). HopAR1 (formerly known as AvrPphB) from *P. syringae* coimmunoprecipitates the PBS1 kinase from *Arabidopsis*, indicating a physical interaction of both proteins (Shao et al., 2003). AvrRpt2, another cysteine protease from *P. syringae*, cleaves RIN4, an interactor of the matching R protein, RPS2. A physical interaction between AvrRpt2 and RIN4 is likely, but involves a plant protein, recently identified as cyclophilin (Coaker et al., 2005; Kim et al., 2005b). The isolation of additional target proteins including targets of the *Xcv* SUMO-proteases will shed

more light on the way of action of these families of type III effectors.

## A family of transcriptional activators

The effectors discussed above most likely interfere with host cellular processes by their enzymatic function as proteases. In contrast, effector proteins of a second large family, AvrBs3 family members presumably manipulate the host cell transcriptome more directly. Interestingly, members of the AvrBs3 family were only found in *Xanthomonas* sp. with the exception of Brg11 from *R. solanacearum* (Cunnac et al., 2004b), which is, however, more distantly related. Members of the *Xanthomonas* AvrBs3 family share 90–97% amino acid sequence identity and are characterized by striking features (Fig. 3B): all members contain a repetitive region in the central part of the polypeptide, imperfect heptad leucine zipper (LZ) repeats, NLSs and an acidic activation domain (AAD) (Herbers et al., 1992; Yang et al., 1994, 2000; Yang and Gabriel, 1995a,b; Van den Ackerveken et al., 1996; Zhu et al., 1998; Gabriel, 1999; Lahaye and Bonas, 2001; Szurek et al., 2001; van 't Slot and Knogge, 2002; Yang and White, 2004). Genetic analyses of AvrBs3-like effectors revealed that in most cases the known virulence and avirulence activities in the plant cell depend on the NLSs and the AAD (see below). The repeat region consists of 5.5 (Avrxa5) to 28.5 (PthXo3) nearly identical direct repeats of 34 amino acid residues and confers the avirulence specificities of these effectors (accession AAQ79773; Herbers et al., 1992; Yang et al., 1994, 2000; Yang and Gabriel, 1995b; Zhu et al., 1998; Lahaye and Bonas, 2001; Yang and White, 2004). The presence of nearly identical tandem repeats in all family members suggests that this family is naturally growing by recombination events, which indeed could be shown (Yang and Gabriel, 1995a; Yang and White, 2004). Interestingly, Hax2 from *X. campestris* pv. *armoraciae* (*Xca*; Kay et al., 2005) and Brg11 from *R. solanacearum* are distinct from all other family members, because their repeats consist of 35 amino acid residues. Strikingly, both genes were isolated from bacteria which are pathogenic on *Arabidopsis* (Deslandes et al., 1998; Kay et al., 2005) suggesting a correlation between repeat structure and host range of the bacteria.

AvrBs3, the founder of this family, was isolated from *Xcv* based on its avirulence activity on resistant pepper plants (Bonas et al., 1989). However, it could be shown for an increasing number of family members that their primary


function is in virulence. Diseases caused by *Xanthomonas* spp., the observed symptoms on plants and specific plant reactions are shown in Fig. 4. For instance, AvrBs3 specifically induces a hypertrophy in susceptible pepper plants and other solanaceous species (Fig. 4A4). The hypertrophy, which is the enlargement of single mesophyll cells, might play a role in bacterial dispersal (Marois et al., 2002). In field studies, it has indeed been shown that AvrBs3 promotes bacterial spreading (Wichmann and Bergelson, 2004). Also PthA, the first effector that was isolated based on its virulence activity from the citrus canker pathogen *Xac*, causes pustles visible on the surface of leaves, stems and fruits (Swarup et al., 1991). This hyperplasia caused by PthA is, in contrast to the hypertrophy, due to multiplication of plant cells (compare Figs. 4A4 and 4B4). Apl1 from *Xac* also induces citrus canker-like symptoms (Ishihara et al., 2003). The recently isolated family members *hax2* to *hax 4* from *Xca* contribute additively to the aggressiveness of the pathogen on *Raphanus sativus* (Kay et al., 2005). Also for other members of the AvrBs3 family a virulence function could be observed: Avr6 from *X. campestris* pv. *malvacearum* increases the release of bacteria to the surface of infected cotton plants (Yang et al., 1994). The rice pathogen *Xoo* encodes several AvrBs3 homologs that contribute to bacterial blight disease (Fig. 4C), e.g., AvrXa7 and, to a lesser extent, Avrxa5 enhance bacterial growth and lesion formation in rice (Bai et al., 2000). Bai et al. (2000), Yang et al. (2000) and Yang and White (2004) described even more family members from *Xoo* contributing to virulence (*aB* and *pthXo* genes). It should be emphasized that studies of *Xoo* strains revealed a large number of AvrBs3-like effectors compared to *Xcv* strains, which bear 1, 2 or no homologs of AvrBs3 (Lee et al., 2005; Ochiai et al., 2005; Thieme et al., 2005). It would be interesting to understand why the rice pathogen has evolved so many *avrBs3* family members to conquer its host. The central questions are how AvrBs3 and its homologs induce the observed plant reactions and which host proteins or genes are targeted.

## Following a type III effector from the pathogenic bacterium into the plant cell


### The destination: visualization of type III effectors at their site of action

Although AvrBs3 has been studied for quite some time its biochemical function is not entirely clear yet. The presence of NLSs and an AAD in the


**A:** Bacterial spot disease caused by *Xanthomonas campestris* pv. *vesicatoria*



**B:** Citrus canker caused by *Xanthomonas axonopodis* pv. *citri*



**C:** Bacterial blight disease caused by *Xanthomonas oryzae* pv. *oryzae*



**Figure 4.** Plant diseases caused by different *Xanthomonas* species. Xanthomonads are phytopathogenic bacteria which cause various disease symptoms on important crop plants: (A) *Xanthomonas campestris* pv. *vesicatoria* causes bacterial spot disease on pepper and tomato. Water-soaked lesions are induced on leaves and fruits which later become necrotic (A1 and A2). AvrBs3 of *Xcv* induces hypertrophy in susceptible plant leaves visible as pustules on the lower epidermis (A3). Light microscopy analysis of a pepper leaf section infected with *Xcv* strain I74A expressing *avrBs3* visualizes the hypertrophy, the enlargement of single mesophyll cells (A4). This photograph is reprinted with permission from Marois et al. (2002). (B) *Xanthomonas axonopodis* pv. *citri* is the causal agent of citrus canker. The symptoms are raised circular lesions, which later becoming pustules which then darken and thicken into brown corky canker visible on leaves (B1), fruits (B2) and stems (B3). Photographs are reprinted with permission from Dean W. Gabriel: <http://www.biotech.ufl.edu/PlantContainment/canker.htm>. The pustule formation is due to tissue hyperplasia, an excessive mitotic cell division (B4). This photograph is reprinted with permission from Swarup et al. (1991). (C) *Xanthomonas oryzae* pv. *oryzae* causes bacterial blight disease in rice. Typical symptoms are water-soaked to yellowish stripes on leaf blades which start at leaf tips and increase in length extending to the sheath (C1). The photograph of rice was provided by Dr. Jan Leach, Bioagricultural Sciences and Pest Management Department at Colorado State University. (C2) Infections of IR24 rice leaves with *Xoo* PXO99. On the left, the wild-type strain was inoculated. The middle leaf is inoculated with a PXO99ME2 mutant missing *pthXo1* and the right leaf with PXO99ME2 with *pthXo1* on a plasmid. Disease symptoms are stronger if *pthXo1* is present. The leaves were photographed 3 d after inoculation. The photographs were provided by Dr. Bing Yang and Dr. Frank White, Department of Plant Pathology, Kansas State University.

C-terminus suggested that AvrBs3 is transported into the plant cell and localizes to the nucleus (Van den Ackerveken et al., 1996; Szurek et al., 2001) at that time providing a first indication that bacterial effectors act inside the host plant cell. Using *Agrobacterium*-mediated transient expression, AvrBs3 was expressed inside the cells of resistant pepper plants and induced a specific HR (Van den Ackerveken et al., 1996). This result strongly suggested that AvrBs3 is recognized inside the plant cell upon delivery by *Xcv*. Since then, the transient expression assays were crucial to demonstrate the activity of a growing number of effectors inside the plant cell.

While the injection of type III effectors from *Yersinia* into the host cell cytosol was first shown in 1994 (YopE; Sory and Cornelis, 1994), a direct proof of translocation by phytopathogenic bacteria was demonstrated only much later. Two approaches using biochemistry and immunocytochemistry were developed to demonstrate type III-dependent translocation into the plant cell. For the biochemical approach (Casper-Lindley et al., 2002) the effector AvrBs2 was fused to the calmodulin-dependent adenylate cyclase domain (Cya) of *Bordetella pertussis* cyclolysin, which is only active in eukaryotic cells, but not in bacteria. The detection of Cya activity in the plant cell strongly indicated that AvrBs2 is delivered into the plant cell (Casper-Lindley et al., 2002). The "Cya-assay" has been used to show the translocation of a number of effectors into the plant cell (Roden et al., 2004b; Schechter et al., 2004). Also for an AvrBs3::Cya fusion protein, a Cya activity in the plant was measured after delivery by *Xcv* (S. Köthke, F. Thieme and U. Bonas, unpublished). Using immunocytochemistry, a more direct proof for the translocation of a type III effector was possible and allowed us to directly visualize the effectors AvrBs3 and AvrBs4 in the plant cell. AvrBs3-specific antibodies detected both proteins in plant nuclei of susceptible and resistant pepper plants after infection with *Xanthomonas* (Szurek et al., 2002; B. Szurek and U. Bonas, unpublished). Furthermore, these experiments clearly demonstrated that the N-terminal amino acids 1–27 of AvrBs3 are essential for the type III-dependent translocation of AvrBs3 by *Xcv* into the plant cell and that localization to the nucleus depends on the NLSs (Szurek et al., 2002). Recently, the bacterial protein HpaB was identified, which is probably necessary for the translocation of AvrBs3 at the right moment. HpaB shows typical features of type III chaperones and physically interacts with the first 50 amino acids of AvrBs3 (see above; Büttner et al., 2004).

### Upon arrival: AvrBs3 dimerizes in the plant cell cytoplasm

As mentioned above, AvrBs3 family members possess a putative LZ and an AAD suggesting a role in transcriptional activation in the eukaryotic host cell. Transcription factors are often post-translationally modified and form oligomers. Indeed, AvrBs3 dimerizes in the cytoplasm prior to nuclear import (Gürlebeck et al., 2005). Surprisingly, the dimerization is not conferred by the LZ-like motif, which is known to be involved in homo- and heterodimerization of plant bZip factors (Landschulz et al., 1988; Meshi and Iwabuchi, 1995), but by the repeat region in AvrBs3. Interaction studies in yeast revealed that a certain number, but not defined repeats of AvrBs3 are necessary for dimerization indicating that the leucine-rich backbone of the central region confers the dimerization rather than the variable residues of the repeats (positions 12 and 13; Gürlebeck et al., 2005). Although the heptad leucine repeats are not essential for homodimerization of AvrBs3, they might be involved in heterodimerization, e.g., with a plant bZip factor. Interestingly, it was shown that the region of Apl1 containing the LZ motif can contribute to recognition specificity in addition to the central repeat region (Ishihara et al., 2004). Although the C-termini of AvrBs3 and AvrBs4 containing the LZ are almost identical and interchangeable, they differ in their recognition specificity (Bonas et al., 1989, 1993; Ballvora et al., 2001). However, the avirulence activities of effector proteins are determined rather by the host than by the pathogen and the LZ motifs of the effectors plays probably a major role in virulence. Interestingly, ATR13, an effector recently cloned from the downy mildew *Hyaloperonospora* (formerly *Pero-nospora*) and recognized by the resistance protein RPP13 from *Arabidopsis* also contains a heptad repeat region, next to four direct repeats (Allen et al., 2004). Although ATR13 is clearly distinct from the AvrBs3 family, the presence of a putative LZ and direct repeats in an oomycete pathogen suggests a more widespread usage and importance of these structural motifs for phytopathogens. The interesting question remains if these motifs indeed confer interaction with plant proteins.

### Searching for proteins targeted by AvrBs3

Potential interaction partners of AvrBs3 were identified by yeast two-hybrid based analysis of pepper and tomato cDNA libraries (Szurek et al., 2001; B. Szurek, D. Gürlebeck, J. Stuttmann and U.


Bonas, unpublished). One of the proteins interacting with AvrBs3 is importin  $\alpha$ , which specifically interacts with the NLSs of AvrBs3 in yeast and in vitro (Szurek et al., 2001; Schornack et al., 2004). Importin  $\alpha$  is the only interactor of an AvrBs3 family member reported so far. Since the NLSs of AvrBs3 family members are conserved, importin  $\alpha$  is supposed to interact with all AvrBs3-like effectors. In agreement with this hypothesis, an interaction between importin  $\alpha$  and AvrBs4 was demonstrated (Szurek et al., 2001; Schornack et al., 2004). Importin  $\alpha$  is part of the nuclear import machinery in eukaryotes and binds to the NLSs of a target protein via its armadillo repeats. Then, importin  $\beta$  interacts with importin  $\alpha$  and mediates the import of the protein complex into the nucleus (Görlich et al., 1994, 1995; Mattaj and Englmeier, 1998; Komeili and O'Shea, 2000). AvrBs3 localizes in both susceptible and resistant plants to the nucleus but induces different plant reactions (hypertrophy or the HR, respectively). Importin  $\alpha$  is most likely expressed in resistant and susceptible plants and therefore it is presumably not the product of the so far unknown resistance gene *Bs3* which recognizes AvrBs3. Indeed, the most simple model to explain gene-for-gene interactions was the receptor-ligand model in which the products of the plant *R* gene and the *avr* gene of the pathogen physically interact with each other (Gabriel and Rolfe, 1990). However, such direct interactions were only observed for AvrPto from *P. syringae* and Pto from tomato, AvrPita from *Magnaporthe grisea* and Pi-ta from rice and for PopP2 from *R. solanacearum* and RRS1-R from *Arabidopsis* (Scofield et al., 1996; Tang et al., 1996; Jia et al., 2000; Deslandes et al., 2003). Interestingly, not only the kinase Pto but also Prf, a typical NB-LRR resistance protein was found to be necessary for the recognition of AvrPto in tomato. This is consistent with the "guard hypothesis" (Van der Biezen and Jones, 1998; also see Schornack et al., this issue), which proposes the interaction of the effector with a plant protein, its virulence target. In resistant plants, the *R* protein monitors the modification of this virulence target which leads to the induction of defense reactions. With regard to the guard model, Pto could be a virulence target of AvrPto, guarded by Prf. However, yeast three-hybrid screens using Pto and AvrPto as baits did not isolate Prf (Bogdanove and Martin, 2000). Also for other type III effectors, guarded virulence targets in *Arabidopsis* were reported: PBS1 interacts with the *Pseudomonas* effector HopAR1, and RIN4 is targeted by AvrRpt2, AvrRpm1 and AvrB1 from *Pseudomonas* (Mackey et al., 2002; Shao et al., 2003; Kim et al., 2005a). While the known guarding *R* protein for PBS1 is

RPS5, RIN4 is protected by at least two *R* proteins: RPM1 and RPS2 (Mackey et al., 2002, 2003; Axtell and Staskawicz, 2003; Mackey, 2003).

Could the AvrBs3-interactor importin  $\alpha$  also be a virulence target that is guarded by different *R* proteins? Since importin  $\alpha$  is necessary for a large number of plant proteins to be imported in the nucleus, it presents an attractive target for bacterial effectors to modify plant reactions. Therefore, importin  $\alpha$  should be well guarded by the plant. Indeed, the corresponding resistance gene of AvrBs3, *Bs3*, only confers recognition of nuclear localized AvrBs3. However, deletion derivatives of AvrBs3 which are still nuclear imported but lack the AAD or the repeat region do not induce the HR in *Bs3* pepper plants. Thus, *Bs3* cannot be the guardian of importin  $\alpha$ . *Bs4*, a typical NB-LRR *R* protein recognizes not only wild-type AvrBs4 (97% sequence identity with AvrBs3) but also derivatives which cannot be imported into the nucleus (e.g., AvrBs4 $\Delta$ NLS; S. Schornack, T. Lahaye and U. Bonas, unpublished data; Ballvora et al., 2001; Schornack et al., 2004). Thus, *Bs4* is active even if importin  $\alpha$  is not targeted by AvrBs4. Based on these data it is likely that importin  $\alpha$  is recruited by the AvrBs3 family members for nuclear localization and not a virulence target guarded by *R* proteins. Besides *Bs4*, several *R* genes that correspond to *avrBs3*-like genes in the pathogen have been isolated. Interestingly, although they recognize very homologous type III effectors, the *R* genes identified so far are quite diverse. For example, *xa5* from rice is a recessive allele which encodes a homolog of the  $\gamma$ -subunit of transcription factor IIA and confers recognition of Avrxa5 from *Xoo* (Blair et al., 2003; Iyer and McCouch, 2004). The recently isolated *Xa27* gene from rice is induced upon infection with *Xoo* strains expressing the AvrBs3 homolog AvrXa27. Strikingly, the only differences between the resistant and susceptible alleles lie within the promoter region (Gu et al., 2004, 2005).

### Searching for genes targeted by AvrBs3

AvrBs3 family members were postulated to act as transcriptional activators. As mentioned above, AvrXa27 induces the expression of the corresponding resistance gene *Xa27*. Our group has investigated the AvrBs3-dependent transcript profile of susceptible pepper plants after *Xcv* infection. More than 20 genes upregulated by AvrBs3 (*upa*) were identified (Marois et al., 2002; Kay and Bonas, unpublished). The homologies of the predicted *upa* gene products to a pectate lyase,  $\alpha$ -expansins and auxin-induced proteins (Marois et al., 2002) suggest



**Figure 5.** Model for the activities of the *Xanthomonas* type III effectors XopD and AvrBs3. (A) Proposed way of action of XopD, a member of the C48 family of cysteine proteases. After secretion and translocation of XopD into the plant cell (2), it probably binds to importin  $\alpha$  via its nuclear localization signal (NLS, 3). Importin  $\beta$  binds to the complex (4) and mediates the nuclear import of XopD (5). Inside the nucleus, XopD localizes to nuclear foci, where it deSUMOylates yet unidentified plant proteins (6; Hotson et al., 2003). (B) Proposed way of action of AvrBs3. AvrBs3 is constitutively expressed in *Xcv* (Bonas et al., 1989). The TTS chaperone HpaB physically interacts with the N-terminal 1–50 amino acids of AvrBs3 and promotes its secretion through the TTS system (Büttner et al., 2004). AvrBs3 is translocated into the plant cell (Szurek et al., 2002), where it dimerizes (Gürlebeck et al., 2005). Importin  $\alpha$  specifically interacts with the NLSs of AvrBs3 (Szurek et al., 2001), importin  $\beta$  probably binds to this complex and mediates transport into the nucleus (6). Here, AvrBs3 either binds directly to DNA (7a) or in a complex with other plant proteins (7b) and modifies the host cell transcriptome resulting in hypertrophy symptoms (Marois et al., 2002) or the HR.

a role in modification of the plant cell wall architecture which is in accordance with the hypertrophy triggered by AvrBs3. Of special interest are *upa* genes that are induced independently of de novo protein synthesis and thus could be direct targets of AvrBs3 (Marois, Kay and Bonas, unpublished). Several *upa* genes encode proteins with homology to enzymes and to transcriptional regulators. The latter are of special interest because they could be primary targets of AvrBs3 at the top of the cascade leading to disease. Detailed analysis of

the *upa* genes is in progress and will give important insights into the mechanism by which AvrBs3-like proteins trigger host gene expression. Interestingly, for AvrXa7 from *Xoo* binding to AT-rich DNA was shown (Yang et al., 2000). However, no specific DNA sequence was defined. Although a known type of DNA-binding motif is missing in AvrBs3-like proteins, we cannot exclude that AvrBs3 and its homologs bind directly to promoter elements. Alternatively, AvrBs3 might interact with a transcription factor that itself can bind to DNA and whose activity is

modulated by AvrBs3 via its AAD (Fig. 5). Strikingly, in recent yeast two-hybrid screens, cDNAs were isolated from tomato, which indeed encode putative transcription factors (Gürlebeck, Raschke and Bonas, unpublished).

## Future prospects

Due to the growing amount of bacterial genome data and the possibility of in silico prediction and high-throughput analyses the number of effector proteins from phytopathogens increases constantly. The major challenge for the future is to assign molecular functions to individual effector proteins. The contribution of putative virulence factors to the host–pathogen interaction will be assayed by genetic and phenotypic analyses. The generation of multiple knockout mutants might be required to eliminate functional redundancy and to reveal the contribution of type III effectors to disease development. Identification of plant virulence targets by protein interactor screens using yeast two-hybrid studies and biochemical approaches might help to get a deeper insight into the molecular function of a given effector protein. The comparison of identified virulence targets, especially for large effector families, could reveal common strategies of phytopathogens to conquer their hosts. And, finally, the identification and analysis of corresponding R proteins will illustrate which diverse or common mechanisms plants use to defend themselves.

## Acknowledgments

Our work discussed here was funded by the Deutsche Forschungsgemeinschaft (SFB 363). We are grateful to the “AvrBs3 family members” Dean Gabriel, Jan Leach, Bing Yang, Frank White, Mr. Takashi Fujikawa and Shinji Tsuyumu for their cooperation and help to create Fig. 4 by providing photographs and permissions to use published photographs. We thank S. Kay and O. Kirchner for critically reading the manuscript.

## References

- Abramovitch RB, Kim Y-J, Chen S, Dickman MB, Martin GB. *Pseudomonas* type III effector AvrPtoB induces plant disease susceptibility by inhibition of host programmed cell death. *EMBO J* 2003;22:60–9.
- Aizawa SI. Bacterial flagella and type III secretion systems. *FEMS Microbiol Lett* 2001;202:157–64.
- Alfano JR, Collmer A. Bacterial pathogens in plants: life up against the wall. *Plant Cell* 1996;8:1683–98.
- Alfano JR, Collmer A. Type III secretion system effector proteins: double agents in bacterial disease and plant defense. *Annu Rev Phytopathol* 2004;42:385–414.
- Alfano JR, Charkowski AO, Deng WL, Badel JL, Petnicki-Ocwieja T, van Dijk K, et al. The *Pseudomonas syringae* Hrp pathogenicity island has a tripartite mosaic structure composed of a cluster of type III secretion genes bounded by exchangeable effector and conserved effector loci that contribute to parasitic fitness and pathogenicity in plants. *Proc Natl Acad Sci USA* 2000;97:4851–61.
- Allen RL, Bittner-Eddy PD, Grenville-Briggs LJ, Meitz JC, Rehmany AP, Rose LE, et al. Host–parasite coevolutionary conflict between *Arabidopsis* and downy mildew. *Science* 2004;306:1957–60.
- Anderson DM, Schneewind O. *Yersinia enterocolitica* type III secretion: an mRNA signal that couples translation and secretion of YopQ. *Mol Microbiol* 1999;31:1139–48.
- Arnold DL, Gibbon MJ, Jackson RW, Wood JR, Brown J, Mansfield JW, et al. Molecular characterization of *avrPphD*, a widely distributed gene from *Pseudomonas syringae* pv. *phaseolicola* involved in non-host recognition by pea (*Pisum sativum*). *Physiol Mol Plant Pathol* 2001;58:55–62.
- Astua-Monge G, Minsavage GV, Stall RE, Vallejos CE, Davis MJ, Jones JB. *Xv4-avrXv4*: a new gene-for-gene interaction identified between *Xanthomonas campestris* pv. *vesicatoria* race T3 and the wild tomato relative *Lycopersicon pennellii*. *Mol Plant-Microbe Interact* 2000;13:1346–55.
- Axtell MJ, Staskawicz BJ. Initiation of RPS2-specified disease resistance in *Arabidopsis* is coupled to the AvrRpt2-directed elimination of RIN4. *Cell* 2003;112:369–77.
- Badel JL, Charkowski AO, Deng W-L, Collmer A. A Gene in the *Pseudomonas syringae* pv. *tomato* Hrp pathogenicity island conserved effector locus, *hopPtoA1*, contributes to efficient formation of bacterial colonies in planta and is duplicated elsewhere in the genome. *Mol Plant-Microbe Interact* 2002;15:1014–24.
- Bai J, Choi SH, Ponciano G, Leung H, Leach JE. *Xanthomonas oryzae* pv. *oryzae* avirulence genes contribute differently and specifically to pathogen aggressiveness. *Mol Plant-Microbe Interact* 2000;13:1322–9.
- Ballvora A, Pierre M, Van den Ackerveken G, Schornack S, Rossier O, Ganai M, et al. Genetic mapping and functional analysis of the tomato *Bs4* locus, governing recognition of the *Xanthomonas campestris* pv. *vesicatoria* AvrBs4 protein. *Mol Plant-Microbe Interact* 2001;14:629–38.
- Bell KS, Sebahia M, Pritchard L, Holden MT, Hyman LJ, Holeva MC, et al. Genome sequence of the enterobacterial phytopathogen *Erwinia carotovora* subsp. *atroseptica* and characterization of virulence factors. *Proc Natl Acad Sci USA* 2004;101:11105–10.
- Blair MW, Garris AJ, Iyer AS, Chapman B, Kresovich S, McCouch SR. High resolution genetic mapping and



- candidate gene identification at the *xa5* locus for bacterial blight resistance in rice (*Oryza sativa* L.). *Theor Appl Genet* 2003;107:62–73.
- Bogdanove AJ, Martin GB. AvrPto-dependent Pto-interacting proteins and AvrPto-interacting proteins in tomato. *Proc Natl Acad Sci USA* 2000;97:8836–40.
- Bogdanove AJ, Beer SV, Bonas U, Boucher CA, Collmer A, Coplin DL, et al. Unified nomenclature for broadly conserved *hrp* genes of phytopathogenic bacteria. *Mol Microbiol* 1996;20:681–3.
- Bogdanove AJ, Bauer DW, Beer SV. *Erwinia amylovora* secretes DspE, a pathogenicity factor and functional AvrE homolog, through the *hrp* (type III secretion) pathway. *J Bacteriol* 1998;180:2244–7.
- Bonas U, Stall RE, Staskawicz B. Genetic and structural characterization of the avirulence gene *avrBs3* from *Xanthomonas campestris* pv. *vesicatoria*. *Mol Gen Genet* 1989;218:127–36.
- Bonas U, Schulte R, Fenselau S, Minsavage GV, Staskawicz BJ, Stall RE. Isolation of a gene-cluster from *Xanthomonas campestris* pv. *vesicatoria* that determines pathogenicity and the hypersensitive response on pepper and tomato. *Mol Plant-Microbe Interact* 1991;4:81–8.
- Bonas U, Conrads-Strauch J, Balbo I. Resistance in tomato to *Xanthomonas campestris* pv. *vesicatoria* is determined by alleles of the pepper-specific avirulence gene *avrBs3*. *Mol Gen Genet* 1993;238:261–9.
- Bonshtien A, Lev A, Gibly A, Debbie P, Avni A, Sessa G. Molecular properties of the *Xanthomonas* AvrRxv effector and global transcriptional changes determined by its expression in resistant tomato plants. *Mol Plant-Microbe Interact* 2005;18:300–10.
- Boyd AP, Lambermont I, Cornelis GR. Competition between the Yops of *Yersinia enterocolitica* for delivery into eukaryotic cells: role of the SycE chaperone binding domain of YopE. *J Bacteriol* 2000;182:4811–21.
- Bretz J, Losada L, Lisboa K, Hutcheson SW. Lon protease functions as a negative regulator of type III protein secretion in *Pseudomonas syringae*. *Mol Microbiol* 2002;45:397–409.
- Brito B, Aldon D, Barberis P, Boucher C, Genin S. A signal transfer system through three compartments transduces the plant cell contact-dependent signal controlling *Ralstonia solanacearum* *hrp* genes. *Mol Plant-Microbe Interact* 2002;15:109–19.
- Buell CR, Joardar V, Lindeberg M, Selengut J, Paulsen IT, Gwinn ML, et al. The complete genome sequence of the *Arabidopsis* and tomato pathogen *Pseudomonas syringae* pv. *tomato* DC3000. *Proc Natl Acad Sci USA* 2003;100:10181–6.
- Büttner D, Bonas U. Port of entry—the type III secretion translocon. *Trends Microbiol* 2002;10:186–92.
- Büttner D, Bonas U. Common infection strategies of plant and animal pathogenic bacteria. *Curr Opin Plant Biol* 2003;6:312–9.
- Büttner D, Nennstiel D, Klüsener B, Bonas U. Functional analysis of HrpF, a putative type III translocon protein from *Xanthomonas campestris* pv. *vesicatoria*. *J Bacteriol* 2002;184:2389–98.
- Büttner D, Noël L, Thieme F, Bonas U. Genomic approaches in *Xanthomonas campestris* pv. *vesicatoria* allow fishing for virulence genes. *J Biotechnol* 2003;106:203–14.
- Büttner D, Gürlebeck D, Noël LD, Bonas U. HpaB from *Xanthomonas campestris* pv. *vesicatoria* acts as an exit control protein in type III-dependent protein secretion. *Mol Microbiol* 2004;54:755–68.
- Casper-Lindley C, Dahlbeck D, Clark ET, Staskawicz BJ. Direct biochemical evidence for type III secretion-dependent translocation of the AvrBs2 effector protein into plant cells. *Proc Natl Acad Sci USA* 2002;99:8336–41.
- Chang JH, Rathjen JP, Bernal AJ, Staskawicz BJ, Michelmore RW. AvrPto enhances growth and necrosis caused by *Pseudomonas syringae* pv. *tomato* in tomato lines lacking either Pto or Prf. *Mol Plant-Microbe Interact* 2000;13:568–71.
- Chang JH, Urbach JM, Law TF, Arnold LW, Hu A, Gombar S, et al. A high-throughput, near-saturating screen for type III effector genes from *Pseudomonas syringae*. *Proc Natl Acad Sci USA* 2005;102:2549–54.
- Chatterjee A, Cui Y, Chatterjee AK. Regulation of *Erwinia carotovora* *hrpL*(Ecc) (sigma-L(Ecc)), which encodes an extracytoplasmic function subfamily of sigma factor required for expression of the HRP regulon. *Mol Plant-Microbe Interact* 2002;15:971–80.
- Chatterjee A, Cui Y, Yang H, Collmer A, Alfano JR, Chatterjee AK. GacA, the response regulator of a two-component system, acts as a master regulator in *Pseudomonas syringae* pv. *tomato* DC3000 by controlling regulatory RNA, transcriptional activators, and alternate sigma factors. *Mol Plant-Microbe Interact* 2003;16:1106–17.
- Chen Z, Kloek AP, Boch J, Katagiri F, Kunkel BN. The *Pseudomonas syringae* *avrRpt2* gene product promotes pathogen virulence from inside plant cells. *Mol Plant-Microbe Interact* 2000;13:1312–21.
- Ciesiolka LD, Hwin T, Gearlds JD, Minsavage GV, Saenz R, Bravo M, et al. Regulation of expression of avirulence gene *avrRxv* and identification of a family of host interaction factors by sequence analysis of *avrBsT*. *Mol Plant-Microbe Interact* 1999;12:35–44.
- Coaker G, Falick A, Staskawicz B. Activation of a phytopathogenic bacterial effector protein by a eukaryotic cyclophilin. *Science* 2005;308:548–50.
- Cornelis G, Van Gijsegem F. Assembly and function of type III secretory systems. *Annu Rev Microbiol* 2000;54:735–74.
- Cunnac S, Boucher C, Genin S. Characterization of the *cis*-acting regulatory element controlling HrpB-mediated activation of the type III secretion system and effector genes in *Ralstonia solanacearum*. *J Bacteriol* 2004a;186:2309–18.
- Cunnac S, Occhialini A, Barberis P, Boucher C, Genin S. Inventory and functional analysis of the large Hrp regulon in *Ralstonia solanacearum*: identification of

- novel effector proteins translocated to plant host cells through the type III secretion system. *Mol Microbiol* 2004b;53:115–28.
- Dangl JL, Jones JDG. Plant pathogens and integrated defence responses to infection. *Nature* 2001;411:826–33.
- da Silva AC, Ferro JA, Reinach FC, Farah CS, Furlan LR, Quaggio RB, et al. Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature* 2002;417:459–63.
- Deng WL, Rehm AH, Charkowski AO, Rojas CM, Collmer A. *Pseudomonas syringae* exchangeable effector loci: sequence diversity in representative pathovars and virulence function in *P. syringae* pv. *syringae* B728a. *J Bacteriol* 2003;185:2592–602.
- Deslandes L, Pileur F, Liaubet L, Camut S, Can C, Williams K, et al. Genetic characterization of *RRS1*, a recessive locus in *Arabidopsis thaliana* that confers resistance to the bacterial soilborne pathogen *Ralstonia solanacearum*. *Mol Plant-Microbe Interact* 1998;11:659–67.
- Deslandes L, Olivier J, Peeters N, Feng DX, Khounlotham M, Boucher C, et al. Physical interaction between *RRS1-R*, a protein conferring resistance to bacterial wilt, and *PopP2*, a type III effector targeted to the plant nucleus. *Proc Natl Acad Sci USA* 2003;100:8024–9.
- Dobrindt U, Hochhut B, Hentschel U, Hacker J. Genomic islands in pathogenic and environmental microorganisms. *Nat Rev Microbiol* 2004;2:414–24.
- Escobar L, van den Ackerveken G, Rossier O, Pieplow S, Bonas U. Type III secretion and *in planta* recognition of the *Xanthomonas* avirulence proteins *AvrBs1* and *AvrBsT*. *Mol Plant Pathol* 2001;2:287–96.
- Feldman MF, Cornelis GR. The multitasking type III chaperones: all you can do with 15 kDa. *FEMS Microbiol Lett* 2003;219:151–8.
- Fenselau S, Bonas U. Sequence and expression analysis of the *hrpB* pathogenicity operon of *Xanthomonas campestris* pv. *vesicatoria* which encodes eight proteins with similarity to components of the *Hrp*, *Ysc*, *Spa*, and *Fli* secretion systems. *Mol Plant-Microbe Interact* 1995;8:845–54.
- Fenselau S, Balbo I, Bonas U. Determinants of pathogenicity in *Xanthomonas campestris* pv. *vesicatoria* are related to proteins involved in secretion in bacterial pathogens of animals. *Mol Plant-Microbe Interact* 1992;5:390–6.
- Flor HH. Current status of the gene-for-gene concept. *Annu Rev Phytopathol* 1971;9:275–96.
- Francis M, Wolf-Watz H, Forsberg Å. Regulation of type III secretion systems. *Curr Opin Microbiol* 2002;5:166–72.
- Frederick RD, Majerczak DR, Coplin DL. *Erwinia stewartii* *WtsA*, a positive regulator of pathogenicity gene expression, is similar to *Pseudomonas syringae* pv. *paseolicola* *HrpS*. *Mol Microbiol* 1993;9:477–85.
- Gabriel DW. The *Xanthomonas avr/pth* gene family. Stacey G, Keen NT, editors. *Plant-microbe interactions*, vol. 4. St. Paul, MN: APS Press; 1999. p. 39–55.
- Gabriel DW, Rolfe BG. Working models of specific recognition in plant-microbe interactions. *Annu Rev Phytopathol* 1990;28:365–91.
- Galyov EE, Håkansson S, Wolf-Watz H. Characterization of the operon encoding the *YpkA Ser/Thr* protein kinase and the *YopJ* protein of *Yersinia pseudotuberculosis*. *J Bacteriol* 1994;176:4543–8.
- Gassmann W, Dahlbeck D, Chesnokova O, Minsavage GV, Jones JB, Staskawicz BJ. Molecular evolution of virulence in natural field strains of *Xanthomonas campestris* pv. *vesicatoria*. *J Bacteriol* 2000;182:7053–9.
- Genin S, Gough CL, Zischek C, Boucher CA. Evidence that the *hrpB* gene encodes a positive regulator of pathogenicity genes from *Pseudomonas solanacearum*. *Mol Microbiol* 1992;6:3065–76.
- Görlich D, Prehn S, Laskey RA, Hartmann E. Isolation of a protein that is essential for the first step of nuclear protein import. *Cell* 1994;79:767–78.
- Görlich D, Vogel F, Mills AD, Hartmann E, Laskey RA. Distinct functions for the two importin subunits in nuclear protein import. *Nature* 1995;377:246–8.
- Grimm C, Panopoulos NJ. The predicted protein product of a pathogenicity locus from *Pseudomonas syringae* pv. *phaseolicola* is homologous to a highly conserved domain of several prokaryotic regulatory proteins. *J Bacteriol* 1989;171:5031–8.
- Grimm C, Aufsatz W, Panopoulos NJ. The *hrpRS* locus of *Pseudomonas syringae* pv. *phaseolicola* constitutes a complex regulatory unit. *Mol Microbiol* 1995;15:155–65.
- Gu K, Tian D, Yang F, Wu L, Sreekala C, Wang D, et al. High-resolution genetic mapping of *Xa27(t)*, a new bacterial blight resistance gene in rice, *Oryza sativa* L. *Theor Appl Genet* 2004;108:800–7.
- Gu K, Yang B, Tian D, Wu L, Wang D, Sreekala C, et al. *R* gene expression induced by a type-III effector triggers disease resistance in rice. *Nature* 2005;435:1122–5.
- Guo M, Manulis S, Mor H, Barash I. The presence of diverse IS elements and an *avrPphD* homologue that acts as a virulence factor on the pathogenicity plasmid of *Erwinia herbicola* pv. *gypsophylae*. *Mol Plant-Microbe Interact* 2002;15:709–16.
- Gürlebeck D, Szurek B, Bonas U. Dimerization of the bacterial effector protein *AvrBs3* in the plant cell cytoplasm prior to nuclear import. *Plant J* 2005;42:175–87.
- Guttman DS, Vinatzer BA, Sarkar SF, Ranall MV, Kettler G, Greenberg JT. A functional screen for the type III (*Hrp*) secretome of the plant pathogen *Pseudomonas syringae*. *Science* 2002;295:1722–6.
- Guttman DS, Greenberg JT. Functional analysis of the type III effectors *AvrRpt2* and *AvrRpm1* of *Pseudomonas syringae* with the use of a single-copy genomic integration system. *Mol Plant-Microbe Interact* 2001;14:145–55.
- Heesemann J, Algermissen B, Laufs R. Genetically manipulated virulence of *Yersinia enterocolitica*. *Infect Immun* 1984;46:105–10.
- Herbers K, Conrads-Strauch J, Bonas U. Race-specificity of plant resistance to bacterial spot disease determined by repetitive motifs in a bacterial avirulence protein. *Nature* 1992;356:172–4.

- Hochstrasser M. Biochemistry: all in the ubiquitin family. *Science* 2000a;289:563–4.
- Hochstrasser M. Evolution and function of ubiquitin-like protein-conjugation systems. *Nat Cell Biol* 2000b;2: E153–7.
- Holeva MC, Bell KS, Hyman LJ, Avrova AO, Whisson SC, Birch PR, et al. Use of a pooled transposon mutation grid to demonstrate roles in disease development for *Erwinia carotovora* subsp. *atroseptica* putative type III secreted effector (DspE/A) and helper (HrpN) proteins. *Mol Plant-Microbe Interact* 2004;17:943–50.
- Holland IB. Translocation of bacterial proteins—an overview. *Biochim Biophys Acta* 2004;1694:5–16.
- Hotson A, Mudgett MB. Cysteine proteases in phytopathogenic bacteria: identification of plant targets and activation of innate immunity. *Curr Opin Plant Biol* 2004;7:384–90.
- Hotson A, Chosed R, Shu H, Orth K, Mudgett MB. *Xanthomonas* type III effector XopD targets SUMO-conjugated proteins *in planta*. *Mol Microbiol* 2003; 50:377–89.
- Hueck CJ. Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol Mol Biol Rev* 1998;62:379–433.
- Huguet E, Hahn K, Wengelnik K, Bonas U. *hpaA* mutants of *Xanthomonas campestris* pv. *vesicatoria* are affected in pathogenicity but retain the ability to induce host-specific hypersensitive reaction. *Mol Microbiol* 1998;29:1379–90.
- Hutcheson SW, Bretz J, Sussan T, Jin S, Pak K. Enhancer-binding proteins HrpR and HrpS interact to regulate *hrp*-encoded type III protein secretion in *Pseudomonas syringae* strains. *J Bacteriol* 2001;183: 5589–98.
- Ishihara H, Ponciano G, Leach JE, Tsuyumu S. Functional analysis of the 3' end of *avrBs3/pthA* genes from two *Xanthomonas* species. *Physiol Mol Plant Pathol* 2003;63:329–38.
- Ishihara H, Ponciano G, Leach JE, Tsuyumu S. Functional analysis of the 3' end of *avrBs3/pthA* genes from two *Xanthomonas* species. *Physiol Mol Plant Pathol* 2004;63:329–38.
- Iyer AS, McCouch SR. The rice bacterial blight resistance gene *xa5* encodes a novel form of disease resistance. *Mol Plant-Microbe Interact* 2004;17:1348–54.
- Jackson RW, Athanassopoulos E, Tsiamis G, Mansfield JW, Sesma A, Arnold DL, et al. Identification of a pathogenicity island, which contains genes for virulence and avirulence, on a large native plasmid in the bean pathogen *Pseudomonas syringae* pathovar *phaseolicola*. *Proc Natl Acad Sci USA* 1999;96: 10875–80.
- Jia Y, McAdams SA, Bryan GT, Hershey HP, Valent B. Direct interaction of resistance gene and avirulence gene products confers rice blast resistance. *EMBO J* 2000;19:4004–14.
- Jones JB, Lacy GH, Bouzari H, Stall RE, Schaad NW. Reclassification of the xanthomonads associated with bacterial spot disease of tomato and pepper. *Syst Appl Microbiol* 2004;27:755–62.
- Kay S, Boch J, Bonas U. Characterization of AvrBs3-like effectors from a *Brassicaceae* pathogen reveals virulence and avirulence activities and a protein with a novel repeat architecture. *Mol Plant-Microbe Interact* 2005;18:838–48.
- Kearney B, Staskawicz BJ. Widespread distribution and fitness contribution of *Xanthomonas campestris* avirulence gene, *avrBs2*. *Nature* 1990;346:385–6.
- Keen NT. Gene-for-gene complementarity in plant-pathogen interactions. *Annu Rev Genet* 1990;24:447–63.
- Kim J-G, Park BK, Yoo C-H, Jeon E, Oh J, Hwang I. Characterization of the *Xanthomonas axonopodis* pv. *glycines* Hrp pathogenicity island. *J Bacteriol* 2003; 185:3155–66.
- Kim JG, Jeon E, Oh J, Moon JS, Hwang I. Mutational analysis of *Xanthomonas* harpin HpaG identifies a key functional region that elicits the hypersensitive response in nonhost plants. *J Bacteriol* 2004;186: 6239–47.
- Kim HS, Desveaux D, Singer AU, Patel P, Sondek J, Dangl JL. The *Pseudomonas syringae* effector AvrRpt2 cleaves its C-terminally acylated target, RIN4, from *Arabidopsis* membranes to block RPM1 activation. *Proc Natl Acad Sci USA* 2005a;102:6496–501.
- Kim MG, da Cunha L, McFall AJ, Belkadir Y, DebRoy S, Dangl JL, et al. Two *Pseudomonas syringae* type III effectors inhibit RIN4-regulated basal defense in *Arabidopsis*. *Cell* 2005b;121:749–59.
- Klement Z. Hypersensitivity. Mount MS, Lacy GH, editors. *Phytopathogenic prokaryotes*, vol. 2. New York: Academic Press; 1982. p. 149–77.
- Knoop V, Staskawicz B, Bonas U. Expression of the avirulence gene *avrBs3* from *Xanthomonas campestris* pv. *vesicatoria* is not under the control of *hrp* genes and is independent of plant factors. *J Bacteriol* 1991;173:7142–50.
- Koebnik R. The role of bacterial pili in protein and DNA translocation. *Trends Microbiol* 2001;9:586–90.
- Komeili A, O'Shea EK. Nuclear transport and transcription. *Curr Opin Cell Biol* 2000;12:355–60.
- Lahaye T. The *Arabidopsis* *RRS1-R* disease resistance gene—uncovering the plant's nucleus as new battlefield of plant defense? *Trends Plant Sci* 2002;7:425–7.
- Lahaye T, Bonas U. Molecular secrets of bacterial type III effector proteins. *Trends Plant Sci* 2001;6:479–85.
- Landschulz WH, Johnson PF, McKnight SL. The leucine zipper: a hypothetical structure common to a new class of DNA binding proteins. *Science* 1988;240: 1759–64.
- Lavie M, Shillington E, Eguiluz C, Grimsley N, Boucher C. PopP1, a new member of the YopJ/AvrRxv family of type III effector proteins, acts as a host-specificity factor and modulates aggressiveness of *Ralstonia solanacearum*. *Mol Plant-Microbe Interact* 2002;15: 1058–68.
- Lee BM, Park YJ, Park DS, Kang HW, Kim JG, Song ES, et al. The genome sequence of *Xanthomonas oryzae* pathovar *oryzae* KACC10331, the bacterial blight pathogen of rice. *Nucleic Acids Res* 2005;33: 577–86.

- Lim MT, Kunkel BN. The *Pseudomonas syringae* *avrRpt2* gene contributes to virulence on tomato. *Mol Plant-Microbe Interact* 2005;18:626–33.
- Lin N-C, Martin GB. An *avrPto/avrPtoB* mutant of *Pseudomonas syringae* pv. *tomato* DC3000 does not elicit Pto-mediated resistance and is less virulent on tomato. *Mol Plant-Microbe Interact* 2005;18:43–51.
- Lindeberg M, Stavrinides J, Chang JH, Alfano JR, Collmer A, Dangl JL, et al. Proposed guidelines for a unified nomenclature and phylogenetic analysis of type III Hop effector proteins in the plant pathogen *Pseudomonas syringae*. *Mol Plant-Microbe Interact* 2005;18:275–82.
- Lindgren PB. The role of *hrp* genes during plant-bacterial interactions. *Annu Rev Phytopathol* 1997;35:129–52.
- Lindgren PB, Peet RC, Panopoulos NJ. Gene-cluster of *Pseudomonas syringae* pv. *phaseolicola* controls pathogenicity of bean plants and hypersensitivity on nonhost plants. *J Bacteriol* 1986;168:512–22.
- Lloyd SA, Forsberg Å, Wolf-Watz H, Francis MS. Targeting exported substrates to the *Yersinia* TTSS: different functions for different signals? *Trends Microbiol* 2001a;9:367–71.
- Lloyd SA, Norman M, Rosqvist R, Wolf-Watz H. *Yersinia* YopE is targeted for type III secretion by N-terminal, not mRNA, signals. *Mol Microbiol* 2001b;39:520–32.
- Lloyd SA, Sjöstrom M, Andersson S, Wolf-Watz H. Molecular characterization of type III secretion signals via analysis of synthetic N-terminal amino acid sequences. *Mol Microbiol* 2002;43:51–9.
- Lopez-Solanilla E, Bronstein PA, Schneider AR, Collmer A. HopPtoN is a *Pseudomonas syringae* Hrp (type III secretion system) cysteine protease effector that suppresses pathogen-induced necrosis associated with both compatible and incompatible plant interactions. *Mol Microbiol* 2004;54:353–65.
- Lorang JM, Shen H, Kobayashi D, Cooksey D, Keen NT. *avrA* and *avrE* in *Pseudomonas syringae* pv. *tomato* PT23 play a role in virulence on tomato plants. *Mol Plant-Microbe Interact* 1994;7:508–15.
- Losada L, Sussan T, Pak K, Zeyad S, Rozenbaum I, Hutcheson SW. Identification of a novel *Pseudomonas syringae* Psy61 effector with virulence and avirulence functions by a HrpL-dependent promoter-trap assay. *Mol Plant-Microbe Interact* 2004;17:254–62.
- Mackey D, Belkhadir Y, Alonso JM, Ecker JR, Dangl JL. *Arabidopsis* RIN4 is a target of the type III virulence effector AvrRpt2 and modulates RPS2-mediated resistance. *Cell* 2003;112:379–89.
- Mackey D, Holt BF, Wiig A, Dangl JL. RIN4 interacts with *Pseudomonas syringae* type III effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. *Cell* 2002;108:743–54.
- Marenda M, Brito B, Callard D, Genin S, Barberis P, Boucher C, et al. PrhA controls a novel regulatory pathway required for the specific induction of genes in the presence of plant cells. *Mol Microbiol* 1998;27:437–53.
- Marois E, Van den Ackerveken G, Bonas U. The *Xanthomonas* type III effector protein AvrBs3 modulates plant gene expression and induces cell hypertrophy in the susceptible host. *Mol Plant-Microbe Interact* 2002;15:637–46.
- Martin GB, Bogdanove AJ, Sessa G. Understanding the functions of plant disease resistance proteins. *Annu Rev Plant Biol* 2003;54:23–61.
- Mattaj JW, Englmeier L. Nucleocytoplasmic transport: the soluble phase. *Annu Rev Biochem* 1998;67:265–306.
- Merighi M, Majerczak DR, Stover EH, Coplin DL. The HrpX/HrpY two-component system activates *hrpS* expression, the first step in the regulatory cascade controlling the Hrp regulon in *Pantoea stewartii* subsp. *stewartii*. *Mol Plant-Microbe Interact* 2003;16:238–48.
- Meshi T, Iwabuchi M. Plant transcription factors. *Plant Cell Physiol* 1995;36:1405–20.
- Metz M, Dahlbeck D, Morales CQ, Al Sady B, Clark ET, Staskawicz BJ. The conserved *Xanthomonas campestris* pv. *vesicatoria* effector protein XopX is a virulence factor and suppresses host defense in *Nicotiana benthamiana*. *Plant J* 2005;41:801–14.
- Mor H, Manulis S, Zuck M, Nizan R, Coplin DL, Barash I. Genetic organization of the *hrp* gene cluster and *dspAE/BF* operon in *Erwinia herbicola* pv. *gypsophylae*. *Mol Plant-Microbe Interact* 2001;14:431–6.
- Morales CQ, Posada J, Macneale E, Franklin D, Rivas I, Bravo M, et al. Functional analysis of the early chlorosis factor gene. *Mol Plant-Microbe Interact* 2005;18:477–86.
- Mudgett MB, Staskawicz BJ. Characterization of the *Pseudomonas syringae* pv. *tomato* AvrRpt2 protein: demonstration of secretion and processing during bacterial pathogenesis. *Mol Microbiol* 1999;32:927–41.
- Mudgett MB, Chesnokova O, Dahlbeck D, Clark ET, Rossier O, Bonas U, et al. Molecular signals required for type III secretion and translocation of the *Xanthomonas campestris* AvrBs2 protein to pepper plants. *Proc Natl Acad Sci USA* 2000;97:13324–9.
- Niepold F, Anderson D, Mills D. Cloning determinants of pathogenesis from *Pseudomonas syringae* pathovar *syringae*. *Proc Natl Acad Sci USA* 1985;82:406–10.
- Nizan-Koren R, Manulis S, Mor H, Iraki NM, Barash I. The regulatory cascade that activates the Hrp regulon in *Erwinia herbicola* pv. *gypsophylae*. *Mol Plant-Microbe Interact* 2003;16:249–60.
- Noël L, Thieme F, Nennstiel D, Bonas U. cDNA-AFLP analysis unravels a genome-wide *hrpG*-regulon in the plant pathogen *Xanthomonas campestris* pv. *vesicatoria*. *Mol Microbiol* 2001;41:1271–81.
- Noël L, Thieme F, Nennstiel D, Bonas U. Two novel type III-secreted proteins of *Xanthomonas campestris* pv. *vesicatoria* are encoded within the *hrp* pathogenicity island. *J Bacteriol* 2002;184:1340–8.
- Noël L, Thieme F, Gäbler J, Büttner D, Bonas U. XopC and XopJ, two novel type III effector proteins from *Xanthomonas campestris* pv. *vesicatoria*. *J Bacteriol* 2003;185:7092–102.
- Ochiai H, Inoue Y, Takeya M, Sasaki A, Kaku H. Genome sequence of *Xanthomonas oryzae* pv. *oryzae* suggests contribution of large numbers of effector genes and

- insertion sequences to its race diversity. *Jpn Agric Res Q* 2005;39:275–87.
- Orth K. Function of the *Yersinia* effector YopJ. *Curr Opin Microbiol* 2002;5:38–43.
- Orth K, Palmer LE, Bao ZQ, Stewart S, Rudolph AE, Bliska JB, et al. Inhibition of the mitogen-activated protein kinase kinase superfamily by a *Yersinia* effector. *Science* 1999;285:1920–3.
- Orth K, Xu Z, Mudgett MB, Bao ZQ, Palmer LE, Bliska JB, et al. Disruption of signaling by *Yersinia* effector YopJ, ubiquitin-like protein protease. *Science* 2000;290:1594–7.
- Page AL, Parsot C. Chaperones of the type III secretion pathway: jacks of all trades. *Mol Microbiol* 2002;46:1–11.
- Parkinson JS, Kofoed EC. Communication modules in bacterial signaling proteins. *Annu Rev Genet* 1992;26:71–112.
- Parsot C, Hamiaux C, Page AL. The various and varying roles of specific chaperones in type III secretion systems. *Curr Opin Microbiol* 2003;6:7–14.
- Petnicki-Ocwieja T, Schneider DJ, Tam VC, Chancey ST, Shan L, Jamir Y, et al. Genomewide identification of proteins secreted by the Hrp type III protein secretion system of *Pseudomonas syringae* pv. *tomato* DC3000. *Proc Natl Acad Sci USA* 2002;99:7652–7.
- Preston G, Deng WL, Huang HC, Collmer A. Negative regulation of *hrp* genes in *Pseudomonas syringae* by HrpV. *J Bacteriol* 1998;180:4532–7.
- Ritter C, Dangl JL. The *avrRpm1* gene of *Pseudomonas syringae* pv. *maculicola* is required for virulence on *Arabidopsis*. *Mol Plant-Microbe Interact* 1995;8:444–53.
- Roden J, Eardley L, Hotson A, Cao Y, Mudgett MB. Characterization of the *Xanthomonas* AvrXv4 effector, a SUMO protease translocated into plant cells. *Mol Plant-Microbe Interact* 2004a;17:633–43.
- Roden JA, Belt B, Ross JB, Tachibana T, Vargas J, Mudgett MB. A genetic screen to isolate type III effectors translocated into pepper cells during *Xanthomonas* infection. *Proc Natl Acad Sci USA* 2004b;101:16624–9.
- Rohmer L, Guttman DS, Dangl JL. Diverse evolutionary mechanisms shape the type III effector virulence factor repertoire in the plant pathogen *Pseudomonas syringae*. *Genetics* 2004;167:1341–60.
- Ronald PC, Staskawicz BJ. The avirulence gene *avrBs1* from *Xanthomonas campestris* pv. *vesicatoria* encodes a 50-kD protein. *Mol Plant-Microbe Interact* 1988;1:191–8.
- Rossier O, Wengelnik K, Hahn K, Bonas U. The *Xanthomonas* Hrp type III system secretes proteins from plant and mammalian pathogens. *Proc Natl Acad Sci USA* 1999;96:9368–73.
- Rossier O, Van den Ackerveken G, Bonas U. HrpB2 and HrpF from *Xanthomonas* are type III-secreted proteins and essential for pathogenicity and recognition by the host plant. *Mol Microbiol* 2000;38:828–38.
- Salanoubat M, Genin S, Artiguenave F, Gouzy J, Mangenot S, Arlat M, et al. Genome sequence of the plant pathogen *Ralstonia solanacearum*. *Nature* 2002;415:497–502.
- Schechter LM, Roberts KA, Jamir Y, Alfano JR, Collmer A. *Pseudomonas syringae* type III secretion system targeting signals and novel effectors studied with a Cya translocation reporter. *J Bacteriol* 2004;186:543–55.
- Schesser K, Frithz-Lindsten E, Wolf-Watz H. Delineation and mutational analysis of the *Yersinia pseudotuberculosis* YopE domains which mediate translocation across bacterial and eukaryotic cellular membranes. *J Bacteriol* 1996;178:7227–33.
- Schorneck S, Ballvora A, Gürlebeck D, Peart J, Baulcombe D, Baker B, et al. The tomato resistance protein Bs4 is a predicted non-nuclear TIR-NB-LRR protein that mediates defense responses to severely truncated derivatives of AvrBs4 and overexpressed AvrBs3. *Plant J* 2004;37:46–60.
- Schorneck S, Meyer A, Römer P, Jordan T, Lahaye T. Gene-for-Gene mediated recognition of nuclear-targeted AvrBs3-like bacterial effector proteins. *J Plant Physiol* 2006;163:233–55.
- Schulte R, Bonas U. Expression of the *Xanthomonas campestris* pv. *vesicatoria* *hrp* gene cluster, which determines pathogenicity and hypersensitivity on pepper and tomato, is plant inducible. *J Bacteriol* 1992a;174:815–23.
- Schulte R, Bonas U. A *Xanthomonas* pathogenicity locus is induced by sucrose and sulfur-containing amino-acids. *Plant Cell* 1992b;4:79–86.
- Scofield SR, Tobias CM, Rathjen JP, Chang JH, Lavelle DT, Michelmore RW, et al. Molecular basis of gene-for-gene specificity in bacterial speck disease of tomato. *Science* 1996;274:2063–5.
- Shan L, He P, Zhou JM, Tang X. A cluster of mutations disrupt the avirulence but not the virulence function of AvrPto. *Mol Plant-Microbe Interact* 2000;13:592–8.
- Shao F, Merritt PM, Bao Z, Innes RW, Dixon JE. A *Yersinia* effector and a *Pseudomonas* avirulence protein define a family of cysteine proteases functioning in bacterial pathogenesis. *Cell* 2002;109:575–88.
- Shao F, Golstein C, Ade J, Stoutemyer M, Dixon JE, Innes RW. Cleavage of *Arabidopsis* PBS1 by a bacterial type III effector. *Science* 2003;301:1230–3.
- Sorg JA, Miller NC, Schneewind O. Substrate recognition of type III secretion machines -testing the RNA signal hypothesis. *Cell Microbiol* 2005;7:1217–25.
- Sory MP, Cornelis GR. Translocation of a hybrid YopE-adenylate cyclase from *Yersinia enterocolitica* into HeLa cells. *Mol Microbiol* 1994;14:583–94.
- Sory MP, Boland A, Lambermont I, Cornelis GR. Identification of the YopE and YopH domains required for secretion and internalization into the cytosol of macrophages, using the *cyaA* gene fusion approach. *Proc Natl Acad Sci USA* 1995;92:11998–2002.
- Stall RE. *Xanthomonas campestris* pv. *vesicatoria*. US Singh RPS, Kohmoto K, editors. Pathogenesis and host-parasite specificity in plant diseases. Prokaryotes, vol. I. Tarrytown, NY: Pergamon, Elsevier Science Inc.; 1995. p. 167–84.
- Staskawicz BJ. Genetics of plant-pathogen interactions specifying plant disease resistance. *Plant Physiol* 2001;125:73–6.

- Staskawicz BJ, Dahlbeck D, Keen NT. Cloned avirulence gene of *Pseudomonas syringae* pv. *glycinea* determines race-specific incompatibility on *Glycine max* (L.) Merr. Proc Natl Acad Sci USA 1984;81:6024–8.
- Stebbins CE, Galán JE. Maintenance of an unfolded polypeptide by a cognate chaperone in bacterial type III secretion. Nature 2001;414:77–81.
- Sugio A, Yang B, White FF. Characterization of the *hrpF* pathogenicity peninsula of *Xanthomonas oryzae* pv. *oryzae*. Mol Plant-Microbe Interact 2005;18:546–54.
- Swarup S, De Feyter R, Brlansky RH, Gabriel DW. A pathogenicity locus from *Xanthomonas citri* enables strains from several pathovars of *Xanthomonas campestris* to elicit canker-like lesions on citrus. Phytopathology 1991;81:802–9.
- Swarup S, Yang YN, Kingsley MT, Gabriel DW. An *Xanthomonas citri* pathogenicity gene, *pthA*, pleiotropically encodes gratuitous avirulence on nonhosts. Mol Plant-Microbe Interact 1992;5:204–13.
- Swords KM, Dahlbeck D, Kearney B, Roy M, Staskawicz BJ. Spontaneous and induced mutations in a single open reading frame alter both virulence and avirulence in *Xanthomonas campestris* pv. *vesicatoria* *avrBs2*. J Bacteriol 1996;178:4661–9.
- Szurek B, Marois E, Bonas U, Van den Ackerveken G. Eukaryotic features of the *Xanthomonas* type III effector *AvrBs3*: protein domains involved in transcriptional activation and the interaction with nuclear import receptors from pepper. Plant J 2001;26:523–34.
- Szurek B, Rossier O, Hause G, Bonas U. Type III-dependent translocation of the *Xanthomonas* *AvrBs3* protein into the plant cell. Mol Microbiol 2002;46:13–23.
- Tampakaki AP, Fadoulglou VE, Gazi AD, Panopoulos NJ, Kokkinidis M. Conserved features of type III secretion. Cell Microbiol 2004;6:805–16.
- Tang X, Frederick RD, Zhou J, Halterman DA, Jia Y, Martin GB. Initiation of plant disease resistance by physical interaction of *AvrPto* and *Pto* kinase. Science 1996;274:2060–3.
- Thieme F, Koebnik R, Bekel T, Berger C, Boch J, Büttner D, et al. Insights into genome plasticity and pathogenicity of the plant pathogenic bacterium *Xanthomonas campestris* pv. *vesicatoria* revealed by the complete genome sequence. J Bacteriol 2005;187:7254–66.
- Thomas NA, Brett Finlay B. Establishing order for type III secretion substrates—a hierarchical process. Trends Microbiol 2003;11:398–403.
- Tsiamis G, Mansfield JW, Hockenull R, Jackson RW, Sesma A, Athanassopoulos E, et al. Cultivar-specific avirulence and virulence functions assigned to *avrPphF* in *Pseudomonas syringae* pv. *phaseolicola*, the cause of bean halo-blight disease. EMBO J 2000;19:3204–14.
- Tsuge S, Terashima S, Furutani A, Ochiai H, Oku T, Tsuno K, et al. Effects on promoter activity of base substitutions in the cis-acting regulatory element of *HrpXo* regulons in *Xanthomonas oryzae* pv. *oryzae*. J Bacteriol 2005;187:2308–14.
- Van den Ackerveken G, Marois E, Bonas U. Recognition of the bacterial avirulence protein *AvrBs3* occurs inside the host plant cell. Cell 1996;87:1307–16.
- Van der Biezen EA, Jones JDG. Plant disease-resistance proteins and the gene-for-gene concept. Trends Biochem Sci 1998;23:454–6.
- van 't Slot KAE, Knogge W. A dual role of microbial pathogen-derived effector proteins in plant disease and resistance. Crit Rev Plant Sci 2002;21:229–71.
- Vauterin L, Rademaker J, Swings J. Synopsis on the taxonomy of the genus *Xanthomonas*. Phytopathology 2000;90:677–82.
- Vera Cruz CM, Bai J, Ona I, Leung H, Nelson RJ, Mew TW, et al. Predicting durability of a disease resistance gene based on an assessment of the fitness loss and epidemiological consequences of avirulence gene mutation. Proc Natl Acad Sci USA 2000;97:13500–5.
- Wattiau P, Bernier B, Deslee P, Michiels T, Cornelis GR. Individual chaperones required for Yop secretion by *Yersinia*. Proc Natl Acad Sci USA 1994;91:10493–7.
- Wattiau P, Woestyn S, Cornelis GR. Customized secretion chaperones in pathogenic bacteria. Mol Microbiol 1996;20:255–62.
- Weber E, Ojanen-Reuhs T, Huguet E, Hause G, Romantschuk M, Korhonen TK, et al. The type III-dependent Hrp pilus is required for productive interaction of *Xanthomonas campestris* pv. *vesicatoria* with pepper host plants. J Bacteriol 2005;187:2458–68.
- Wei Z-M, Beer SV. *hrpL* activates *Erwinia amylovora* *hrp* gene transcription and is a member of the ECF subfamily of sigma factors. J Bacteriol 1995;177:6201–10.
- Wei W, Plovianich-Jones A, Deng WL, Jin QL, Collmer A, Huang HC, et al. The gene coding for the Hrp pilus structural protein is required for type III secretion of Hrp and Avr proteins in *Pseudomonas syringae* pv. *tomato*. Proc Natl Acad Sci USA 2000a;97:2247–52.
- Wei Z, Kim JF, Beer SV. Regulation of *hrp* genes and type III protein secretion in *Erwinia amylovora* by HrpX/HrpY, a novel two-component system, and HrpS. Mol Plant-Microbe Interact 2000b;13:1251–62.
- Wei CF, Deng WL, Huang HC. A chaperone-like HrpG protein acts as a suppressor of HrpV in regulation of the *Pseudomonas syringae* pv. *syringae* type III secretion system. Mol Microbiol 2005;57:520–36.
- Wengelnik K, Bonas U. HrpX, an AraC-type regulator, activates expression of five of the six loci in the *hrp* cluster of *Xanthomonas campestris* pv. *vesicatoria*. J Bacteriol 1996;178:3462–9.
- Wengelnik K, Rossier O, Bonas U. Mutations in the regulatory gene *hrpG* of *Xanthomonas campestris* pv. *vesicatoria* result in constitutive expression of all *hrp* genes. J Bacteriol 1999;181:6828–31.
- Wengelnik K, Van den Ackerveken G, Bonas U. HrpG, a key *hrp* regulatory protein of *Xanthomonas campestris* pv. *vesicatoria* is homologous to two-component response regulators. Mol Plant-Microbe Interact 1996;9:704–12.

- Whalen MC, Wang JF, Carland FM, Heiskell ME, Dahlbeck D, Minsavage GV, et al. Avirulence gene *avrRxv* from *Xanthomonas campestris* pv. *vesicatoria* specifies resistance on tomato line Hawaii 7998. *Mol Plant-Microbe Interact* 1993;6:616–27.
- Wichmann G, Bergelson J. Effector genes of *Xanthomonas axonopodis* pv. *vesicatoria* promote transmission and enhance other fitness traits in the field. *Genetics* 2004;166:693–706.
- Wright CA, Beattie GA. *Pseudomonas syringae* pv. *tomato* cells encounter inhibitory levels of water stress during the hypersensitive response of *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 2004;101:3269–74.
- Wulff-Strobel CR, Williams AW, Straley SC. LcrQ and Sych function together at the Ysc type III secretion system in *Yersinia pestis* to impose a hierarchy of secretion. *Mol Microbiol* 2002;43:411–23.
- Xiao Y, Heu S, Yi J, Lu Y, Hutcheson SW. Identification of a putative alternate sigma factor and characterization of a multicomponent regulatory cascade controlling the expression of *Pseudomonas syringae* pv. *syringae* Pss61 *hrp* and *hrmA* genes. *J Bacteriol* 1994;176:1025–36.
- Xiao Y, Hutcheson SW. A single promoter sequence recognized by a newly identified alternate sigma factor directs expression of pathogenicity and host range determinants in *Pseudomonas syringae* [published erratum appears in *J Bacteriol* 176:6158]. *J Bacteriol* 1994;176:3089–91.
- Yang B, White FF. Diverse members of the AvrBs3/PthA family of type III effectors are major virulence determinants in bacterial blight disease of rice. *Mol Plant-Microbe Interact* 2004;17:1192–200.
- Yang B, Zhu W, Johnson LB, White FF. The virulence factor AvrXa7 of *Xanthomonas oryzae* pv. *oryzae* is a type III secretion pathway-dependent nuclear-localized double-stranded DNA-binding protein. *Proc Natl Acad Sci USA* 2000;97:9807–12.
- Yang Y, Gabriel DW. Intragenic recombination of a single plant pathogen gene provides a mechanism for the evolution of new host specificities. *J Bacteriol* 1995a;177:4963–8.
- Yang Y, Gabriel DW. *Xanthomonas* avirulence/pathogenicity gene family encodes functional plant nuclear targeting signals. *Mol Plant-Microbe Interact* 1995b;8:627–31.
- Yang YN, De Feyter R, Gabriel DW. Host-specific symptoms and increased release of *Xanthomonas citri* and *X. campestris* pv. *malvacearum* from leaves are determined by the 102-bp tandem repeats of *pthA* and *avrb6*, respectively. *Mol Plant-Microbe Interact* 1994;7:345–55.
- Zhu WG, Magbanua MM, White FF. Identification of two novel *hrp*-associated genes in the *hrp* gene cluster of *Xanthomonas oryzae* pv. *oryzae*. *J Bacteriol* 2000;182:1844–53.
- Zhu WG, Yang B, Chittoor JM, Johnson LB, White FF. AvrXa10 contains an acidic transcriptional activation domain in the functionally conserved C terminus. *Mol Plant-Microbe Interact* 1998;11:824–32.