



Biomolecules: lipids

Organic biomolecules: lipids

- ◆ Organic amphiphilic compounds insoluble in water
- ◆ Easily extracted from animal and vegetal cells using apolar solvents
- ◆ Fundamental to build cell's shape and organelles
- ◆ They form a heterogeneous class, difficult to classify in a simple manner




A short list of lipid classes

- ◆ **Fatty acids (FA):** organic carboxylic acids with long carbon chains ($C > 7$), saturated, unsaturated, subjected to esterification
- ◆ **Glycerides:** mono-, di-, tri-esters of fatty acids and glycerol
- ◆ **Ceramides:** esters of sphingosine and fatty acids
- ◆ **Phospholipids:** di-esters of glycerol-3-Phosphate or esters of sphingosine-phosphate
- ◆ **Cerebrosides (glycolipids):** esters of sphingosine with 1 FA and 1 sugar with 6-C (hexose)
- ◆ **Gangliosides:** glycolipids in which the sugar moiety is complex and branched
- ◆ **Steroids:** derivatives of cholesterol (cicle-pentane-peri-hydro-phenanthrene)
- ◆ **Terpenoids:** compounds made by repetitions of isoprenes units (2-methyl-butadiene)

Some naturally occurring Fatty Acids

Carbon skeleton	Structure*	Systematic name†	Common name (derivation)	Melting point (°C)
12:0	CH ₃ (CH ₂) ₁₀ COOH	<i>n</i> -Dodecanoic acid	Lauric acid (Latin <i>laurus</i> , "laurel plant")	44.2
14:0	CH ₃ (CH ₂) ₁₂ COOH	<i>n</i> -Tetradecanoic acid	Myristic acid (Latin <i>Myristica</i> , nutmeg genus)	53.9
16:0	CH ₃ (CH ₂) ₁₄ COOH	<i>n</i> -Hexadecanoic acid	Palmitic acid (Latin <i>palma</i> , "palm tree")	63.1
18:0	CH ₃ (CH ₂) ₁₆ COOH	<i>n</i> -Octadecanoic acid	Stearic acid (Greek <i>stear</i> , "hard fat")	69.6
20:0	CH ₃ (CH ₂) ₁₈ COOH	<i>n</i> -Eicosanoic acid	Arachidic acid (Latin <i>Arachis</i> , legume genus)	76.5
24:0	CH ₃ (CH ₂) ₂₂ COOH	<i>n</i> -Tetracosanoic acid	Lignoceric acid (Latin <i>lignum</i> , "wood" + <i>cera</i> , "wax")	86.0
16:1(Δ ⁹)	CH ₃ (CH ₂) ₅ CH=CH(CH ₂) ₇ COOH	<i>cis</i> -9-Hexadecenoic acid	Palmitoleic acid	-0.5
18:1(Δ ⁹)	CH ₃ (CH ₂) ₇ CH=CH(CH ₂) ₇ COOH	<i>cis</i> -9-Octadecenoic acid	Oleic acid (Latin <i>oleum</i> , "oil")	13.4
18:2(Δ ^{9,12})	CH ₃ (CH ₂) ₄ CH=CHCH ₂ CH=CH(CH ₂) ₇ COOH	<i>cis</i> -, <i>cis</i> -9,12-Octadecadienoic acid	Linoleic acid (Greek <i>linon</i> , "flax")	-5
18:3(Δ ^{9,12,15})	CH ₃ CH ₂ CH=CHCH ₂ CH=CHCH ₂ CH=CH(CH ₂) ₇ COOH	<i>cis</i> -, <i>cis</i> -, <i>cis</i> -9,12,15-Octadecatrienoic acid	α-Linolenic acid	-11
20:4(Δ ^{5,8,11,14})	CH ₃ (CH ₂) ₄ CH=CHCH ₂ CH=CHCH ₂ CH=CHCH ₂ CH=CH(CH ₂) ₃ COOH	<i>cis</i> -, <i>cis</i> -, <i>cis</i> -, <i>cis</i> -5,8,11,14-Icosatetraenoic acid	Arachidonic acid	-49.5

The most common FAs in biological membranes



In the membrane there are always FA with an even number of C.


In case there are, double bonds are always in cis configuration.

The old notation ω -3, ω -6 and ω -9 refers to the position of the first double bond, counting how many carbon atoms are left before the end

FAs packing


- ◆ 3D packing or assembly of FAs depends on the degree of unsaturation (no. of double bonds)
- ◆ The main consequence of this packing is the melting temperature: saturated FAs are solid at RT, unsaturated FAs have lower T_m)

- ◆ FAs s... in or-
der to



FAs packing

- ◆ 3D packing or assembly of FAs depends on the degree of unsaturation (no. of double bonds)
- ◆ The main consequence of this packing is the melting temperature: saturated FAs are solid at RT, unsaturated FAs have lower T_m)
- ◆ FAs spontaneously assemble in aqueous solutions in order to decrease the hydrophobic effect of water




Which reactions do FAs undergo?

- ◆ They are weak acids with K_a $10^{-5} - 10^{-6}$, decreasing with increasing chain length
- ◆ They can form:
 - esters after reacting with alcohols
 - amides after reacting with amines
 - salts after reacting with a strong base (NaOH / KOH)
- ◆ They can be oxidised by enzymes to release energy (beta-oxidations)



Which reactions do FAs undergo?

- ◆ Unsaturated FAs can also be oxidised by oxygen radicals, giving rise to epoxides, di-oles, peroxides
- ◆ These byproducts are degraded into aldehydes and short chain FA, and are responsible for the bad smell/taste of out-of-date fat



Mono-, di- and tri-glycerides

- ◆ Esters of FA(s) with glycerol (1,2,3-propan-tri-ole)
- ◆ Triglycerides are neutral lipids used for storage, they are confined in specialised vacuoles of special cells (adipose cells)




1-stearoyl-2-linoleyl-3-palmitoyl-glycerol

Soap production


- ◆ Soap is the sodium / potassium salt of long chain FAs
- ◆ It is produced from either animal or vegetal fat
- ◆ The reaction is an alkaline hydrolysis performed at high temperature
- ◆ This reaction produces 1 mol of glycerol and 3 mol of salt per 1 mol of triglyceride
- ◆ The % of unsaturated FA and presence of additives give solid or liquid soap at room temperature



Phospholipids

- ◆ These molecules are the scaffold of the membrane lipid bilayer and contain a polar head
- ◆ The term phospholipids include two types of polar lipids present in biological membranes:
 - **Glycerol-phospholipids**, synthesized starting from glycerol-3-phosphate (G3P, an intermediate of glucose metabolism)
 - **Sphingolipids**, synthesized from sphingosine (a complex aliphatic amino-alcohol)




Common classes of glycerophospholipids




Name of X—OH	Formula of -X	Name of Phospholipid
Water	—H	Phosphatidic acid
Ethanolamine	—CH ₂ CH ₂ NH ₃ ⁺	Phosphatidylethanolamine
Choline	—CH ₂ CH ₂ N(CH ₃) ₃ ⁺	Phosphatidylcholine (lecithin)
Serine	—CH ₂ CH(NH ₃ ⁺)COO ⁻	Phosphatidylserine
<i>myo</i> -Inositol		Phosphatidylinositol
Glycerol	—CH ₂ CH(OH)CH ₂ OH	Phosphatidylglycerol
Phosphatidylglycerol		Diphosphatidylglycerol (cardiolipin)

*A typical mammalian phospholipid:
phosphatidyl-choline (PC)*




Phospholipases are enzymes specialised in the hydrolysis of single ester bonds



Sphingolipids


- ◆ These lipids account for 20-30% of plasma membrane, especially in cells of the central nervous system
- ◆ In this case the scaffold is the trans unsaturated 18-C amino alcohol called **sphingosine**
- ◆ When the amine is conjugated with a FA → ceramide is produced
- ◆ When the OH is esterified → sphingolipid (e.g. sphingomyelin, sphingolipid with neutral sugars) or sphingolipid with accharides)




Spingolipids

- ◆ These lipids account for 20-30% of plasma membrane, especially in cells of the central nervous system
- ◆ In this case the scaffold is the trans unsaturated 18-C amino alcohol called sphingosine
- ◆ When the amine is conjugated with a FA → ceramide is produced
- ◆ When the OH in position 1 of Ceramide is esterified → sev-

iced:





Spingolipids




- ◆ When the OH in position 1 of Ceramide is esterified → several sphingolipids are produced:
 - Sphingomyelin (X = phosphocholine)
 - Cerebrosides (glucose or other neutral sugars)
 - Gangliosides (complex oligosaccharides)

Glycerol-phospholipids vs sphingolipids

- ◆ Despite the different chemical origin, the two molecules have a similar 3D spatial arrangement and a similar charge (polar) distribution




Gangliosides



- ◆ Both Gangliosides and Cerebrosides contain sugars → they are always present only on the external leaflet of the membrane bilayer
- ◆ Gangliosides are responsible for the blood groups


Sterols

- ◆ This class is found almost exclusively in eukaryotes
- ◆ They have a common cyclic non-aromatic scaffold derived from cicle-pentane-peri-hydro-phenanthrene
- ◆ They have both structural and hormonal/signalling function
- ◆ The presence of 4 fused rings rigidifies the molecule → once inserted into the membranes it affects fluidity
- ◆ In mammals the precursor form all steroid hormones is cholesterol, whose OH group in C3 makes it slightly polar



Sterols

- ◆ This class is found almost exclusively in eukaryotes
- ◆ They have a common cyclic non-aromatic scaffold derived from cicle-pentane-peri-hydro-phenanthrene
- ◆ They have both structural and hormonal/signalling function
- ◆ The presence of 4 fused rings rigidifies the molecule → once inserted into the membranes it affects fluidity
- ◆ In mammals the precursor of all steroid hormones is cholesterol, whose OH group in C3 makes it slightly polar






Steroid hormones

- ◆ **Glucocorticoids** (ex. Cortisol): hormones synthesised by the surrenal cortex glands, affecting the metabolism of carbohydrates, proteins and lipids. They also affect inflammatory and stress responses.
- ◆ **Mineral-corticoids** (ex. Aldosterone): hormones synthesised by the surrenal cortex glands, affecting water and salt excretion from kidneys.
- ◆ **Androgens** (ex. Testosterone) and **estrogens** (ex. beta-estradiol): crucial hormones for a correct sexual development and functionality, they are synthesised by either testes (male) or ovaries (female)
- ◆ They are all slightly more soluble than cholesterol

Vitamin D


- ◆ Vitamin D is a regulator of Ca^{2+} homeostasis
- ◆ It is formed after photolysis of C9-C10 bond by UV light, followed by a spontaneous isomerization to vitamin D2 or D3
- ◆ Both these forms must be hydroxylated in the kidney (position C1) and in the liver (position C25) to give the active compound: 1-alpha-25-di-hydroxy-colecalciferol.



R = X vitamin D3
R = Y vitamin D2

Vitamin D

- ◆ The active form of vitamin D (1,25-dihydroxyvitamin D) induces an increase in the plasma concentration of Ca^{2+} , thus favouring its uptake by intestinal cells.
- ◆ Without vitamin D, only 10 to 15% of dietary calcium and about 60% of phosphorus is absorbed. The interaction with its receptor increases the efficiency of intestinal calcium absorption to 30 to 40% and phosphorus absorption to approximately 80%
- ◆ In this way these ions are deposited into bones and teeth.




Vitamin D

- ◆ Very low levels of vitamin D, especially in pregnant women and infants, result in low level of growth and mental retardation.
- ◆ Vitamin D deficiency also causes muscle weakness. Skeletal muscles have a vitamin D receptor and require this vitamin for maximum function.
- ◆ Brain, prostate, breast, and colon tissues, among others, as well as immune cells have a vitamin D receptor. In addition, some of these tissues and cells express the 25-hydroxyvitamin D-1 α -hydroxylase.
- ◆ Directly or indirectly, it controls more than 200 genes, including genes responsible for the regulation of cellular proliferation, differentiation, apoptosis, and angiogenesis.
- ◆ It decreases cellular proliferation of both normal cells and cancer cells and induces their terminal differentiation.
- ◆ One practical application is the use of 1,25-dihydroxyvitamin D3 and its active analogues for the treatment of psoriasis.


Terpenes and terpenoids

- ◆ Heterogenous class of hormones, co-enzymes, vitamins sharing only a part of the scaffold: a repeat of isoprene units




Coenzyme Q (ubiquinone)

- ◆ Shuttle of electrons in the respiratory chain in mitochondria, chloroplasts and in prokaryotes
- ◆ It is stably inserted in the membranes



Vitamin A (retinol)

- ◆ Vitamin A is a derivative of beta-carotene, once oxidised into the aldehyde form (retinal) it is incorporated into the proteins of vision, where it helps transducing the light impulse into images
- ◆ Low levels supplied by the diet induce a reduced by-night vision
- ◆ Extremely low levels induce blindness (poverty-related blindness)





X = CH_2OH retinol


X = CHO retinal

Vitamin K

- ◆ This vitamin is only produced by plants and bacteria
- ◆ In humans it is produced by the intestinal commensals
- ◆ It is crucial for a proper blood coagulation



Vitamin K1 (plants)



Vitamin K2 (bacteria)

Vitamin E

- ◆ Also known as alpha-tocopherol is an anti-oxidant, able to scavenge reactive oxygen species (ROS)
- ◆ It prevents oxidation of membranes and proteins

